


09-01-00

A

08/31/00



08/31/00 09:39:36 001

Please type a plus sign (+) inside this box → ☐
 Approved for use through 09/30/2000. OMB 0651-0032
 Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
 Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))

Attorney Docket No. 5594.US.P9

First Inventor or Application Identifier M. Winn et al.

Title Endothelin Antagonists

Express Mail Label No. EL 384 168 615 US

APPLICATION ELEMENTS

See MPEP chapter 900 concerning utility patent application contents.

1. ☒ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original and a duplicate for fee processing)
2. ☒ Specification [Total Pages 865]
(preferred arrangement set forth below)
- Descriptive title of the Invention
- Cross References to Related Applications
- Statement Regarding Fed sponsored R & D
- Reference to Microfiche Appendix
- Background of the Invention
- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure
3. ☐ Drawing(s) (35 U.S.C. 113) [Total Sheets ☐
4. Oath or Declaration [Total Pages ☐

- a. ☐ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 C.F.R. § 1.63(d))
(for continuation/divisional with Box 16 completed)
- i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).

* NOTE FOR ITEMS 1 & 13 IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

ADDRESS TO:

 Assistant Commissioner for Patent
 Box Patent Application
 Washington, DC 20231

5. ☐ Microfiche Computer Program (Appendix)
6. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- a. ☐ Computer Readable Copy
- b. ☐ Paper Copy (identical to computer copy)
- c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

7. ☐ Assignment Papers (cover sheet & document(s))
8. ☐ 37 C.F.R. § 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)
9. ☐ English Translation Document (if applicable)
10. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
11. ☐ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
13. ☐ * Small Entry ☐ Statement filed in prior application
Statement(s) (PTO/SB/02-12) Status still proper and desired
14. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
15. ☒ Other: Unexecuted Declaration & POA

16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No. _____

Prior application information: Examiner _____ Group / Art Unit _____

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

17. CORRESPONDENCE ADDRESS

☐ Customer Number or Bar Code Label

(Insert Customer No. or Attach bar code label here)

or ☐ Correspondence address below

Name Steven F. Weinstock
Abbott Laboratories
Department 377 / AP6D-2

Address 100 Abbott Park Road

City Abbott Park State IL Zip Code 60064-6050

Country USA Telephone (847) 937-4556 Fax (847) 938-2623

Name (Print/Type) James D. McNeil Registration No. (Attorney/Agent) 26,204

Signature *James D. McNeil* Date 08/31/2000

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: M. Winn *et al.*

Serial No.: (Not assigned yet)

Group No.: (Not assigned yet)

Filed: August 31, 2000

Examiner: (Not assigned yet)

For: **Endothelin Antagonists**

Box Patent Application

Assistant Commissioner for Patents

Washington, D.C. 20231

EXPRESS MAIL CERTIFICATE

"Express Mail" label number EL384168615US

Date of Deposit: August 31, 2000

I hereby state that the following *attached* papers or fees

Specification (791 pages); Claims (73 pages); Abstract (1 page); **Total: 865**

Utility Patent Application Transmittal

Unexecuted, Declaration and Power of Attorney

Fee Transmittal— 1 pg. in duplicate

Two Return-Receipt Postcards

are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10, on the date indicated above and is addressed to the Box Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

Deborah L. Siedlecki



Signature of person mailing paper or fee

(Express Mail Certificate)

ENDOTHELIN ANTAGONISTS

This application is a continuation-in-part application of U.S. patent application Serial No. 09/634, 661, filed August 7, 2000, which is a continuation-in-part application of U.S. patent application Serial No. 09/048,955, filed March 27, 1998, which is a continuation-in-part application of U.S. patent application Serial No. 08/794,506, filed February 4, 1997 which is a continuation-in-part of U.S. patent application Serial No. 08/600,625, filed February 13, 1996, which is a continuation-in-part of U.S. patent application Serial No. 08/497,998, filed August 2, 1995, which is a continuation-in-part of U.S. patent application Serial No. 08/442,575, filed May 30, 1995, which is a continuation-in-part of U.S. patent application Serial No. 08/334,717, filed November 4, 1994, which is a continuation-in-part of U.S. patent application Serial No. 08/293,349, filed August 19, 1994.

Technical Field

The present invention relates to compounds which are endothelin antagonists, processes for making such compounds, synthetic intermediates employed in these processes, methods and compositions for antagonizing endothelin, methods for the inhibition of bone metastases, methods for the prevention of growth of new metastases, methods for the inhibition of bone turnover, and methods for the prevention of bone loss in patients, including cancer patients, using an endothelin ET receptor antagonist.

Background of the Invention

Endothelin (ET) is a 21 amino acid peptide that is produced by endothelial cells. ET is produced by enzymatic cleavage of a Trp-Val bond in the precursor peptide big endothelin (Big ET). This cleavage is caused by an endothelin converting enzyme (ECE). Endothelin has been shown to constrict arteries and veins, increase mean arterial blood pressure, decrease cardiac output, increase cardiac contractility in vitro, stimulate mitogenesis in vascular smooth muscle cells in vitro, contract non-vascular smooth muscle including guinea pig trachea, human urinary bladder strips and rat uterus in vitro, increase airway resistance in vivo, induce formation of gastric ulcers, stimulate release of atrial natriuretic factor in vitro and in vivo, increase plasma levels of vasopressin, aldosterone and catecholamines, inhibit release of renin in vitro and stimulate release of gonadotropins in vitro.

It has been shown that vasoconstriction is caused by binding of endothelin to its receptors on vascular smooth muscle (Nature 332 411 (1988), FEBS Letters 231 440 (1988) and Biochem. Biophys. Res. Commun. 154 868 (1988)). An agent which suppresses endothelin production or an agent which binds to endothelin or which inhibits the binding of endothelin to an endothelin receptor will produce beneficial effects in a variety of therapeutic

areas. In fact, an anti-endothelin antibody has been shown, upon intrarenal infusion, to ameliorate the adverse effects of renal ischemia on renal vascular resistance and glomerular filtration rate (Kon, et al., J. Clin. Invest. 83 1762 (1989)). In addition, an anti-endothelin antibody attenuated the nephrotoxic effects of intravenously administered cyclosporin (Kon, et al., Kidney Int. 37 1487 (1990)) and attenuated infarct size in a coronary artery ligation-induced myocardial infarction model (Watanabe, et al., Nature 344 114 (1990)).

Clozel et al. (Nature 365: 759-761 (1993)) report that Ro 46-2005, a nonpeptide ET-A/B antagonist, prevents post-ischaemic renal vasoconstriction in rats, prevents the decrease in cerebral blood flow due to subarachnoid hemorrhage (SAH) in rats, and decreases MAP in sodium-depleted squirrel monkeys when dosed orally. A similar effect of a linear tripeptide-like ET-A antagonist, BQ-485, on arterial caliber after SAH has also been recently reported (S.Itoh, T. Sasaki, K. Ide, K. Ishikawa, M. Nishikibe, and M. Yano, Biochem. Biophys. Res. Comm. , 195: 969-75 (1993)). These results indicate that agents which antagonize ET/ET receptor binding will provide therapeutic benefit in the indicated disease states.


Agents with the ability to antagonize ET/ET receptor binding have been shown to be active in a number of animal models of human disease. For example, Hogaboam et al (EUR. J. Pharmacol. 1996, 309, 261-269), have shown that an endothelin receptor antagonist reduced injury in a rat model of colitis. Aktan et al (Transplant Int 1996, 9, 201-207) have demonstrated that a similar agent prevents ischemia-reperfusion injury in kidney transplantation. Similar studies have suggested the use of endothelin antagonists in the treatment of angina, pulmonary hypertension, Raynaud's disease, and migraine. (Ferro and Webb, Drugs 1996, 51,12-27).

Abnormal levels of endothelin or endothelin receptors have also been associated with a number of disease states, including prostate cancer (Nelson et al, Nature Medicine 1995, 1, 944-949) and as a modulator in osteoblastic bone lesion (UROLOGY 53:1063-1069, 1999), suggesting a role of endothelin in the pathophysiology of these diseases.

Wu-Wong et al (Life Sciences 1996, 58, 1839-1847) have shown that both endothelin and endothelin antagonists bind tightly to plasma proteins, e.g., serum albumin. This plasma protein binding can decrease the effectiveness with which the antagonists inhibit endothelin's action. Thus, endothelin antagonists with reduced plasma protein binding may be more effective than highly bound congeners.

Disclosure of the Invention

In accordance with the present invention there are compounds of the formula (I):






wherein


Z is -C(R₁₈)(R₁₉)- or -C(O)- wherein R₁₈ and R₁₉ are independently selected from hydrogen and loweralkyl;

n is 0 or 1;

R is -(CH₂)_m-W wherein m is an integer from 0 to 6 and W is

- (a) -C(O)₂-G wherein G is hydrogen or a carboxy protecting group,
- (b) -PO₃H₂,
- (c) -P(O)(OH)E wherein E is hydrogen, loweralkyl or arylalkyl,
- (d) -CN,
- (e) -C(O)NHR₁₇ wherein R₁₇ is loweralkyl,
- (f) alkylaminocarbonyl,
- (g) dialkylaminocarbonyl,
- (h) tetrazolyl,
- (i) hydroxy,
- (j) alkoxy,
- (k) sulfonamido,
- (l) -C(O)NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl, haloalkyl, aryl or dialkylamino,
- (m) -S(O)₂NHC(O)R₁₆ wherein R₁₆ is defined as above,





- R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen;
- 15 R_3 is (a) $R_4-C(O)-R_5$ -, R_4-R_{5a} -, $R_4-C(O)-R_5-N(R_6)$ -, $R_6-S(O)_2-R_7$ - or $R_{26}-S(O)-R_{27}$ - wherein R_5 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene, (iv) $-N(R_{20})-R_8$ - or $-R_{8a}-N(R_{20})-R_8$ - wherein R_8 and R_{8a} are independently selected from the group consisting of alkylene and alkenylene and R_{20} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl or cycloalkylalkyl or (v) $-O-R_9$ - or $-R_{9a}-O-R_9$ - wherein R_9 and R_{9a} are independently selected from alkylene;

R_{5a} is (i) alkylene or (ii) alkenylene;

R_7 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene or

(iv) $-N(R_{21})-R_{10}-$ or $-R_{10a}-N(R_{21})-R_{10}-$ wherein R_{10} and R_{10a} are independently selected from the group consisting of alkylene and alkenylene and R_{21} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl; R_4 and R_6 are independently selected from the group consisting of

5 (i) $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from

- (1) hydrogen,
- (2) loweralkyl,
- (3) haloalkyl,
- (4) alkoxyalkyl,
- (5) haloalkoxyalkyl,
- (6) alkenyl,
- (7) alkynyl,
- (8) cycloalkyl,
- (9) cycloalkylalkyl,
- (10) aryl,
- (11) heterocyclic,
- (12) arylalkyl,
- (13) (heterocyclic)alkyl,
- (14) hydroxyalkyl,
- (15) alkoxy,
- (16) aminoalkyl,
- (17) trialkylaminoalkyl,
- (18) alkylaminoalkyl,
- (19) dialkylaminoalkyl,
- (20) carboxyalkyl, and
- (21) diarylalkyl,

(ii) loweralkyl,

(iii) alkenyl,

(iv) alkynyl,

(v) cycloalkyl,

(vi) cycloalkylalkyl,

(vii) aryl,


(viii) arylalkyl,

(ix) heterocyclic,

(x) (heterocyclic)alkyl,

(xi) alkoxyalkyl,

- (xii) hydroxyalkyl,
- (xiii) haloalkyl,
- (xiv) haloalkenyl,
- (xv) haloalkoxyalkyl,
- (xvi) haloalkoxy,
- (xvii) alkoxyhaloalkyl,
- (xviii) alkylaminoalkyl,
- (xix) dialkylaminoalkyl,
- (xx) alkoxy,



(xxi) wherein z is 0-5 and R_{7a} is alkylene, and

(xxii) $(R_{11a})(R_{12a})N-N(H)-$

wherein R_{11a} and R_{12a} are independently selected from aryl and alkyl,

R_{26} is (i) loweralkyl, (ii) haloalkyl, (iii) alkenyl, (iv) alkynyl, (v) cycloalkyl, (vi) cycloalkylalkyl, (vii) aryl, (viii) arylalkyl, (ix) heterocyclic, (x) (heterocyclic)alkyl, (xi) alkoxyalkyl or (xii) alkoxy-substituted haloalkyl; and

R_{27} is alkylene or alkenylene;


(b) $R_{22}-O-C(O)-R_{23}-$ wherein R_{22} is a carboxy protecting group or heterocyclic and R_{23} is (i) a covalent bond, (ii) alkylene, (iii) alkenylene or (iv) $-N(R_{24})-R_{25}-$ wherein R_{25} is alkylene and R_{24} is hydrogen or loweralkyl,

- (c) loweralkyl,
- (d) alkenyl,
- (e) alkynyl,
- (f) cycloalkyl,
- (g) cycloalkylalkyl,
- (h) aryl,
- (i) arylalkyl,
- (j) aryloxyalkyl,
- (k) heterocyclic,
- (l) (heterocyclic)alkyl,
- (m) alkoxyalkyl,
- (n) alkoxyalkoxyalkyl, or

(o) $R_{13}-C(O)-CH(R_{14})-$

wherein R_{13} is amino, alkylamino or dialkylamino and R_{14} is aryl or $R_{15}-C(O)-$ wherein R_{15} is amino, alkylamino or dialkylamino; or a pharmaceutically acceptable salt thereof.

A preferred embodiment of the invention is a compound of formula (II)



wherein the substituents $-R_2$, $-R$ and $-R_1$ exist in a *trans,trans* relationship and Z , n , R , R_1 , R_2 , and R_3 are as defined above.

Another preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0 and Z is $-CH_2-$.

Another preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 1 and Z is $-CH_2-$.

Another preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-CH_2-$, and R_3 is $R_4-C(O)-R_5-$, $R_6-S(O)_2-R_7-$ or $R_{26}-S(O)-R_{27}-$ wherein R_4 , R_5 , R_6 , R_7 , R_{26} and R_{27} are as defined above.

Another preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-CH_2-$, and R_3 is alkoxyalkyl or alkoxyalkoxyalkyl.

Another preferred embodiment of the invention is a compound of formula (I) or formula (II) wherein n is zero; Z is $-CH_2-$ wherein R_{18} and R_{19} are hydrogen; R is $C(O)-G$ wherein G is hydrogen; R_1 is aryl substituted with one substituent selected from methoxy, methoxyethoxy, and isopropoxyethoxy; R_2 is 1,3-benzodiox-5-yl; R_3 is $R_4-C(O)-R_5-$ wherein R_5 is methylene and R_4 is selected from $(R_{11})(R_{12})N-$ and $(R_{11a})(R_{12a})N-N(H)-$; one of R_{11} and R_{12} is hydrogen and the other is selected from arylalkyl and diarylalkyl wherein each

aryl group of the diarylalkyl is substituted with methyl or ethyl; and one of R_{11a} or R_{12a} is alkyl and the other is aryl.

Another preferred embodiment of the invention is a compound of formula (I) or
 5 formula (II) wherein n is zero; Z is $-CH_2-$ wherein R_{18} and R_{19} are hydrogen; R is $C(O)-G$ wherein G is hydrogen; R_1 is phenyl substituted with one substituent selected from methoxy, methoxyethoxy, and isopropoxyethoxy; R_2 is 1,3-benzodiox-5-yl; R_3 is $R_4-C(O)-R_5-$ wherein
 R_5 is methylene and R_4 is selected from $(R_{11})(R_{12})N-$ and $(R_{11a})(R_{12a})N-N(H)-$; one of R_{11}
 10 and R_{12} is hydrogen and the other is selected from phenylalkyl and diphenylalkyl wherein each phenyl group of the diphenylalkyl is substituted with methyl or ethyl; and one of R_{11a} or
 R_{12a} is alkyl and the other is phenyl.

A more preferred embodiment of the invention is a compound of formula (I) or (II)
 15 wherein n is 0, Z is $-CH_2-$, and R_3 is $R_4-C(O)-R_5-$ wherein R_4 is $(R_{11})(R_{12})N-$ as defined above and R_5 is alkylene or R_3 is $R_6-S(O)_2-R_7-$ or $R_{26}-S(O)-R_{27}-$ wherein R_7 is alkylene,
 R_{27} is alkylene and R_6 and R_{26} are defined as above.

Another more preferred embodiment of the invention is a compound of formula (I) or
 20 (II) wherein n is 0, Z is $-CH_2-$ and R_3 is $R_4-C(O)-N(R_{20})-R_8-$ or $R_6-S(O)_2-N(R_{21})-R_{10}-$ wherein R_8 and R_{10} are alkylene and R_4 , R_6 , R_{20} and R_{21} are defined as above.

An even more preferred embodiment of the invention is a compound of formula (I) or
 25 (II) wherein n is 0, R is tetrazolyl or $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group or R is tetrazolyl or R is
 $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$,
 R_1 and R_2 are independently selected from (i) loweralkyl, (ii) cycloalkyl, (iii) substituted aryl
 wherein aryl is phenyl substituted with one, two or three substituents independently selected
 from loweralkyl, alkoxy, halo, alkoxyalkoxy and carboxyalkoxy, (iv) substituted or
 30 unsubstituted heterocyclic, (v) alkenyl, (vi) heterocyclic (alkyl), (vii) arylalkyl, (viii)
 aryloxyalkyl, (ix) (N-alkanoyl-N-alkyl)aminoalkyl and (x) alkylsulfonylamidoalkyl, and R_3 is
 $R_4-C(O)-R_5-$ wherein R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected
 from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl, arylalkyl, heterocyclic,
 hydroxyalkyl, alkoxy, aminoalkyl, and trialkylaminoalkyl, and R_5 is alkylene; or R_3 is
 $R_4-C(O)-N(R_{20})-R_8-$ or $R_6-S(O)_2-N(R_{21})-R_{10}-$ wherein R_4 is loweralkyl, aryl, alkoxy,
 35 alkylamino, aryloxy or arylalkoxy and R_6 is loweralkyl, haloalkyl, alkoxyalkyl,
 haloalkoxyalkyl, aryl or arylalkyl, R_8 and R_{10} are alkylene and R_{20} and R_{21} are loweralkyl; or

R_3 is $R_6-S(O)_2-R_7$ - or $R_{26}-S(O)-R_{27}$ - wherein R_6 is loweralkyl or haloalkyl, R_7 is alkylene, R_{26} is loweralkyl and R_{27} is alkylene.

- A yet more preferred embodiment of the invention is a compound of formula (I) or (II)
- 5 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i) loweralkyl, (ii) alkenyl,
- (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl,
- (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl,
- 10 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl,
- 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl,
- 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl,
- 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or
- dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and
- 15 carboxyalkoxy, (ix) heterocyclic (alkyl), (x) arylalkyl, (xi) aryloxyalkyl, (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonamidoalkyl, R_2 is substituted or unsubstituted
- 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl,
- 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl,
- 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is
- 20 $R_4-C(O)-N(R_{20})-R_8$ - or $R_6-S(O)_2-N(R_{21})-R_{10}$ - wherein R_8 and R_{10} are alkylene, R_{20} and R_{21} are loweralkyl, R_4 is loweralkyl, aryl, alkoxy, alkylamino, aryloxy or arylalkoxy and R_6 is loweralkyl, haloalkyl, alkoxyalkyl, aryl or arylalkyl.

- Another yet more preferred embodiment of the invention is a compound of formula (I)
- 25 or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i) loweralkyl, (ii) alkenyl,
- (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl,
- (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl,
- 30 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl,
- 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl,
- 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl,
- 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or
- dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and
- 35 carboxyalkoxy, (ix) heterocyclic (alkyl), (x) arylalkyl, (xi) aryloxyalkyl, (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonamidoalkyl, R_2 is substituted or unsubstituted
- 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl,

8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofurnayl,
 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ -
 wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N$ - wherein R_{11} and R_{12} are independently
 selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl, arylalkyl,
 5 heterocyclic, hydroxyalkyl, alkoxy, aminoalkyl, and trialkylaminoalkyl.

Another yet more preferred embodiment of the invention is a compound of formula (I)
 or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,
 tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1
 10 is (i) loweralkyl, (ii) alkenyl, (iii) heterocyclic (alkyl), (iv) aryloxyalkyl, (v) arylalkyl, (vi)
 aryl, (vii) (N-alkanoyl-N-alkyl)aminoalkyl, or (viii) alkylsulfonylamidoalkyl, R_2 is substituted
 or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-
 methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofurnayl, 4-methoxyphenyl,
 dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from
 15 loweralkyl, alkoxy and halogen and R_3 is $R_4-C(O)-R_5$ - wherein R_5 is alkylene and R_4 is
 $(R_{11})(R_{12})N$ - wherein R_{11} is loweralkyl and R_{12} is aryl, arylalkyl, hydroxyalkyl, alkoxy,
 aminoalkyl, trialkylaminoalkyl, or heterocyclic.

Another yet more preferred embodiment of the invention is a compound of formula (I)
 20 or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,
 tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1
 is (i) loweralkyl, (ii) alkenyl, (iii) heterocyclic (alkyl), (iv) aryloxyalkyl, (v) arylalkyl, (vi) (N-
 alkanoyl-N-alkyl)aminoalkyl, or (vii) alkylsulfonylamidoalkyl, (viii) phenyl, or (ix) substituted
 or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl,
 25 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl,
 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is
 selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, R_2 is
 substituted or unsubstituted 1,3-benzodioxolyl,
 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl,
 30 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl,
 fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy
 and halogen and R_3 is $R_6-S(O)_2-N(R_{21})-R_{10}$ - wherein R_{10} is alkylene, R_6 is loweralkyl,
 haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl and R_{21} is loweralkyl, haloalkyl,
 alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl.

Another yet more preferred embodiment of the invention is a compound of formula (I)
 or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,

tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i) substituted or unsubstituted

4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl,

3-fluoro-4-ethoxyphenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl or

- 5 1,4-benzodioxanyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy and alkoxyalkoxy, (ii) loweralkyl, (iii) alkenyl, (iv) heterocyclic (alkyl), (v) aryloxyalkyl, (vi) arylalkyl, (vii) (N-alkanoyl-N-alkyl)aminoalkyl, (viii) alkylsulfonamidalkyl, or (ix) phenyl, R_2 is substituted or unsubstituted

1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl,

- 10 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R_3 is alkoxy carbonyl or $R_6-S(O)_2-N(R_{21})-R_{10}$ wherein R_{10} is alkylene, R_6 is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{21} is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl.

15

Another yet more preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl or haloalkyl, Z is $-CH_2-$, R_1 is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, arylalkyl, aryl, (N-alkanoyl-N-alkyl)aminoalkyl, or alkylsulfonamidalkyl, and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from alkyl, aryl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

20

A still more preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl or haloalkyl, Z is $-CH_2-$, R_1 is substituted or unsubstituted

4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl,

4-trifluoromethylphenyl, 4-pentafluoroethylphenyl,

- 30 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, (ii) loweralkyl, (iii) alkenyl, (iv) heterocyclic (alkyl), (v) aryloxyalkyl, (vi) arylalkyl, (vii) (N-alkanoyl-N-alkyl)aminoalkyl, (viii) alkylsulfonamidalkyl, or (ix) phenyl, R_2 is 1,3-benzodioxolyl,

- 35 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is

(R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl, aryl, arylalkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, or heterocyclic.

Another still more preferred embodiment of the invention is a compound of formula
 5 (I) or (II) wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl or haloalkyl, Z is -CH₂-, R₁ is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, arylalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidoalkyl, phenyl, or alkoxyalkyl, R₂ is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl,
 10 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl, aryl, arylalkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, or heterocyclic.

A most highly preferred embodiment of the invention is a compound of formula (I) or
 15 (II) wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, Z is -CH₂-, R₁ is substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl,
 20 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl.

Another most highly preferred embodiment of the invention is a compound of formula
 30 (I) or (II) wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, Z is -CH₂-, R₁ is substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl,
 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is 1,3-benzodioxolyl,
 35 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ is loweralkyl and R₁₂ is aryl.

Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R₁ is substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl,

- 5 3-fluorophenyl, 2-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is substituted or unsubstituted 1,3-benzodioxolyl,
- 10 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R₃ is $R_6-S(O)_2-N(R_{21})-R_{10}-$ wherein R₁₀ is alkylene, R₆ is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R₂₁ is loweralkyl, haloalkyl or alkoxyalkyl.

- 15 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R₁ is substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 2-fluorophenyl, 3-fluoro-4-ethoxyphenyl,

- 20 4-methoxymethoxyphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl,
- 25 dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R₃ is $R_4-C(O)-R_5-$ wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ is alkyl and R₁₂ is selected from aryl, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

- 30 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R₁ is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, arylalkyl, aryl, (N-alkanoyl-N-alkyl)aminoalkyl, or alkylsulfonylamidoalkyl, and R₃ is $R_4-C(O)-R_5-$ wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from
- 35 alkyl, aryl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic, with the proviso that one of R₁₁ and R₁₂ is alkyl.

Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

5 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is loweralkyl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

10 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is alkenyl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

15 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is heterocyclic (alkyl), and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

20 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is aryloxyalkyl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

25 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is arylalkyl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

30 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is aryl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

35 Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is (N-alkanoyl-N-alkyl)aminoalkyl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

Another most highly preferred embodiment of the invention is a compound of formula (I) or (II) wherein n is 0, Z is $-\text{CH}_2-$, R_1 is alkylsulfonylamidoalkyl, and R_3 is $\text{R}_4-\text{C}(\text{O})-\text{R}_5$ - wherein R_4 is $(\text{R}_{11})(\text{R}_{12})\text{N}-$ as defined therein and R_5 is alkylene.

The present invention also relates to processes for preparing the compounds of formula (I) and (II) and to the synthetic intermediates employed in these processes.

The present invention also relates to a method of antagonizing endothelin in a mammal (preferably, a human) in need of such treatment, comprising administering to the mammal a therapeutically effective amount of a compound of formula (I) or (II).

The invention further relates to endothelin antagonizing compositions comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of formula (I) or (II).

In another embodiment of the invention is disclosed a method for inhibiting bone metastases in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

In another embodiment of the invention is disclosed a method for preventing new bone metastases in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

In another embodiment of the instant invention, therefore, is disclosed a method for inhibiting metastatic growth in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

In another embodiment of the invention is disclosed a method for inhibiting bone loss in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

In another embodiment of the instant invention, is disclosed a method for inhibiting bone turnover in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

In another embodiment of the invention is disclosed a method for the reduction of cancer related pain in a patient in need thereof which comprises administering to the patient a therapeutically effective amount of an endothelin ET-A receptor antagonist.

In another embodiment of the instant invention is disclosed therapeutically acceptable formulations of an endothelin ET-A receptor antagonist, optionally in the presence of a co-therapeutic agent, for use in these methods.

Brief Description of the Drawings

Figure 1 illustrates levels of interleukin-6 (IL-6) in a subject population treated with a placebo or 2.5 mg or 10 mg ABT-627.

Figure 2 illustrates levels of prostate specific antigen (PSA) in a subject population treated with a placebo or 2.5 mg or 10 mg of ABT-627.

Figure 3 illustrates VAS score levels relating to pain assessment in a subject population treated with a placebo or 2.5 mg or 10 mg of ABT-627.

Figure 4 illustrates crosslinked N-telopeptides (degradation) in a subject population treated with a placebo or 10 mg ABT-627.

Figure 5 illustrates bone alkaline phosphatase (BAP) formation in a subject population treated with a placebo or 10 mg ABT-627.

Figure 6 illustrates skeletal involvement in a subject population treated with a placebo or 10 mg ABT-627.

Figure 7 illustrates acid phosphatase levels in a subject population treated with a placebo or 10 mg ABT-627.

The compounds of the invention comprise two or more asymmetrically substituted carbon atoms. As a result, racemic mixtures, mixtures of diastereomers, as well as single diastereomers of the compounds of the invention are included in the present invention. The terms "S" and "R" configuration are as defined by the IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem. (1976) 45, 13 - 30.

The term "carboxy protecting group" as used herein refers to a carboxylic acid protecting ester group employed to block or protect the carboxylic acid functionality while the reactions involving other functional sites of the compound are carried out. Carboxy protecting groups are disclosed in Greene, "Protective Groups in Organic Synthesis" pp. 152-186 (1981), which is hereby incorporated herein by reference. In addition, a carboxy protecting group can be used as a prodrug whereby the carboxy protecting group can be readily cleaved *in vivo*, for example by enzymatic hydrolysis, to release the biologically active parent. T. Higuchi and V. Stella provide a thorough discussion of the prodrug concept in "Pro-drugs as Novel Delivery Systems", Vol 14 of the A.C.S. Symposium Series, American Chemical Society (1975), which is hereby incorporated herein by reference. Such carboxy protecting groups are well known to those skilled in the art, having been extensively used in the protection of carboxyl groups in the penicillin and cephalosporin fields, as described in U.S. Pat. No. 3,840,556 and 3,719,667, the disclosures of which are hereby incorporated herein by reference. Examples of esters useful as prodrugs for compounds containing

carboxyl groups can be found on pages 14-21 of "Bioreversible Carriers in Drug Design: Theory and Application", edited by E.B. Roche, Pergamon Press, New York (1987), which is hereby incorporated herein by reference. Representative carboxy protecting groups are C₁ to C₈ alkyl (e.g., methyl, ethyl or tertiary butyl and the like); haloalkyl; alkenyl; cycloalkyl and substituted derivatives thereof such as cyclohexyl, cyclopentyl and the like; cycloalkylalkyl and substituted derivatives thereof such as cyclohexylmethyl, cyclopentylmethyl and the like; arylalkyl, for example, phenethyl or benzyl and substituted derivatives thereof such as alkoxybenzyl or nitrobenzyl groups and the like; arylalkenyl, for example, phenylethenyl and the like; aryl and substituted derivatives thereof, for example, 5-indanyl and the like;

5 dialkylaminoalkyl (e.g., dimethylaminoethyl and the like); alkanoyloxyalkyl groups such as acetoxymethyl, butyryloxymethyl, valeryloxymethyl, isobutyryloxymethyl, isovaleryloxymethyl, 1-(propionyloxy)-1-ethyl, 1-(pivaloyloxy)-1-ethyl, 1-methyl-1-(propionyloxy)-1-ethyl, pivaloyloxymethyl, propionyloxymethyl and the like;

10 cycloalkanoyloxyalkyl groups such as cyclopropylcarbonyloxymethyl, cyclobutylcarbonyloxymethyl, cyclopentylcarbonyloxymethyl, cyclohexylcarbonyloxymethyl and the like; aroyloxyalkyl, such as benzoyloxymethyl, benzoyloxyethyl and the like;

15 arylalkylcarbonyloxyalkyl, such as benzylcarbonyloxymethyl, 2-benzylcarbonyloxyethyl and the like; alkoxycarbonylalkyl, such as methoxycarbonylmethyl, cyclohexyloxy carbonylmethyl, 1-methoxycarbonyl-1-ethyl, and the like;

20 alkoxycarbonyloxyalkyl, such as methoxycarbonyloxymethyl, t-butyloxy carbonyloxymethyl, 1-ethoxycarbonyloxy-1-ethyl, 1-cyclohexyloxy carbonyloxy-1-ethyl and the like; alkoxycarbonylaminoalkyl, such as t-butyloxy carbonylaminomethyl and the like; alkylaminocarbonylaminoalkyl, such as methylaminocarbonylaminomethyl and the like; alkanoylaminoalkyl, such as

25 acetylaminomethyl and the like; heterocycliccarbonyloxyalkyl, such as 4-methylpiperaziny carbonyloxymethyl and the like; dialkylaminocarbonylalkyl, such as dimethylaminocarbonylmethyl, diethylaminocarbonylmethyl and the like; (5-(loweralkyl)-2-oxo-1,3-dioxolen-4-yl)alkyl, such as (5-t-butyl-2-oxo-1,3-dioxolen-4-yl)methyl and the like;

30 and (5-phenyl-2-oxo-1,3-dioxolen-4-yl)alkyl, such as (5-phenyl-2-oxo-1,3-dioxolen-4-yl)methyl and the like.

The term "N-protecting group" or "N-protected" as used herein refers to those groups intended to protect the N-terminus of an amino acid or peptide or to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, "Protective Groups In Organic Synthesis," (John Wiley & Sons, New York (1981)), which is hereby incorporated by reference. N-protecting groups

35 comprise acyl groups such as formyl, acetyl, propionyl, pivaloyl,

t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxyacetyl, chlorobutyl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl and the like; carbamate forming groups such as benzyloxycarbonyl, p-chlorobenzoyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzoyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butylloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, phenylthiocarbonyl and the like; alkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like; and silyl groups such as trimethylsilyl and the like. Preferred N-protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, phenylsulfonyl, benzyl, t-butylloxycarbonyl (Boc) and benzyloxycarbonyl (Cbz).

The term "alkanoyl" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a carbonyl ($-C(O)-$) group. Examples of alkanoyl include acetyl, propionyl and the like.

The term "alkanoylamino" as used herein refers to an alkanoyl group as previously defined appended to an amino group. Examples alkanoylamino include acetamido, propionylamido and the like.

The term "alkanoylaminoalkyl" as used herein refers to $R_{43}-NH-R_{44}$ - wherein R_{43} is an alkanoyl group and R_{44} is an alkylene group.

The term "alkanoyloxyalkyl" as used herein refers to $R_{30}-O-R_{31}$ - wherein R_{30} is an alkanoyl group and R_{31} is an alkylene group. Examples of alkanoyloxyalkyl include acetoxymethyl, acetoxylethyl and the like.

The term "alkenyl" as used herein refers to a straight or branched chain hydrocarbon radical containing from 2 to 15 carbon atoms and also containing at least one carbon-carbon double bond. Alkenyl groups include, for example, vinyl (ethenyl), allyl (propenyl), butenyl, 1-methyl-2-buten-1-yl and the like.

The term "alkenylene" denotes a divalent group derived from a straight or branched chain hydrocarbon containing from 2 to 15 carbon atoms and also containing at least one carbon-carbon double bond. Examples of alkenylene include $-\text{CH}=\text{CH}-$, $-\text{CH}_2\text{CH}=\text{CH}-$, $-\text{C}(\text{CH}_3)=\text{CH}-$, $-\text{CH}_2\text{CH}=\text{CHCH}_2-$, and the like.

5 The term "alkenyloxy" as used herein refers to an alkenyl group, as previously defined, connected to the parent molecular moiety through an oxygen ($-\text{O}-$) linkage. Examples of alkenyloxy include allyloxy, butenyloxy and the like.

The term "alkoxy" as used herein refers to $\text{R}_{41}\text{O}-$ wherein R_{41} is a loweralkyl group, as defined herein. Examples of alkoxy include, but are not limited to, ethoxy, tert-butoxy, and
10 the like.

The term "alkoxyalkoxy" as used herein refers to $\text{R}_{80}\text{O}-\text{R}_{81}\text{O}-$ wherein R_{80} is loweralkyl as defined above and R_{81} is alkylene. Representative examples of alkoxyalkoxy groups include methoxymethoxy, ethoxymethoxy, t-butoxymethoxy and the like.

15 The term "alkoxyalkoxyalkyl" as used herein refers to an alkoxyalkoxy group as previously defined appended to an alkyl radical. Representative examples of alkoxyalkoxyalkyl groups include methoxyethoxyethyl, methoxymethoxymethyl, and the like.

The term "alkoxyalkyl" as used herein refers to an alkoxy group as previously defined appended to an alkyl radical as previously defined. Examples of alkoxyalkyl include, but are not limited to, methoxymethyl, methoxyethyl, isopropoxymethyl and the like.

20 The term "alkoxycarbonyl" as used herein refers to an alkoxyl group as previously defined appended to the parent molecular moiety through a carbonyl group. Examples of alkoxycarbonyl include methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl and the like.

The term "alkoxycarbonylalkenyl" as used herein refers to an alkoxycarbonyl group as previously defined appended to an alkenyl radical. Examples of alkoxycarbonylalkenyl
25 include methoxycarbonylethenyl, ethoxycarbonylethenyl and the like.

The term "alkoxycarbonylalkyl" as used herein refers to $\text{R}_{34}-\text{C}(\text{O})-\text{R}_{35}-$ wherein R_{34} is an alkoxy group and R_{35} is an alkylene group. Examples of alkoxycarbonylalkyl include methoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylmethyl and the like.

30 The term "alkoxycarbonylaminoalkyl" as used herein refers to $\text{R}_{38}-\text{C}(\text{O})-\text{NH}-\text{R}_{39}-$ wherein R_{38} is an alkoxy group and R_{39} is an alkylene group.

The term "alkoxycarbonyloxyalkyl" as used herein refers to $\text{R}_{36}-\text{C}(\text{O})-\text{O}-\text{R}_{37}-$ wherein R_{36} is an alkoxy group and R_{37} is an alkylene group.

The term "(alkoxycarbonyl)thioalkoxy" as used herein refers to an alkoxycarbonyl group as previously defined appended to a thioalkoxy radical. Examples of
35 (alkoxycarbonyl)thioalkoxy include methoxycarbonylthiomethoxy, ethoxycarbonylthiomethoxy and the like.

The term "alkoxyhaloalkyl" as used herein refers to a haloalkyl radical to which is appended an alkoxy group.

The terms "alkyl" and "loweralkyl" as used herein refer to straight or branched chain alkyl radicals containing from 1 to 15 carbon atoms including, but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl, n-pentyl, 1-methylbutyl, 2,2-dimethylbutyl, 2-methylpentyl, 2,2-dimethylpropyl, n-hexyl and the like.

The term "(N-alkanoyl-N-alkyl)aminoalkyl" as used herein refers to $R_{85}C(O)N(R_{86})R_{87}$ - wherein R_{85} is an alkanoyl as previously defined, R_{86} is loweralkyl, and R_{87} is alkylene.

The term "alkylamino" as used herein refers to $R_{51}NH$ - wherein R_{51} is a loweralkyl group, for example, ethylamino, butylamino, and the like.

The term "alkylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended an alkylamino group.

The term "alkylaminocarbonyl" as used herein refers to an alkylamino group, as previously defined, appended to the parent molecular moiety through a carbonyl ($-C(O)-$) linkage. Examples of alkylaminocarbonyl include methylaminocarbonyl, ethylaminocarbonyl, isopropylaminocarbonyl and the like.

The term "alkylaminocarbonylalkenyl" as used herein refers to an alkenyl radical to which is appended an alkylaminocarbonyl group.

The term "alkylaminocarbonylalkyl" as used herein refers to a loweralkyl radical to which is appended an alkylaminocarbonyl group.

The term "alkylaminocarbonylaminoalkyl" as used herein refers to $R_{40}-C(O)-NH-R_{41}$ - wherein R_{40} is an alkylamino group and R_{41} is an alkylene group.

The term "alkylene" denotes a divalent group derived from a straight or branched chain saturated hydrocarbon having from 1 to 15 carbon atoms by the removal of two hydrogen atoms, for example $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)_2CH_2-$ and the like.

The term "alkylsulfonylamidoalkyl" as used herein refers $R_{88}S(O)_2NHR_{89}$ - wherein R_{88} is loweralkyl and R_{89} is alkylene.

The term "alkylsulfonylamino" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a sulfonylamino ($-S(O)_2-NH-$) group. Examples of alkylsulfonylamino include methylsulfonylamino, ethylsulfonylamino, isopropylsulfonylamino and the like.

The term "alkynyl" as used herein refers to a straight or branched chain hydrocarbon radical containing from 2 to 15 carbon atoms and also containing at least one carbon-carbon triple bond. Examples of alkynyl include $-C\equiv C-H$, $H-C\equiv C-CH_2-$, $H-C\equiv C-CH(CH_3)-$ and the like.

The term "alkynylene" refers to a divalent group derived by the removal of two hydrogen atoms from a straight or branched chain acyclic hydrocarbon group containing from 2 to 15 carbon atoms and also containing a carbon-carbon triple bond. Examples of alkynylene include $-C\equiv C-$, $-C\equiv C-CH_2-$, $-C\equiv C-CH(CH_3)-$ and the like.

The term "aminoalkyl" as used herein refers to a $-NH_2$, alkylamino, or dialkylamino group appended to the parent molecular moiety through an alkylene.

The term "aminocarbonyl" as used herein refers to $H_2N-C(O)-$.

The term "aminocarbonylalkenyl" as used herein refers to an alkenyl radical to which is appended an aminocarbonyl ($NH_2C(O)-$) group.

The term "aminocarbonylalkoxy" as used herein refers to $H_2N-C(O)-$ appended to an alkoxy group as previously defined. Examples of aminocarbonylalkoxy include aminocarbonylmethoxy, aminocarbonylethoxy and the like.

The term "aminocarbonylalkyl" as used herein refers to a loweralkyl radical to which is appended an aminocarbonyl ($NH_2C(O)-$) group.

The term "trialkylaminoalkyl" as used herein refers to $(R_{90})(R_{91})(R_{92})N(R_{93})-$ wherein R_{90} , R_{91} , and R_{92} are independently selected from loweralkyl and R_{93} is alkylene.

The term "aryloxyalkyl" as used herein refers to $R_{32}-C(O)-O-R_{33}-$ wherein R_{32} is an aryl group and R_{33} is an alkylene group. Examples of aryloxyalkyl include benzoyloxymethyl, benzoyloxyethyl and the like.

The term "aryl" as used herein refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like. Aryl groups can be unsubstituted or substituted with one, two or three substituents independently selected from loweralkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxy carbonyl, alkoxy carbonylalkenyl, (alkoxy carbonyl)thioalkoxy, thioalkoxy, amino, alkylamino, dialkylamino, aminoalkyl, trialkylaminoalkyl, aminocarbonyl, aminocarbonylalkoxy, alkanoylamino, arylalkoxy, aryloxy, mercapto, cyano, nitro, carboxaldehyde, carboxy, carboxyalkenyl, carboxyalkoxy, alkylsulfonfylamino, cyanoalkoxy, (heterocyclic)alkoxy, hydroxy, hydroxalkoxy, phenyl and tetrazolylalkoxy. In addition, substituted aryl groups include tetrafluorophenyl and pentafluorophenyl.

The term "arylalkenyl" as used herein refers to an alkenyl radical to which is appended an aryl group, for example, phenylethenyl and the like.

The term "arylalkoxy" as used herein refers to $R_{42}O-$ wherein R_{42} is an arylalkyl group, for example, benzyloxy, and the like.

The term "arylalkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended an arylalkoxy group, for example, benzyloxymethyl and the like.

The term "arylalkyl" as used herein refers to an aryl group as previously defined, appended to a loweralkyl radical, for example, benzyl, phenethyl, 2,2-dimethyl-1-phenyl-1-propyl, 3,3-dimethyl-1-phenyl-1-butyl, and the like.

The term "aryloxy" as used herein refers to $R_{45}O$ - wherein R_{45} is an aryl group, for example, phenoxy, and the like.

The term "arylalkylcarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended an arylalkylcarbonyloxy group (i.e., $R_{62}C(O)O$ - wherein R_{62} is an arylalkyl group).

The term "aryloxyalkyl" refers to an aryloxy group as previously defined appended to an alkyl radical. Examples of aryloxyalkyl include phenoxymethyl, 2-phenoxyethyl and the like.

The term "carboxaldehyde" as used herein refers to a formaldehyde radical, $-C(O)H$.

The term "carboxy" as used herein refers to a carboxylic acid radical, $-C(O)OH$.

The term "carboxyalkenyl" as used herein refers to a carboxy group as previously defined appended to an alkenyl radical as previously defined. Examples of carboxyalkenyl include 2-carboxyethenyl, 3-carboxy-1-ethenyl and the like.

The term "carboxyalkoxy" as used herein refers to a carboxy group as previously defined appended to an alkoxy radical as previously defined. Examples of carboxyalkoxy include carboxymethoxy, carboxyethoxy and the like.

The term "cyanoalkoxy" as used herein refers to an alkoxy radical as previously defined to which is appended a cyano ($-CN$) group. Examples of cyanoalkoxy include 3-cyanopropoxy, 4-cyanobutoxy and the like.

The term "cycloalkanoyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended a cycloalkanoyloxy group (i.e., $R_{60}C(O)O$ - wherein R_{60} is a cycloalkyl group).

The term "cycloalkyl" as used herein refers to an aliphatic ring system having 3 to 10 carbon atoms and 1 to 3 rings including, but not limited to, cyclopropyl, cyclopentyl, cyclohexyl, norbornyl, adamantyl, and the like. Cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from loweralkyl, haloalkyl, alkoxy, thioalkoxy, amino, alkylamino, dialkylamino, hydroxy, halo, mercapto, nitro, carboxaldehyde, carboxy, alkoxy carbonyl and carboxamide.

The term "cycloalkylalkyl" as used herein refers to a cycloalkyl group appended to a loweralkyl radical, including but not limited to cyclohexylmethyl.

The term "dialkylamino" as used herein refers to $R_{56}R_{57}N$ - wherein R_{56} and R_{57} are independently selected from loweralkyl, for example diethylamino, methyl propylamino, and the like.

The term "dialkylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended a dialkylamino group.

The term "dialkylaminocarbonyl" as used herein refers to a dialkylamino group, as previously defined, appended to the parent molecular moiety through a carbonyl ($-C(O)-$) linkage. Examples of dialkylaminocarbonyl include dimethylaminocarbonyl, diethylaminocarbonyl and the like.

The term "dialkylaminocarbonylalkenyl" as used herein refers to an alkenyl radical to which is appended a dialkylaminocarbonyl group.

The term "dialkylaminocarbonylalkyl" as used herein refers to $R_{50}-C(O)-R_{51}$ wherein R_{50} is a dialkylamino group and R_{51} is an alkylene group.

The term "diarylalkyl," as used herein, refers to two aryl groups, as defined herein, attached to the parent molecular moiety through an alkyl group. The aryl groups of the diaryl can be optionally substituted with 1-5 alkyl substituents. Examples of "diaryl" include diphenylmethyl (benzhydryl), 2,2-diphenylethyl, 1,2-diphenylethyl, bis(2-methylphenyl)methyl, and the like.

The term "halo" or "halogen" as used herein refers to I, Br, Cl or F.

The term "haloalkenyl" as used herein refers to an alkenyl radical to which is appended at least one halogen substituent.

The term "haloalkoxy" as used herein refers to an alkoxy radical as defined above, bearing at least one halogen substituent, for example, 2-fluoroethoxy, 2,2,2-trifluoroethoxy, trifluoromethoxy, 2,2,3,3,3-pentafluoropropoxy and the like.

The term "haloalkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended a haloalkoxy group.

The term "haloalkyl" as used herein refers to a lower alkyl radical, as defined above, to which is appended at least one halogen substituent, for example, chloromethyl, fluoroethyl, trifluoromethyl or pentafluoroethyl and the like.

The term "heterocyclic ring" or "heterocyclic" or "heterocycle" as used herein refers to any 3- or 4-membered ring containing a heteroatom selected from oxygen, nitrogen and sulfur; or a 5-, 6- or 7-membered ring containing one, two or three nitrogen atoms; one oxygen atom; one sulfur atom; one nitrogen and one sulfur atom; one nitrogen and one oxygen atom; two oxygen atoms in non-adjacent positions; one oxygen and one sulfur atom in non-adjacent positions; or two sulfur atoms in non-adjacent positions. The 5-membered ring has 0-2 double bonds and the 6- and 7-membered rings have 0-3 double bonds. The nitrogen heteroatoms can be optionally quaternized. The term "heterocyclic" also includes bicyclic groups in which any of the above heterocyclic rings is fused to a benzene ring or a cyclohexane ring or another heterocyclic ring (for example, indolyl, dihydroindolyl, quinolyl,

isoquinolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, decahydroquinolyl, decahydroisoquinolyl, benzofuryl, dihydrobenzofuryl or benzothienyl and the like).

Heterocyclics include: aziridinyl, azetidiny, pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, piperidinyl, homopiperidinyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, quinoliny, isoquinoliny, benzimidazolyl, benzothiazolyl, benzoxazolyl, oxetanyl, furyl, tetrahydrofuranyl, thienyl, thiazolidinyl, isothiazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, pyrrolyl, pyrimidyl and benzothienyl.



Heterocyclics also include compounds of the formula $\text{X}^* \text{Y}^* \text{O}$ where X^* is $-\text{CH}_2-$ or $-\text{O}-$ and Y^* is $-\text{C}(\text{O})-$ or $[\text{C}(\text{R}'')_2]_v$ where R'' is hydrogen or $\text{C}_1\text{-C}_4$ -alkyl and v is 1, 2 or 3 such as 1,3-benzodioxolyl, 1,4-benzodioxanyl and the like. Heterocyclics also include bicyclic rings such as quinuclidinyl and the like.

Heterocyclics can be unsubstituted or monosubstituted or disubstituted with substituents independently selected from hydroxy, halo, oxo ($=\text{O}$), alkylimino ($\text{R}^*\text{N}=$ wherein R^* is a loweralkyl group), amino, alkylamino, dialkylamino, alkoxy, alkoxyalkoxy, aminoalkyl, trialkylaminoalkyl, haloalkyl, cycloalkyl, aryl, arylalkyl, $-\text{COOH}$, $-\text{SO}_3\text{H}$, alkoxycarbonyl, nitro, cyano and loweralkyl. In addition, nitrogen containing heterocycles can be N-protected.

The term "(heterocyclic)alkoxy" as used herein refers to a heterocyclic group as defined above appended to an alkoxy radical as defined above. Examples of (heterocyclic)alkoxy include 4-pyridylmethoxy, 2-pyridylmethoxy and the like.

The term "(heterocyclic)alkyl" as used herein refers to a heterocyclic group as defined above appended to a loweralkyl radical as defined above.

The term "heterocycliccarbonyloxyalkyl" as used herein refers to $\text{R}_{46}\text{-C}(\text{O})\text{-O-R}_{47}$ wherein R_{46} is a heterocyclic group and R_{47} is an alkylene group.

The term "hydroxy" as used herein refers to $-\text{OH}$.

The term "hydroxyalkenyl" as used herein refers to an alkenyl radical to which is appended a hydroxy group.

The term "hydroxyalkoxy" as used herein refers to an alkoxy radical as previously defined to which is appended a hydroxy ($-\text{OH}$) group. Examples of hydroxyalkoxy include 3-hydroxypropoxy, 4-hydroxybutoxy and the like.

The term "hydroxyalkyl" as used herein refers to a loweralkyl radical to which is appended a hydroxy group.

The term "leaving group" as used herein refers to a halide (for example, Cl, Br or I) or a sulfonate (for example, mesylate, tosylate, triflate and the like).

The term "mercapto" as used herein refers to -SH.

- 5 The terms "methylenedioxy" and "ethylenedioxy" refer to one or two carbon chains attached to the parent molecular moiety through two oxygen atoms. In the case of methylenedioxy, a fused 5 membered ring is formed. In the case of ethylenedioxy, a fused 6 membered ring is formed. Methylenedioxy substituted on a phenyl ring results in the



formation of a benzodioxolyl radical.



Ethylenedioxy substituted on a phenyl

ring results in the formation of a benzodioxanyl radical.

- 10 The term "substantially pure" as used herein means 95% or more of the specified compound.

The term "tetrazolyl" as used herein refers to a radical of the formula



or a tautomer thereof.


- 15 The term "tetrazolylalkoxy" as used herein refers to a tetrazolyl radical as defined above appended to an alkoxy group as defined above. Examples of tetrazolylalkoxy include tetrazolylmethoxy, tetrazolyloxy and the like.

The term "thioalkoxy" as used herein refers to $R_{70}S$ - wherein R_{70} is loweralkyl. Examples of thioalkoxy include, but are not limited to, methylthio, ethylthio and the like.

- 20 The term "thioalkoxyalkoxy" as used herein refers to $R_{80}S-R_{81}O$ - wherein R_{80} is loweralkyl as defined above and R_{81} is alkylene. Representative examples of alkoxyalkoxy groups include CH_3SCH_2O -, $EtSCH_2O$ -, $t-BuSCH_2O$ - and the like.


The term "thioalkoxyalkoxyalkyl" as used herein refers to a thioalkoxyalkoxy group appended to an alkyl radical. Representative examples of alkoxyalkoxyalkyl groups include $CH_3SCH_2CH_2OCH_2CH_2$ -, $CH_3SCH_2OCH_2$ -, and the like.

The term "*trans,trans*" as used herein refers to the orientation of substituents (R_1 and



R_2) relative to the central substituent R as shown

The term "*trans,cis*" as used herein refers to the orientation of substituents (R_1 and R_2)




relative to the central substituent R as shown

or

This definition encompasses both the case where R and R_2 are *cis* and R and R_1 are *trans* and the case where R_2 and R are *trans* and R and R_1 are *cis*.

The term "*cis,cis*" as used herein refers to the orientation of substituents (R_1 and R_2)

relative to the central substituent R as shown



Preferred compounds of the invention are selected from the group consisting of:

- trans-trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[3-(N-propyl-N-n-pentanesulfonylamino)propyl]-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(4-Methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-[2-(N-propyl-N-n-pentanesulfonylamino)ethyl]pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-n-pentanesulfonylamino)ethyl]pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-n-hexanesulfonylamino)ethyl]pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(4-Propoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-n-pentanesulfonylamino)ethyl]pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3,4-Difluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3,4-Difluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-n-pentanesulfonylamino)ethyl]pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-*n*-hexanesulfonylamino)ethyl]pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(3-chloropropanesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid;

5 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(3-chloropropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(4-methylbutanesulfonyl)amino)ethyl]pyrrolidine-3-carboxylic acid;

10 *trans,trans*-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(*n*-pentanesulfonyl)amino)ethyl]pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(2,2,3,3,3-pentafluoropropoxyethanesulfonyl)-amino)ethyl]pyrrolidine-3-carboxylic acid;

15 *trans,trans*-2-(1,4-Benzodioxan-6-yl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(*n*-pentanesulfonyl)amino)ethyl]pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(3-chloropropanesulfonyl)amino)-ethyl)pyrrolidine-3-carboxylic acid;

20 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-((2,2,2-trifluoroethoxyethane)sulfonyl)amino)-ethyl]pyrrolidine-3-carboxylic acid;

25 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(butanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(2-methylpropanesulfonyl)amino)ethyl]pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(butanesulfonylamino))ethyl)pyrrolidine-3-carboxylic acid;

30 *trans,trans*-2-(2-Methylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

35 *trans,trans*-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

- trans,trans*-2-(2-(2-Tetrahydro-2H-pyran)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2,2,4-Trimethyl-3-pentenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 5 *trans,trans*-2-(2,2,-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[[N-4-heptyl-N(2-methyl-3-fluorophenyl)] amino carbonylmethyl]-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 10 *trans,trans*-2-(2-(2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 15 *trans,trans*-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 20 *trans,trans*-2-(2,2-dimethylpentyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2,2,-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 25 *trans,trans*-2-(2-(2-Methoxyphenyl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2,2-Dimethyl-3-(*E*)-pentenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 30 (2*S*, 3*R*, 4*S*)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- (2*S*, 3*R*, 4*S*)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 35 *trans,trans*-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

- trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
 (2*R*,3*R*,4*S*)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid;
- 5 *trans,trans*-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylamino)butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-(4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylamino)butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;
- 10 *trans,trans*-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylamino)butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;
- 15 *trans,trans*-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-(4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-butyl-N-(4-dimethylamino)butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 20 *trans,trans*-2-(2,2,4-Trimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-(N-butyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2,4-Trimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylamino)butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;
- 25 *trans,trans*-2-(2,2,4-Trimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2,2,4-Trimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-(4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 30 *trans,trans*-2-(2,2,4-Trimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylamino)butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
- 35

trans,trans-2-(2,2-Dimethyl-2-(1,3-Dioxol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;

5 *trans,trans*-2-(2,2-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2,-Dimethyl-2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;

10 *trans,trans*-2-(2,2-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Methoxyphenyl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

15 *trans,trans*-2-(2-(2-Methoxyphenyl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Methoxyphenyl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

20 *trans,trans*-2-(2-(2-Methoxyphenyl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Methoxyphenyl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

25 *trans,trans*-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-((2-Methoxyphenoxy)-methyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

30 *trans,trans*-2-((2-Methoxyphenoxy)-methyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Methoxyphenoxy)-methyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

35 *trans,trans*-2-(2-(2-Oxo 1,2-dihydro pyridin-1-yl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyridin-1-yl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyridin-1-yl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

5 *trans,trans*-2-(2-(2-Oxopyridin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyridin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

10 *trans,trans*-2-(2-(2-Oxopyridin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopiperidin-1-yl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopiperidin-1-yl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

15 *trans,trans*-2-(2-(2-Oxopiperidin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopiperidin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

20 *trans,trans*-2-(2-(2-Oxopiperidin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopiperidin-1-yl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

25 *trans,trans*-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

30 *trans,trans*-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-trimethylammonio)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

35 *trans,trans*-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-trimethylammonibutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(3,3-Dimethyl-2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(3,3-Dimethyl-2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(3,3-Dimethyl-2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(4,4-Dimethyl-2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(N-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(4,4-Dimethyl-2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(4,4-Dimethyl-2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-dibutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

5 *trans,trans*-2-(2-(1-propanesultamyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

10 *trans,trans*-2-(2-(1-propanesultamyl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

15 *trans,trans*-2-(2-(1-propanesultamyl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-propanesultamyl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

20 *trans,trans*-2-(2-(1-propanesultamyl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

25 *trans,trans*-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

30 *trans,trans*-2-(2-(1-pyrazolyl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-dibutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

35 *trans,trans*-2-(2-(1-pyrazolyl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-(N,N-dibutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

5 *trans,trans*-2-(2-(2-oxazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(Oxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(Oxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

10 *trans,trans*-2-(2-(Oxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(Oxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

15 *trans,trans*-2-(2-(Oxazol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(Oxazol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

20 *trans,trans*-2-(2-(5-Methyloxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(5-Methyloxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

25 *trans,trans*-2-(2-(5-Methyloxazol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

30 *trans,trans*-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

35 *trans,trans*-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(2,5-Dioxopyrrolidin-1-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(propoxy)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyridin-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyrimidin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyrimidin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(Pyrimidin-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(1,3-benzodioxol-4-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
trans,trans-2-(2-(1,3-benzodioxol-4-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-4-heptyl-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid; and
trans,trans-2-(2-(1,3-benzodioxol-4-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[(N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid;
(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
(2S,3R,4S)-2-(2,2-Dimethylpent-(E)-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
(2S,3R,4S)-2-(2,2-Dimethylpent-(E)-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S,3R,4S)-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S,3R,4S)-2-(2-(2-Methoxyphenyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

5 or a pharmaceutically acceptable salt.

Most preferred compounds of the invention are selected from the group consisting of:

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2,-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-[[N-4-heptyl-N-(2-methyl-3-fluorophenyl)] aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2,-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Methoxyphenyl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2-Dimethyl-3-(E)-pentenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S, 3R, 4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S, 3R, 4S)-2-(2,2 Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S, 3R, 4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

5 (2R, 3R, 4S)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(((N-propyl-N-pentanesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid;

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

10 (2S,3R,4S)-2-(2,2-Dimethylpent-(E)-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S,3R,4S)-2-(2,2-Dimethylpent-(E)-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2S,3R,4S)-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid; and

15 (2S,3R,4S)-2-(2-(2-Methoxyphenyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-((bis-(o-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

20 *trans,trans*-2-[4-(2-methoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-(2,2-dimethyl-1-phenylpropyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-methoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-((bis-(o-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-isopropoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-(2,2-dimethyl-1-phenylpropyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

25 *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,3-dimethyl-1-phenylbutyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-isopropoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-((1-(o-tolyl)-1-(o-ethylphenyl)-methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(4-(2-(2-propoxyethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-N-phenyl-

30 N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid; and

trans,trans-2-(4-(2-methoxyethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-phenyl-N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid;

or a pharmaceutically acceptable salt thereof.

35

Methods for preparing the compounds of the invention are shown in Schemes I-XXIII.

Scheme I illustrates the general procedure for preparing the compounds of the invention when n and m are 0, Z is $-\text{CH}_2-$ and W is $-\text{CO}_2\text{H}$. A α -ketoester 1, where E is loweralkyl or a carboxy protecting group is reacted with a nitro vinyl compound 2, in the presence of a base (for example, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) or sodium ethoxide or sodium hydride and the like) in an inert solvent such as toluene, benzene, tetrahydrofuran or ethanol and the like. The condensation product 3 is reduced (for example, hydrogenation using a Raney nickel or platinum catalyst). The resulting amine cyclizes to give the dihydro pyrrole 4. Reduction of 4 (for example, sodium cyanoborohydride or catalytic hydrogenation and the like) in a protic solvent such as ethanol or methanol and the like gives the pyrrolidine compound 5 as a mixture of *cis-cis*, *trans,trans* and *cis,trans* products. Chromatographic separation removes the *cis-cis* isomer leaving a mixture of the *trans,trans* and *cis,trans* isomers which is further elaborated. The *cis-cis* isomer can be epimerized (for example, using sodium ethoxide in ethanol) to give the *trans,trans* isomer and then carried on as described below. The pyrrolidine nitrogen is (1) acylated or sulfonylated with $\text{R}_3\text{-X}$ (R_3 is $\text{R}_4\text{-C(O)-}$ or $\text{R}_6\text{-S(O)}_2\text{-}$ and X is a leaving group such as a halide (Cl is preferred) or X taken together with $\text{R}_4\text{-C(O)-}$ or $\text{R}_6\text{-S(O)}_2\text{-}$ forms an activated ester including esters or anhydrides derived from formic acid, acetic acid and the like, alkoxycarbonyl halides, N-hydroxysuccinimide, N-hydroxyphthalimide, N-hydroxybenzotriazole, N-hydroxy-5-norbornene-2,3-dicarboxamide, 2,4,5-trichlorophenol and the like) or (2) alkylated with $\text{R}_3\text{-X}$ where X is a leaving group (for example, X is a halide (for example, Cl, Br or I) or X is a leaving group such as a sulfonate (for example, mesylate, tosylate, triflate and the like)) in the presence of a base such as diisopropyl ethylamine or triethylamine and the like to give the N-derivatized pyrrolidine 6 which is still a mixture of *trans,trans* and *cis,trans* isomers. Hydrolysis of the ester 6 (for example, using a base such as sodium hydroxide in $\text{EtOH}/\text{H}_2\text{O}$) selectively hydrolyzes the *trans,trans* ester to give a mixture of 7 and 8, which are readily separated.

Scheme II illustrates a general procedure for preparing the compounds of the invention when n is 1, m is 0, Z is $-\text{CH}_2-$ and W is $-\text{CO}_2\text{H}$. A substituted benzyl chloride 9 is reacted with a lithio dithiane 10 in an inert solvent such as THF or dimethoxyethane to give the alkylated adduct 11. The anion of compound 11 is formed using a base such as n-butyllithium and then reacted with $\text{R}_1\text{-CH}_2\text{-X'}$ wherein X' is a leaving group such as a halide or sulfonate to give compound 12. The dithiane protecting group is cleaved (for example, using a mercuric salt in water) to give the keto compound 13. Reaction of ketone 13 with benzyl amine and formaldehyde gives the keto piperidine compound 14. Treatment of compound 14 with an activated nitrile such as trimethylsilyl cyanide followed by a dehydrating agent such as phosphorous oxychloride provides the isomeric ene nitriles 15. Reduction of the double

bond (for example, using sodium borohydride) affords the piperidiny nitrile 16. Hydrolysis of the nitrile using hydrochloric acid in the presence of a carboxy protecting reagent (for example, an alkyl alcohol) affords ester 17 (where E is a carboxy protecting group).

Debenzylation by catalytic hydrogenation under acidic conditions affords the free piperidine compound 18. Compound 18 is further elaborated by the procedures described in Scheme I for compound 5 to give the final product compound 19.

Scheme III illustrates a general procedure for preparing the compounds of the invention when m and n are 0, Z is -C(O)- and W is -CO₂H. alpha-Keto ester 20 (wherein E is loweralkyl or a carboxy protecting group) is reacted with an alpha-haloester 21 (where J is lower alkyl or a carboxy protecting group and the halogen is bromine, iodine or chlorine) in the presence of a base such as NaH or potassium tert-butoxide or lithium diisopropylamide in an inert solvent such as THF or dimethoxyethane to give diester 22. Treating compound 22 with R₃-NH₂ and heating in acetic acid gives the cyclic compound 23. The double bond is reduced (for example, by catalytic hydrogenation using a palladium on carbon catalyst or sodium cyanoborohydride reduction) to give pyrrolidone 24. Epimerization with sodium ethoxide in ethanol to give the desired *trans,trans* configuration, followed by sodium hydroxide hydrolysis of the ester, affords the desired *trans,trans* carboxylic acid 25.

Scheme IV illustrates a general procedure for preparing the compounds of the invention when n is 0, m is 1, Z is -CH₂- and W is -CO₂H. The *trans,trans* compound 7, prepared in Scheme I, is homologated by the Arndt-Eistert synthesis. The carboxy terminus is activated (for example, by making the acid chloride using thionyl chloride) to give compound 52, where L is a leaving group (in the case of an acid chloride, L is Cl). Compound 52 is treated with diazomethane to give the diazo ketone 53. Rearrangement of compound 53 (for example, using water or an alcohol and silver oxide or silver benzoate and triethylamine, or heating or photolysis in the presence of water or an alcohol) affords the acetic acid compound 54 or an ester which may be hydrolyzed. Compounds where m is from 2 to 6 can be obtained by repetition of the above described process.

A preferred embodiment is shown in Schemes V and VI. A benzoyl acetate 26 is reacted with a nitro vinyl benzodioxolyl compound 27 using 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) as the base in toluene to give compound 28. Catalytic hydrogenation using Raney nickel leads to reduction of the nitro group to an amine and subsequent cyclization to give the dihydropyrrole 29. The double bond is reduced with sodium cyanoborohydride to give the pyrrolidine compound 30 as a mixture of *cis-cis*, *trans,trans* and *cis,trans* isomers. Chromatography separates out the *cis-cis* isomer, leaving a mixture of the *trans,trans* and *cis,trans* isomers (31).

Scheme VI illustrates the further elaboration of the *trans,trans* isomer. The mixture (31) of *trans,trans* and *cis,trans* pyrrolidines described in Scheme IV is reacted with N-propyl

bromoacetamide in acetonitrile in the presence of ethyldiisopropylamine to give the alkylated pyrrolidine compound 32, still as a mixture of *trans,trans* and *cis,trans* isomers. Sodium hydroxide in ethanol-water hydrolyzes the ethyl ester of the *trans,trans* compound but leaves the ethyl ester of the *cis,trans* compound untouched, thus allowing separation of the

5 *trans,trans* carboxylic acid 33 from the *cis,trans* ester 34.

Scheme VII illustrates the preparation of a specific piperidinyll compound. Benzodioxolyl methyl chloride 35 is reacted with lithio dithiane 36 to give the alkylated compound 37. Treatment of compound 37 with 4-methoxybenzyl chloride in the presence of lithium diisopropylamide gives compound 38. Cleavage of the dithiane protecting group

10 using a mercuric salt in aqueous solution gives ketone 39. Treatment of 39 with benzylamine and formaldehyde gives the keto piperidine 40. Treatment of compound 40 with trimethylsilyl cyanide followed by phosphorous oxychloride gives the ene nitrile as a mixture of isomers 41. Sodium borohydride reduction of the double bond gives the piperidinyll nitrile 42. Hydrochloric acid hydrolysis in the presence of ethanol gives ethyl ester 43. The N-

15 benzyl protecting group is removed by catalytic hydrogenation to give the free piperidine compound 44. Compound 44 is further elaborated by the procedures described in Scheme V for compound 31 resulting in the formation of the N-derivatized carboxylic acid 45.

A preferred embodiment of the process shown in Scheme III is shown in Scheme VIII. 4-Methoxybenzoylacetate 46 (wherein E is loweralkyl or a carboxy protecting group) is

20 reacted with an benzodioxolyl alpha-bromoacetate 47 (wherein E is lower alkyl or a carboxy protecting group) in the presence of NaH in THF to give diester 48. Treating compound 48 with ethoxypropylamine and heating in acetic acid gives the cyclic compound 49. The double bond is reduced by catalytic hydrogenation using a palladium on carbon catalyst to give pyrrolidone 50. Epimerization with sodium ethoxide in ethanol to give the desired *trans,trans*

25 configuration is followed by sodium hydroxide hydrolysis of the ester to afford the desired *trans,trans* carboxylic acid 51.

Scheme IX illustrates the preparation of compounds where n is 0, Z is -CH₂-, and W is other than carboxylic acid. Compound 55, which can be prepared by the procedures described in Scheme IV, is converted (for example, using peptide coupling condition, e.g. N-

30 methylmorpholine, EDCI and HOBt, in the presence of ammonia or other amide forming reactions) to give carboxamide 56. The carboxamide is dehydrated (for example, using phosphorus oxychloride in pyridine) to give nitrile 57. Nitrile 57 under standard tetrazole forming conditions (sodium azide and triethylamine hydrochloride or trimethylsilylazide and tin oxide) is reacted to give tetrazole 58. Alternatively nitrile 57 is reacted with

35 hydroxylamine hydrochloride in the presence of a base (for example, potassium carbonate, sodium carbonate, sodium hydroxide, triethylamine, sodium methoxide or NaH) in a solvent such as DMF, DMSO, or dimethylacetamide to give amidoxime 59. The amidoxime 59 is

allowed to react with a methyl or ethyl chloroformate in a conventional organic solvent (such as, chloroform, methylene chloride, dioxane, THF, acetonitrile or pyridine) in the presence of a base (for example, triethylamine, pyridine, potassium carbonate and sodium carbonate) to give an O-acyl compound. Heating of the O-acyl amidoxime in an inert solvent (such as benzene, toluene, xylene, dioxane, THF, dichloroethane, or chloroform and the like) results in cyclization to compound 60. Alternatively reacting the amidoxime 59 with thionyl chloride in an inert solvent (for example, chloroform, dichloromethane, dioxane and THF and the like) affords the oxathiadiazole 61.

Scheme X illustrates the preparation of compounds in which R₃ is an acylmethylene group. A carboxylic acid 62 (where R₄ is as previously defined herein) is treated with oxalyl chloride in a solution of methylene chloride containing a catalytic amount of N,N-dimethylformamide to give the acid chloride. Treatment of the acid chloride with excess ethereal diazomethane affords a diazoketone, and then treatment with anhydrous HCl in dioxane gives the alpha-chloroketone 63. Pyrrolidine ester 5 where E is lower alkyl or a carboxy protecting group, prepared in Scheme I, is alkylated with the alpha-chloroketone 63 to provide alkylated pyrrolidine 64. Carboxy deprotection (for example, hydrolysis of an alkyl ester using lithium or sodium hydroxide in ethanol-water) gives the alkylated pyrrolidine acid 65.

Scheme XI illustrates the preparation of "reverse amides and sulfonamides". The carboxy protected pyrrolidine 5, prepared in Scheme I, is reacted with a difunctionalized compound X-R₈-X where R₈ is alkylene and X is a leaving group (for example a halide where Br is preferred) to give N-alkylated compound 66. Treatment of 66 with an amine (R₂₀NH₂) affords secondary amine 67. This amine (67) can be reacted with an activated acyl compound (for example, R₄-C(O)-Cl) and then carboxy deprotected (for example, hydrolysis of an ester or hydrogenation of a benzyl moiety) to afford amide 68. Alternatively amine 67 can be reacted with an activated sulfonyl compound (for example, R₆-S(O)₂-Cl) and then carboxy deprotected (for example, hydrolysis of an ester or hydrogenation of a benzyl moiety) to afford sulfonamide 69.

Scheme XII illustrates a method for synthesizing pyrrolidines by an azomethine ylide type [3+2]-cycloaddition to an acrylate. General structures such as compound 70 are known to add to unsaturated esters such as 71 to provide pyrrolidines such as compound 72 (O. Tsuge, S. Kanemasa, K. Matsuda, Chem. Lett. 1131-4 (1983), O. Tsuge, S. Kanemasa, T. Yamada, K. Matsuda, J. Org. Chem. 52 2523-30 (1987), and S. Kanemasa, K. Skamoto, O. Tsuge, Bull. Chem. Soc. Jpn. 62 1960-68 (1989)). A specific example is also shown in Scheme XII. Silylimine 73 is reacted with acrylate 74 in the presence of trimethylsilyl triflate and tetrabutylammonium fluoride to give the desired pyrrolidine 75 as a mixture of isomers. This method can be modified to provide the N-acetamido derivatives directly by reacting 73

and 74 with the appropriate bromoacetamide (for example, dibutyl bromoacetamide) in the presence of tetrabutylammonium iodide and cesium fluoride to give compound 76.

Scheme XIII illustrates a method for producing an enantiomerically pure pyrrolidine 80, which can be further elaborated on the pyrrolidine nitrogen. Intermediate racemic pyrrolidine ester 77 (for example, prepared by the procedure described in Scheme V) is Boc-nitrogen protected (for example, by treatment with Boc_2O) and then the ester is hydrolyzed (for example, using sodium or lithium hydroxide in ethanol and water) to give t-butyl carbamoyl pyrrolidine carboxylic acid 78. The carboxylic acid is converted to its (+)-cinchonine salt, which can be recrystallized (for example from ethyl acetate and hexane or chloroform and hexane) to afford the diastereomerically pure salt. This diastereomerically pure salt can be neutralized (for example, with sodium carbonate or citric acid) to afford enantiomerically pure carboxylic acid 79. The pyrrolidine nitrogen can be deprotected (for example, using trifluoroacetic acid) and the ester reformed by the use of ethanolic hydrochloric acid to give salt 80. Alternatively one can use ethanol HCl to cleave the protecting group and form the ester in one step. The pyrrolidine nitrogen can be further elaborated (for example, by treatment with the dibutyl amide of bromoacetamide in acetonitrile in the presence of diisopropylethylamine) to give optically active compound 81. The use of (-)-cinchonine will give the opposite enantiomer.

Scheme XIV describes another procedure for preparation of pyrrolidines. Pyrrolidines may be synthesized by the use of an azomethine ylide cycloaddition to an acrylate derivative as described by Cottrell, I. F., et al., J. Chem. Soc., Perkin Trans. 1, 5: 1091-97 (1991). Thus, the azomethine ylide precursor 82 (where R_{55} is hydrogen or methyl) is condensed with a substituted acrylate 83 (wherein R_2 is as described herein and R_{56} is loweralkyl) under acidic conditions to afford the substituted pyrrolidine 84. The N-protecting group can be removed (for example, by hydrogenolysis of an N-benzyl group) to give 85, which can be alkylated under the conditions described above to provide the N-substituted pyrrolidine 86. Standard ester hydrolysis of 86 produces the desired pyrrolidine carboxylic acid 87.

A preferred process is shown in Scheme XV. Nitro vinyl compound (88) is reacted with beta-keto ester 89 in the presence of a base such as sodium ethoxide and the like or a trialkylamine such as triethylamine or diisopropylethylamine and the like or an amidine such as DBU and the like in an inert solvent such as THF, toluene, DMF, acetonitrile, ethyl acetate, isopropyl acetate or methylene chloride and the like at a temperature of from about 0°C to about 100°C for a period of time from about 15 minutes to overnight to give compound 90. Reduction of the nitro group followed by cyclization was effected for example by catalytic hydrogenation with a hydrogen pressure of from about atmospheric pressure to 300 p.s.i. over from about 1 hour to about 1 day of compound 90 in an inert solvent such as THF, ethyl acetate, toluene, ethanol, isopropanol, DMF or acetonitrile and the like, using a hydrogenation

catalyst such as Raney nickel, palladium on carbon, a platinum catalyst, such as platinum oxide, platinum on carbon or platinum on alumina and the like, or a rhodium catalyst, such as rhodium on carbon or rhodium on alumina and the like, and the like affords intermediate nitrone 91a or a mixture of nitrone 91a and imine 91b. The reaction mixture comprising the

5 nitrone or nitrone/imine mixture is treated with an acid such as trifluoroacetic acid or acetic acid or sulfuric acid or phosphoric acid or methanesulfonic acid and the like, and the hydrogenation is continued to give pyrrolidine compound 92 as the *cis,cis*-isomer. Epimerization at C-3 is effected by treatment of compound 92 with a base such as sodium ethoxide, potassium t-butoxide, lithium t-butoxide or potassium t-amylxide and the like or a

10 trialkylamine such as triethylamine or diisopropylethylamine and the like or an amidine such as DBU and the like in an inert solvent such as ethanol, ethyl acetate, isopropyl acetate, THF, toluene or DMF and the like at a temperature of from about -20° C to about 120° C to give the *trans,trans* compound 93. Compound 93 itself can optionally be resolved into enantiomers prior to reacting with X-R₃. The substantially pure (i.e., at least 95% of the desired isomer)

15 optically active (+)-isomer of compound 93 is obtained by treatment of a mixture of the (+)-isomer and the (-)-isomer of 93 with S-(+)-mandelic acid, D-tartaric acid or D-dibenzoyl tartaric acid and the like in a solvent such as acetonitrile, ethyl acetate, isopropyl acetate, ethanol or isopropanol and the like. The (+)-isomer of 93 selectively crystallizes as the salt, leaving the (-)-isomer of 93 in solution. Alternatively, the substantially pure (i.e., at

20 least 95% of the desired isomer) optically active (-)-isomer of compound 93 can be selectively crystallized by reaction of a mixture of the (+)-isomer and the (-)-isomer of 93 with L-tartaric acid, L-dibenzoyl tartaric acid or L-pyrogutamic acid and the like, leaving the desired (+)-isomer of compound 93 in solution.

Compound 93 (racemic or optically active) is reacted with X-R₃ (where X is a leaving

25 group (for example, a halide or a sulfonate) and R₃ is as previously defined) using a base such as diisopropylethylamine, triethylamine, sodium bicarbonate or potassium carbonate and the like in an inert solvent such as acetonitrile, THF, toluene, DMF or ethanol and the like at a temperature of from about 0° C to about 100° C to give the intermediate ester 94. The ester can be isolated or converted *in situ* to the carboxylic acid (95) using hydrolysis conditions

30 such as a base such as sodium hydroxide or lithium hydroxide or potassium hydroxide and the like in a solvent such as ethanol-water or THF-ethanol and the like.

A more detailed description of the preparation of some specific analogs is provided in Schemes XVI-XXI. Aliphatic alpha-ketoesters (Scheme XVI) may be prepared by copper-catalyzed addition of a Grignard reagent (for example, propylmagnesium bromide) to an

35 unsaturated ester, for example, ethyl 3,3-dimethylacrylate. The resultant ester is hydrolyzed, for example with sodium hydroxide in aqueous alcohol, and is homologated in stepwise fashion to the corresponding alpha-ketoester, for example by activation using

carbonyldiimidazole and condensation with magnesio-ethoxymalonate. Alternatively, olefinic alpha-ketoesters may be prepared by Claisen rearrangement of the corresponding allylic alcohols; hydrolysis and homologation as described above produce the desired alpha-ketoester.

5 N-alkyl,O-alkyl bromohydroxamates are prepared according to Scheme XVII. N-Boc-O-allyl hydroxylamine is alkylated with an alkyl halide, for example using sodium hydride as base; the double bond is selectively reduced, for example using hydrogen and a palladium catalyst. After removal of the Boc protecting group, for example with TFA, the resultant amine is acylated, for example using bromoacetyl bromide.

10 The alpha-ketoesters described in Scheme XVI may be converted to pyrrolidine derivatives as described in Scheme XVIII. Michael addition onto a nitrostyrene derivative can be catalyzed with base, for example DBU or potassium t-butoxide; the resultant adduct is hydrogenated, for example using Raney Nickel as catalyst, to give an imine, which is reduced further, for example using sodium cyanoborohydride under controlled pH. A mixture of
15 isomers are generated, in which the trans-trans is generally preferred.

Scheme XIX describes several strategies for resolving the racemic pyrrolidines described above. Treatment with a chiral acid, for example (S)-(+)-mandelic acid, may provide a crystalline derivative, which can be further enriched through recrystallization. The salt may be washed with base to extract the resolving agent and return the optically active
20 pyrrolidine product. Alternatively, the amino ester can be N-protected (for example with Boc-anhydride) and hydrolyzed (for example with sodium hydroxide) to give the corresponding N-protected amino acid. Activation of the acid, for example as the pentafluorophenyl ester, followed by coupling with a chiral nonracemic oxazolidinone anion, provides the corresponding acyloxazolidinone diastereomers, which may be separated
25 chromatographically. Alcoholysis of one acyloxazolidinone diastereomer, followed by cleavage of the N-protecting group, returns an optically enriched amino ester. A similar transformation may be accomplished through coupling of the protected amino acid with a chiral nonracemic amino alcohol. After chromatographic separation of the resultant diastereomers, the amide is cleaved and the protecting group is removed to provide optically
30 enriched product.

Optically active amino esters prepared as described above may be alkylated (Scheme XX) with a variety of electrophiles, for example dibutyl bromoacetamide, N-butyl,N-alkoxy bromoacetamide, N-(4-heptyl)-N-(3-methyl-4-fluorophenyl) bromoacetamide, or N-(Ω -hydroxyalkyl)-N-alkyl haloacetamide. Hydrolysis of the resultant ester, for example using
35 sodium hydroxide in aqueous alcohol, provides the product.

For one particular class of electrophile, N-(Ω -hydroxyalkyl)-N-alkyl haloacetamides, further transformations of the alkylation product are possible (Scheme XXI). Activation (for


example using methanesulfonyl chloride) of the alcohol, followed by displacement with halogen (for example, using lithium bromide) provides the corresponding halide.

Displacement of halide with an amine, for example dimethylamine, provides the corresponding amino ester, which may be hydrolyzed as previously described to provide product.


5

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000


Scheme I




Scheme II




Scheme III




Scheme IV




Scheme V




Scheme VI




Scheme VII




Scheme VII cont.




Scheme VIII




Scheme IX





Scheme X




Scheme XI




Scheme XII




Scheme XIII




Scheme XIV




Scheme XV




SCHEME XVI






SCHEME XVII




SCHEME XVIII




SCHEME XIX



SCHEME XX



SCHEME XXI




Other amines may be prepared according to Scheme XXII. An aryl aldehyde or ketone (aldehyde shown), which may be acquired commercially or prepared, for example, through a Friedel-Crafts acylation of a benzene derivative with an acyl halide, is reacted with an amine, for example ammonia, hydroxylamine or the like. The resultant imine is reduced,

for example using sodium borohydride or sodium cyanoborohydride or a metal like zinc or tin or the like, to give the corresponding optionally substituted carbonylamine, which is converted to the target compound according to the procedures described above.

5


Scheme XXII



Still other amines may be prepared according to Scheme XXIII. An optionally substituted aryl halide (R_{11a} is the optionally substituted aryl and X is bromo or iodo) is reacted with a metallated amine (for example, lithium tert-butylamide, or sodium benzylamide, or the like) to provide an optionally substituted aniline. This compound is reacted with an oxidized nitrogen compound, for example nitrous acid or the like, and the resultant compound is reduced using a metal like zinc or tin or palladium or the like to provide an N,N-disubstituted hydrazine, which is converted to the target compound according to the procedures described above.

15

Scheme XXIII









5



m is 0 to 6;

- 20






; and

R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl,


alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen;

or a salt thereof;

or a compound of the formula:



or



wherein n is 0 or 1;

m is 0 to 6;

W is (a) $-C(O)_2-G$ where G is hydrogen or a carboxy protecting group, (b) $-PO_3H_2$,

(c) $-P(O)(OH)E$ where E is hydrogen, loweralkyl or arylalkyl,

(d) $-CN$,

5 (e) $-C(O)NHR_{17}$ where R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,

(i) hydroxy,

10 (j) alkoxy,


(k) sulfonamido,

(l) $-C(O)NHS(O)_2R_{16}$ where R_{16} is loweralkyl, haloalkyl,

phenyl or

dialkylamino,

(m) $-S(O)_2NHC(O)R_{16}$,



15 (n) ,



(o) ,



(p) ,




(q) ,



(r) ,



20 (s) ,



R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl,

- 5 alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and
- 10 $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen; or a salt thereof.

Preferred intermediates include compounds of formula (III), (IV) and (V) wherein m is zero or 1;


- 15 W is $-CO_2-G$ wherein G is hydrogen or a carboxy protecting group, and R_1 and R_2 are as defined above; or the substantially pure (+)- or (-)-isomer thereof.

Particularly preferred intermediates are compounds of formula (III), (IV) and (V) wherein

- 20 n and m are both 0;
- W is $-CO_2-G$ wherein G is hydrogen or a carboxy protecting group; and R_1 is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl,
- 25 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii)
- 30 (N-alkanoyl-N-alkyl)aminoalkyl, and (xiii) alkylsulfonylamidoalkyl, and R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl,

1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofurnayl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen; or the substantially pure (+)- or (-)-isomer thereof.

- 5 Other compounds which are useful as intermediates for the preparation of compounds of the invention are:



wherein n is 0 or 1;

m is 0 to 6;

R_{5b} is alkylene;

Q is a leaving group;

- 5 W is (a) -C(O)₂-G where G is hydrogen or a carboxy protecting group, (b) -PO₃H₂,
 (c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,
 (d) -CN,
 (e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,
 (f) alkylaminocarbonyl,
 10 (g) dialkylaminocarbonyl,
 (h) tetrazolyl,
 (i) hydroxy,
 (j) alkoxy,
 (k) sulfonamido,
 15 (l) -C(O)NHS(O)₂R₁₆ where R₁₆ is loweralkyl, haloalkyl, phenyl or

dialkylamino,

(m) -S(O)₂NHC(O)R₁₆,



(n) ,



(o) ,






(p) ,



(q) ,



(r) ,




R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl,

- 5 alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, 10 heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ - wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen;


or a salt thereof;

or a compound of the formula:

15



or



wherein n is 0 or 1;

20 m is 0 to 6;

R_{5b} is alkylene;

Q is a leaving group;

W is (a) $-C(O)_2-G$ where G is hydrogen or a carboxy protecting group, (b) $-PO_3H_2$,


(c) $-P(O)(OH)E$ where E is hydrogen, loweralkyl or arylalkyl,

25 (d) $-CN$,

(e) $-C(O)NHR_{17}$ where R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

- (g) dialkylaminocarbonyl,
 (h) tetrazolyl,
 (i) hydroxy,
 (j) alkoxy,
 5 (k) sulfonamido,
 (l) $-\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}_{16}$ where R_{16} is loweralkyl, haloalkyl, phenyl or dialkylamino,
 (m) $-\text{S}(\text{O})_2\text{NHC}(\text{O})\text{R}_{16}$.



(n) ,



(o) ,



(p) ,




(q) ,




(r) ,



(s) ,



(t) , or



(u) ; and

5

10

15

R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, 5 alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidalkyl, heterocyclic, (heterocyclic)alkyl and (R_{aa})(R_{bb})N-R_{cc}- wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R₁ and R₂ is other than hydrogen;

10 or a salt thereof.

Preferred intermediates include compounds of formula (VI), (VII) and (VIII) wherein m is zero or 1;

R_{5b} is alkylene;

Q is a leaving group;

15 W is -CO₂-G wherein G is hydrogen or a carboxy protecting group, and R₁ and R₂ are as defined above; or the substantially pure (+)- or (-)-isomer thereof.

Particularly preferred intermediates are compounds of formula (VI), (VII) and (VIII) wherein

20 n and m are both 0;

R_{5b} is alkylene;

Q is a leaving group;

W is -CO₂-G wherein G is hydrogen or a carboxy protecting group;

and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl,

25 (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl,

4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl,

4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl,

3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl,

4-hydroxyphenyl, 4-t-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or


30 dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, and (xiii) alkylsulfonamidalkyl, and R₂ is

substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl,

1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-

35 methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen; or the substantially pure (+)- or (-)-isomer thereof.

Other compounds which are useful as intermediates for the preparation of compounds of the invention are:



wherein n is 0 or 1;

m is 0 to 6;

R_{5b} is alkylene;

R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl;

W is (a) -C(O)₂-G where G is hydrogen or a carboxy protecting group, (b) -PO₃H₂,

(c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,

(d) -CN,

(e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,


(i) hydroxy,







(j) alkoxy,

(k) sulfonamido,

(l) -C(O)NHS(O)₂R₁₆ where R₁₆ is loweralkyl, haloalkyl, phenyl or dialkylamino,

(m) -S(O)₂NHC(O)R₁₆,







R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen;

or a salt thereof;

or a compound of the formula:



or



(X)

(XI)

wherein n is 0 or 1;

m is 0 to 6;

5 R_{5b} is alkylene;

R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl,

cycloalkylalkyl, aryl or arylalkyl;

W is (a) $-C(O)_2-G$ where G is hydrogen or a carboxy protecting group, (b) $-PO_3H_2$,

(c) $-P(O)(OH)E$ where E is hydrogen, loweralkyl or arylalkyl,

10 (d) $-CN$,

(e) $-C(O)NHR_{17}$ where R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,

15 (i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

(l) $-C(O)NHS(O)_2R_{16}$ where R_{16} is loweralkyl, haloalkyl, phenyl or dialkylamino,

(m) $-S(O)_2NHC(O)R_{16}$,



(n) ,







(o) ,



(p) ,



(q) ,



- 5 R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen; or a salt thereof.

- 15 Preferred intermediates include compounds of formula (IX), (X) and (XI) wherein m is zero or 1; R_{5b} is alkylene; R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl;
- 20 W is $-CO_2-G$ wherein G is hydrogen or a carboxy protecting group, and R_1 and R_2 are as defined above; or the substantially pure (+)- or (-)-isomer thereof.

Particularly preferred intermediates are compounds of formula (IX), (X) and (XI) wherein

- 25 n and m are both 0; R_{5b} is alkylene; R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl;

W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, and (xiii) alkylsulfonylamidoalkyl, and R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen; or the substantially pure (+)- or (-)-isomer thereof.

The foregoing may be better understood by reference to the following examples which are provided for illustration and not intended to limit the scope of the inventive concept. The following abbreviations are used: Boc for tert-butyloxycarbonyl, Cbz for benzyloxycarbonyl, DBU for 1,8-diazabicyclo[5.4.0]undec-7-ene, EDCI for 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, EtOAc for ethyl acetate, EtOH for ethanol, HOBT for 1-hydroxybenzotriazole, Et₃N for triethylamine, TFA for trifluoroacetic acid and THF for tetrahydrofuran.

Example 1

trans,trans- 2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 1A

Ethyl 2-(4-methoxybenzoyl)-4-nitromethyl-3-(1,3-benzodioxole-5-yl)butyrate

To ethyl (4-methoxybenzoyl)acetate (23.0 g, 0.104 mol), prepared by the method of Krapcho *et al.*, Org. Syn. 47, 20 (1967), and 5-(2-nitrovinyl)-1,3-benzodioxole (17.0 g, 0.088 mol) dissolved in 180 mL of toluene and heated to 80 °C was added 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.65 g) with stirring. The mixture was heated until all the nitro starting material dissolved. The solution was stirred without heating for 30 minutes (min) and then an additional 0.65 g of DBU was added. After stirring an additional 45 minutes, thin layer chromatography (5% ethyl acetate in methylene chloride) indicated the absence of nitro starting material. Toluene (200 mL) was added, and the organic phase was washed with

dilute hydrochloric acid and NaCl solution. The organic phase was dried over sodium sulfate and then concentrated under reduced pressure. The residue obtained was chromatographed on silica gel eluting with 3:1 hexane-ethyl acetate to give 21.22 g of the desired product as a mixture of isomers and 9.98 g. of recovered ethyl (4-methoxybenzoyl)acetate.

Example 1B

Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-4,5-dihydro-3H-pyrrole-3-carboxylate

The compound resulting from Example 1A (21 g) in 500 mL of ethanol was hydrogenated under 4 atmospheres of hydrogen pressure using a Raney nickel 2800 catalyst (51 g). (The Raney nickel was washed with ethanol three times before use.) The catalyst was removed by filtration, and the solution was concentrated under reduced pressure. The residue obtained was chromatographed on silica gel eluting with 8.5% ethyl acetate in methylene chloride to give 12.34 g of the desired product.

Example 1C

Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate as a mixture of *cis-cis*; *trans,trans*; and *cis,trans*-isomers

The compound resulting from Example 1B (11.89 g, 0.324 mol) was dissolved in 27 mL of tetrahydrofuran and 54 mL of ethanol. Sodium cyanoborohydride (2.35 g, 0.374 mol) and 5 mg bromocresol green were added. To this blue solution was added dropwise a solution of 1:2 concentrated HCl in ethanol at such a rate that the color was kept at light yellow-green. After the yellow color persisted without additional HCl, the solution was stirred an additional 20 minutes. The solution was concentrated *in vacuo* and then partitioned between chloroform and an aqueous potassium bicarbonate solution. The organic phase was separated, dried over sodium sulfate, and concentrated under reduced pressure. The residue was chromatographed on silica gel eluting with 85:15 ethyl acetate-hexane to give 5.96 g. of a mixture of 64% *trans,trans*-compound and 34% *cis,trans*-compound. Further elution with pure ethyl acetate gave 0.505 g of an unknown solid followed by 3.044 g of pure *cis,cis*-compound.

Example 1D

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The mixture of 64% *trans,trans*- and 34% *cis,trans*-pyrrolidines (the mixture resulting from Example 1C) (5.72 g, 15.50 mmol), ethyldiisopropylamine (4.20 g, 32.56 mmol), and N-propyl bromoacetamide (3.42 g, 19.0 mmol), prepared by the method of Weaver, W.E. and

Whaley, W.M., J. Amer. Chem. Soc., 69: 515 (1947), in 30 mL of acetonitrile was heated at 50 °C for 1 hour. The solution was concentrated *in vacuo*. The residue was dissolved in toluene, shaken with potassium bicarbonate solution, dried over sodium sulfate and concentrated *in vacuo* to give 7.16 g of product as a mixture of *trans,trans*- and *cis,trans*-ethyl esters.

This mixture was dissolved in a solution of 50 mL of ethanol and 15 mL of water containing 5.00 g of sodium hydroxide and stirred for 3 hours at room temperature. The solution was concentrated *in vacuo* and 60 mL of water added. The mixture was extracted with ether to remove the unreacted *cis,trans*-ethyl ester. The aqueous phase was treated with hydrochloric acid until slightly cloudy. It was then further neutralized with acetic acid to give the crude acid product. The crude product was filtered and purified by dissolving it in tetrahydrofuran, drying over sodium sulfate, concentrating *in vacuo*, and crystallizing from ether to give 3.230 g of the title compound. m.p. 151-153 °C. ¹H NMR (CD₃OD, 300 MHz) δ 0.87 (t, J = 7 Hz, 3H), 1.49 (sextet, J = 7 Hz, 2H), 2.84 (d, J = 16 Hz, 1H), 2.95-3.20 (m, 4H), 3.20 (d, J = 16 Hz, 1H), 3.34-3.42 (m, 1H), 3.58-3.66 (m, 1H), 3.78 (s, 3H), 3.88 (d, J = 10 Hz, 1H), 5.92 (s, 2H), 6.75 (d, J = 8 Hz, 1H), 6.86 (dd, J = 8 Hz, J = 1 Hz, 1H), 6.90 (d, J = 9 Hz, 2H), 7.02 (d, J = 1 Hz, 1H), 7.40 (d, J = 9 Hz, 2H).

Example 2

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the method described in Example 1D, 300 mg of the mixture of 64% *trans,trans*- and 34% *cis,trans*-pyrrolidines (the mixture resulting from Example 1C), 220 mg of diisopropylethylamine and 184 mg iodoacetamide were reacted at 45 °C in 1 mL acetonitrile to give 291 mg of a mixture of *trans,trans*- and *cis,trans*- N-alkylated esters. A portion (270 mg.) was hydrolyzed with 200 mg NaOH in 1 mL of water and 3 mL of ethanol; a chloroform extraction was used to remove the unreacted *cis,trans*-ethyl ester. The isolation and purification procedures described in Example 1D were used to give 134 mg of the title compound. m.p. 246-248 °C. ¹H NMR (DMSO-d₆, 300 MHz) δ 2.61 (d, J = 16 Hz, 1H), 2.71 (t, J = 9 Hz, 1H), 2.90 (t, J = 9 Hz, 1H), 2.98 (d, J = 16 Hz, 1H), 3.25-3.35 (m, 1H), 3.45-3.55 (m, 1H), 3.71 (s, 3H), 3.75 (d, J = 10 Hz, 1H), 6.00 (s, 2H), 6.81 (s, 2H), 6.90 (d, J = 8 Hz, 2H), 7.10 (s, 1H), 7.17 (s, 1H), 7.34 (s, 1H), 7.38 (d, J = 8 Hz, 2H).

Example 3

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-fluorobenzyl)-pyrrolidine-3-carboxylic acid

Using the method described in Example 1D, 300 mg of the mixture of 64% *trans,trans*- and 34% *cis,trans*- pyrrolidines (the mixture resulting from Example 1C), 220 mg of diisopropylethylamine and 185 mg of 4-fluorobenzyl bromide were reacted at room temperature for 3 hours in 1 mL of acetonitrile to give 387 mg of a mixture of *trans,trans*- and *cis,trans*-N-alkylated esters. A portion (360 mg) was hydrolyzed with 250 mg NaOH in 1 mL of water and 4 mL of ethanol to give 160 mg of the title compound as an amorphous powder. ¹H NMR (CDCl₃, 300 MHz) δ 2.74 (t, J = 9 Hz, 1H), 2.95 (t, J = 7 Hz, 1H), 2.98 (d, J = 14, 1H), 3.07 (dd, J = 9 Hz, 1 Hz, 1H), 3.42-3.53 (m, 1H), 3.70 (d, J = 9 Hz, 1H), 3.78 (d, J = 14, 1H), 3.81 (s, 3H), 5.92 (s, 2H), 6.70 (d, J = 8 Hz, 1H), 6.77 (dd, J = 8 Hz, 1 Hz, 1H), 6.91 (d, J = 9 Hz, 2H), 6.94 - 7.00 (m, 3H), 7.20 - 7.25 (M, 1H), 7.44 (d, J = 9 Hz, 2H).

Example 4

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-ethoxyethyl)-pyrrolidine-3-carboxylic acid

Using the method described in Example 1D, 300 mg. of the mixture of 64% *trans,trans*- and 34% *cis,trans*-pyrrolidines (the mixture resulting from Example 1C), 220 mg of diisopropylethylamine and 152 mg of 2-bromoethyl ethyl ether were refluxed in 1.5 mL acetonitrile for 3 hours (bath temperature at 95 °C) to give 346 mg of a mixture of *trans,trans*- and *cis,trans*-esters. Hydrolysis with 250 mg NaOH in 1 mL of water and 3 mL of ethanol afforded 140 mg of the title compound. m.p. 88 - 90 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.25 (t, J = 7 Hz, 3H), 2.21-2.32 (m, 1H), 2.70-2.80 (m, 1H), 2.85-2.94 (m, 2H), 3.38-3.55 (m, 6H), 3.67 (d, J = 10 Hz, 1H), 3.79 (s, 3H), 5.94 (s, 2H), 6.72 (d, J = 8 Hz, 1H), 6.84 (m, 1H), 6.84 (d, J = 9 Hz, 2H), 7.08 (d, J = 1 Hz, 1H), 7.33 (d, J = 9 Hz, 2H).

Example 5

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-propoxyethyl)-pyrrolidine-3-carboxylic acid

Using the method described in Example 1D, 520 mg of the mixture resulting from Example 1C, 364 mg of diisopropylethylamine, 50 mg potassium iodide and 350 mg 2-chloroethyl propyl ether were reacted at 125 °C in 0.5 mL acetonitrile for 4 hours to give 517 mg of a mixture of *trans,trans*- and *cis,trans*-esters. A portion (500 mg) was hydrolyzed with 315 mg NaOH in 1 mL of water and 4 mL of ethanol to give 225 mg of the title compound as an amorphous powder. ¹H NMR (CDCl₃, 300 MHz) δ 0.87 (t, J = 7 Hz, 3H), 1.53 (sextet, J = 7 Hz, 2H), 2.28-2.41 (m, 1H), 2.71-2.83 (m, 1H), 2.92-3.08 (m, 2H), 3.30 (t, J = 7 Hz, 2H), 3.40-3.60 (m, 4H), 3.72-3.83 (m, 1H), 3.76 (s, 3H), 5.92 (s, 2H), 6.71 (d, J = 8 Hz, 2H), 6.74 (dd, J = 8 Hz, 1 Hz), 6.71 (d, J = 9 Hz, 2H), 7.07 (d, J = 9 Hz, 2H), 7.73 (d, J = 9 Hz, 2H).

Example 6

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(2-methoxyethoxy)ethyl]-pyrrolidine-3-carboxylic acid

Example 6A

Ethyl trans,trans-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl) pyrrolidine-3-carboxylate

To the pure *cis,cis*-compound resulting from Example 1C (3.02 g) dissolved in 10 mL of ethanol was added 20 drops of a solution of 21% sodium ethoxide in ethanol. The reaction mixture was refluxed overnight, at which time thin layer chromatography in ethyl acetate indicated the absence of starting material. The NaOEt was neutralized with HCl in ethanol, and the solution was concentrated *in vacuo*. The residue was taken up in toluene and extracted with potassium bicarbonate in water. The toluene was dried over sodium sulfate and concentrated under reduced pressure to give 2.775 of the title compound which was pure by TLC (ethyl acetate).

Example 6B

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(2-methoxyethoxy)ethyl]-pyrrolidine-3-carboxylic acid

Using the method described in Example 1D, 250 mg of the compound resulting from Example 6A, 150 mg of 2-(2-methoxyethoxy)ethyl bromide and 175 mg diisopropylethylamine in 1 mL acetonitrile were heated at 100 °C for 3 hours to give 229 mg of the *trans,trans*-ester. A portion (200 mg) was hydrolyzed with 125 mg NaOH in 1 mL of water and 2 mL of ethanol to give 151 mg of the title compound as an amorphous powder. ¹H NMR (CD₃OD, 300 MHz) δ 2.9-3.9 (m, 13H), 3.81 (s, 3H), 4.49 (d, J = 10 Hz, 1H), 5.94 (s, 2H), 6.79 (d, J = 8 Hz, 1H), 6.89 (dd, J = 8 Hz, 1 Hz, 1H), 7.00 (d, J = 9 Hz, 2H), 7.05 (d, J = 1 Hz, 1H), 7.49 (d, J = 9 Hz, 2H).

Example 7

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(2-pyridyl)ethyl]-pyrrolidine-3-carboxylic acid

The compound resulting from Example 6A (250 mg), 2-vinyl pyridine (355 mg) and one drop of acetic acid were dissolved in 2-methoxyethanol, and stirred at 100 °C for 2.5 hours. Toluene was added, and the solution was washed with potassium bicarbonate solution. The solution was dried over potassium bicarbonate and concentrated *in vacuo*. Toluene was added and the solution re-concentrated. This was done until the odor of 2-vinylpyridine was gone. The residue was taken up in hot heptane, filtered to remove a small amount of insoluble

impurity, and concentrated *in vacuo* to give 225 mg of intermediate ester. The ester was hydrolyzed by the method described in Example 1D to give 202 mg of the title compound as the dihydrate. m.p. 77-80 °C. ¹H NMR (CD₃OD, 300 MHz) δ 2.8 - 3.3 (m, 6H), 3.55-3.70 (m, 2H), 3.76 (s, 3H), 3.99 (d, J = 10 Hz, 1H), 5.92 (d, J = 1 Hz, 2H), 6.72 (d, J = 8 Hz, 1H),
 5 6.80 (dd, J = 8 Hz, 1 Hz), 6.85 (d, J = 9 Hz, 2H), 6.92 (d, J = 1 Hz, 1H), 7.20 (d, J = 9 Hz, 2H), 7.20-7.32 (m, 2H), 7.70-7.80 (m, 2H), 8.40 (d, J = 4 Hz, 1H).

Example 8

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(morpholin-4-ylcarbonyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 6A (300 mg) and 164 mg triethylamine dissolved in 2 mL of methylene chloride and cooled in an ice bath was added 146 mg 1-morpholinocarbonyl chloride. The mixture was stirred 3 hours at room temperature. Toluene was added and the solution was washed with potassium bicarbonate solution, dried over
 15 sodium sulfate and concentrated *in vacuo* to give the intermediate ester. The ester was hydrolyzed by the method described in Example 1D to give 288 mg of the title compound. m.p. 244-246 °C. ¹H NMR (DMSO-d₆, 300 MHz) δ 2.96 (dd, J = 12, 1 Hz, 13 Hz, 1H), 3.03-3.13 (m, 2H), 3.20-3.30 (m, 2H), 3.40-3.60 (m, 5H), 3.74 (s, 3H), 3.70-3.85 (m, 3H), 5.10 (d, J = 10 Hz, 1H), 5.99 (d, J = 1 Hz, 2H), 6.80-6.90 (m, 2H), 6.87 (d, J = 9 Hz, 2H), 7.07 (s, 1H),
 20 7.25 (d, J = 9 Hz, 2H).

Example 9

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxole-5-yl)-1-(butylaminocarbonyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 6A (300 mg) dissolved in 2 mL tetrahydrofuran and cooled in an ice bath was added 88 mg of butyl isocyanate. After 40 minutes at room temperature, toluene was added, and the solution was concentrated *in vacuo* to give the intermediate ester. The ester was hydrolyzed by the method described in Example
 25 1D to give 232 mg of the title compound. m.p. 220-221 °C. ¹H NMR (DMSO-d₆, 300 MHz) δ 0.78 (t, J = 7 Hz, 3H), 1.10 (sextet, J = 7 Hz, 2H), 1.22 (quintet, J = 7 Hz, 2H), 2.78-3.05 (m, 3H), 3.40-3.56 (m, 2H), 3.74 (s, 3H), 3.95-4.05 (m, 1H), 4.93 (d, J = 9 Hz, 1H), 5.80 (t, broad, J = 7 Hz, 1H), 5.99 (s, 2H), 6.78-6.86 (m, 2H), 6.88 (d, J = 9 Hz, 2H), 7.00 (d, J = 1 Hz, 1H), 7.12 (d, J = 9 Hz, 2H).

Example 10*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-methoxyphenylaminocarbonyl)-3-pyrrolidine-3-carboxylic acid

The compound resulting from Example 6A (300 mg) was treated with 133 mg of 4-methoxyphenyl isocyanate by the procedure described in Example 9. The resulting ester was hydrolyzed with NaOH using the method described in Example 1D to give 279 mg of the title compound. m.p. 185-187 °C. ¹H NMR (CDCl₃, 300 MHz) δ 3.23 (dd, J = 12 Hz, 13 Hz, 1H), 3.55-3.68 (m, 2H), 3.72 (s, 3H), 3.83 (s, 3H), 4.50-4.65 (m, 1H), 5.06 (d, J = 10 Hz, 1H), 5.90 (s, 1H), 5.95 (s, 1H), 6.72 (d, J = 9 Hz, 2H), 6.7-6.8 (m, 3H), 6.92 (d, J = 9 Hz, 2H), 6.97 (d, J = 9 Hz, 2H), 7.37 (d, J = 9 Hz, 2H).

Example 11*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-acetylpyrrolidine-3-carboxylic acid

The compound resulting from Example 6A (250 mg) in 0.5 mL of toluene was treated with 200 mg of acetic anhydride. After stirring 2 hours at room temperature, water was added and the acetic acid neutralized with potassium bicarbonate. The mixture was extracted with toluene to give 273 mg of the intermediate ester. A portion of the ester (200 mg) was hydrolyzed using the method of Example 1D to give 211 mg of the title compound. m.p. 248-250 °C. Rotational isomers are seen in the NMR. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 1.55 and 2.00 (s, 3H), 2.94 and 3.03 (dd, J = 12 Hz, 13 Hz, 1H), 3.3-3.6 (m, 2H), 3.72 and 3.76 (s, 3H), 4.12 and 4.28 (dd, J = 12 Hz, 7 Hz, 1H), 4.95 and 5.04 (d, J = 10 Hz, 1H), 6.00 (s, 2H), 6.75-6.87 (m, 3H), 6.95 and 7.04 (d, J = 9 Hz, 2H), 7.18 and 7.32 (d, J = 9 Hz, 2H).

Example 12*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-furoyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 6A (300 mg) and 164 mg triethylamine dissolved in 2 mL methylene chloride and cooled in an ice bath was added 138 mg of 2-furoyl chloride. The mixture was stirred 30 minutes at room temperature and then worked up by the procedures described in Example 8 to give the intermediate ester. The ester was hydrolyzed by the procedure described in Example 1D to give 269 mg of the title compound as an amorphous powder. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 3.06 (dd, J = 12 Hz, 13 Hz, 1H), 3.3-3.6 (m, 2H), 4.25 (m, 1H), 5.19 (d, J = 10 Hz, 1H), 6.67.4 (m, 8H), 7.8-7.9 (m, 1H).

Example 13

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(phenylaminocarbonyl)-pyrrolidine-3-carboxylic acid

Starting with the compound resulting from Example 6A, phenyl isocyanate and the procedures described in Example 9, the title compound was prepared. m.p. 209-211 °C. ¹H NMR (DMSO-d₆, 300 MHz) δ 3.03 (dd, 1H), 3.55 (m, 1H), 3.70 (m, 1H), 3.72 (s, 3H), 4.15 (m, 1H), 5.13 (d, 1H), 6.00 (s, 2H), 6.88 (m, 5H), 7.07-7.20 (m, 3H), 7.30 (d, 2H), 7.38 (d, 2H), 8.20 (bs, 1H).

Example 14

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(allylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared. m.p. 138-140 °C. ¹H NMR (CDCl₃, 300 MHz) δ 2.84 (d, 1H), 2.90-3.10 (dt, 2H), 3.28 (d, 1H), 3.35 (dd, 1H), 3.62 (m, 1H), 3.72-3.97 (m, 3H), 3.80 (s, 3H), 5.13 (bd, 2H), 5.80 (m, 1H), 5.97 (s, 2H), 6.74-6.97 (m, 5H), 7.38 (d, 2H).

Example 15

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(n-butylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared. m.p. 105-107 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.90 (t, 3H), 1.30 (m, 2H), 1.45 (m, 2H), 2.80 (d, 1H), 2.87-3.35 (m, 6H), 3.62 (m, 1H), 3.80 (s, 3H), 5.97 (s, 2H), 6.75-6.92 (m, 5H), 7.28 (d, 2H).

Example 16

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(n-propyl)-N-methylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared as an amorphous solid. Rotational isomers are seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) δ 0.73, 0.84 (2t, 3H), 1.49 (m, 2H), 2.80 (dd, 1H), 2.85 (2s, 3H), 2.95-3.20 (m, 3H), 3.20-3.40 (m, 1H), 3.40 (d, 1H), 3.60 (m, 1H), 3.79 (s, 3H), 5.93 (s, 2H), 6.73 (d, 1H), 6.86 (m, 2H), 7.03 (m, 1H), 7.32 (d, 2H).

Example 17

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(pyrrolidin-1-ylcarbonylmethyl)-pyrrolidine-3-carboxylic acid

5

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-
(isobutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

10

175-177 °C. ¹H NMR (CD₃OD, 300 MHz) δ 0.87 (dd, 6H), 1.75 (septet, 1H), 2.85 (d, 1H), 2.90-3.10 (m, 4H), 3.23 (d, 1H), 3.40 (m, 1H), 3.58-3.67 (m, 1H), 3.78 (s, 3H), 3.89 (d, 1H), 5.92 (s, 2H), 6.76 (d, 1H), 6.86 (dd, 1H), 6.91 (d, 2H), 7.02 (d, 1H), 7.40 (d, 2H).

15

Using the procedures described in Example 1 the title compound was prepared. m.p.

137-139 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.34 (m, 2H), 1.62 (m, 4H), 1.90 (m, 2H), 2.76 (d, 1H), 2.90 (t, 1H), 3.04 (dd, 1H), 3.22 (d, 1H), 3.28 (dd, 1H), 3.40 (m, 1H), 3.80 (s, 3H), 4.15 (m, 1H), 5.97 (d, 2H), 6.75-6.95 (m, 5H), 7.27 (m, 2H).

20

Example 20*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(morpholin-4-ylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 2.82 (d, 1H), 3.00 (m, 2H), 3.24 (m, 1H), 3.30-3.52 (m, 4H), 3.52-3.75 (m, 8H), 3.80 (s, 3H), 5.95 (s, 2H), 6.75 (d, 1H), 6.84 (d, 3H), 7.00 (s, 1H), 7.28 (d, 2H).

Example 21*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-phenoxyethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 4 the title compound was prepared as an amorphous solid. ¹H NMR (CD₃OD, 300 MHz) δ 2.82 (m, 1H), 2.96 (dd, 1H), 3.13 (m, 1H), 3.32 (m, 1H), 3.51-3.70 (m, 2H), 3.77 (s, 3H), 4.00 (d, 1H), 4.07 (m, 2H), 5.91 (s, 2H), 6.72 (d, 1H), 6.80-6.95 (m, 6H), 7.03 (d, 1H), 7.22 (dd, 2H), 7.39 (d, 2H).

Example 22*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-methoxyethylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared. m.p. 107-109 °C. ¹H NMR (CD₃OD, 300 MHz) δ 2.82 (d, 1H), 2.97 (q, 2H), 3.21 (d, 1H), 3.38 (m, 1H), 3.32 (s, 3H), 3.44 (m, 4H), 3.62 (m, 1H), 3.79 (s, 3H), 3.86 (d, 1H), 5.93 (s, 2H), 6.76 (d, 1H), 6.85 (dd, 1H), 6.91 (d, 2H), 7.01 (d, 1H), 7.38 (d, 2H).

Example 23*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-butoxyethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 4 the title compound was prepared. m.p. 53-55 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.88 (t, J=7Hz, 3H), 1.32 (sextet, J=7Hz, 2H), 1.50 (pentet, J=7Hz, 2H), 2.27 (tt, J=6Hz, 6Hz, 1H), 2.92 (q, J=10Hz, 2H), 3.35 (t, J=7Hz, 2H), 3.42-3.56 (m, 4H), 3.68 (d, J=10Hz, 1H), 3.78 (s, 3H), 5.94 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.83 (d, J=9Hz, 2H), 6.82-6.87 (m, 1H), 7.06 (d, J=2Hz, 1H), 7.32 (d, J=9Hz, 2H). MS m/e 442 (M+H)⁺.

Example 24

trans,trans-2-(1,3-Benzodioxol-5-yl)-4-(4-methoxyphenyl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 and substituting ethyl (1,3-benzodioxol-5-ylcarbonyl)acetate for ethyl (4-methoxybenzoyl)acetate and 4-(2-nitrovinyl)anisole for 5-(2-nitrovinyl)-1,3-benzodioxol-5-yl afforded the title compound. m.p. 97-99 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.78 (t, J=7Hz, 3H), 1.39 (sextet, J=7Hz, 2H), 2.72 (d, J=16Hz, 1H), 2.74 (t, J=10Hz, 1H), 2.80-3.10 (m, 4H), 3.26-3.38 (m, 1H), 3.53 (m, 1H), 3.73 (s, 3H), 3.80 (d, J=10Hz, 2H), 7.80 (t, J=6Hz, 1H). MS (DCI/NH₃) m/e 441 (M+H)⁺.

Example 25

trans,trans-2-(1,3-Benzodioxol-5-yl)-4-(4-methoxyphenyl)-1-(2-propoxyethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 5 and substituting ethyl (1,3-benzodioxol-5-ylcarbonyl)acetate for ethyl (4-methoxybenzoyl)acetate and 4-(2-nitrovinyl)anisole for 5-(2-nitrovinyl)-1,3-benzodioxol-5-yl afforded the title compound. m.p. 67-69 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.89 (t, J=7Hz, 3H), 1.56 (sextet, J=7Hz, 2H), 2.33 (m, 1H), 2.78-3.00 (m, 3H), 3.32 (t, J=7Hz, 2H), 3.45-3.57 (m, 4H), 3.73 (m, 1H), 3.79 (s, 3H), 5.93 (s, 2H), 6.22 (d, J=8Hz, 1H), 6.85 (d, J=8Hz, 3H), 6.98 (s, 1H), 7.37 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 428 (M+H)⁺.

Example 26

trans,trans-2-(1,3-Benzodioxol-5-yl)-4-(4-methoxyphenyl)-1-[2-(2-methoxyethoxy)ethyl]-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 4 and substituting the starting materials described in Example 25 and using 2-(2-methoxyethoxy)ethylbromide to alkylate the pyrrolidine nitrogen afforded the title compound. m.p. 85-86 °C. ¹H NMR (CD₃OD, 300 MHz) δ 3.18-3.90 (m, 15H), 3.79 (s, 3H), 4.57 (d, J=10Hz, 1H), 6.02 (s, 2H), 6.91 (d, J=8Hz, 1H), 6.95 (d, J=9Hz, 2H), 7.06 (dd, J=8Hz, 1H), 7.12 (dd, J=1Hz, 1H), 7.37 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 444 (M+H)⁺.

Example 27

trans,trans-2-(1,3-Benzodioxol-5-yl)-4-(4-methoxyphenyl)-1-(butoxyethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 4, substituting the starting materials described in Example 25 and using 2-ethoxyethylbromide to alkylate the pyrrolidine nitrogen afforded the title compound. m.p. 54-56 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.89 (t, J=7Hz,

3H), 1.44 (sextet, J=7Hz, 2H), 1.52 (pentet, J=7Hz, 2H), 2.40 (m, 1H), 2.74-2.98 (m, 3H), 3.46 (t, J=7Hz, 2H), 3.42-3.56 (m, 4H), 3.68 (d, J=10Hz, 1H), 3.80 (s, 3H), 5.93 (dd, J=6Hz, 1H, 2H), 6.72 (d, J=8Hz, 1H), 6.74 (dd, J=9Hz, 3H), 6.96 (s, 1H), 7.36 (d, J=9Hz, 2H).

5

Example 28

trans,trans-2-(4-Methoxyphenyl)-4-(1,4-benzodioxan-6-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 and substituting 6-(2-nitrovinyl)-1,4-benzodioxane for 5-(2-nitrovinyl)-1,3-benzodioxole afforded the title compound. m.p. 80-81 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.89 (t, J=7Hz, 3H), 1.49 (sextet, J=7Hz, 2H), 2.78 (d, J=16Hz, 1H), 2.92 (t, J=10Hz, 1H), 3.05-3.43 (m, 5H), 3.24 (d, J=16Hz, 1H), 3.52-3.62 (m, 1H), 3.80 (s, 3H), 3.80 (t, J=10Hz, 1H), 4.27 (s, 4H), 6.74-6.93 (m, 5H), 7.29 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 455 (M+H)⁺.

15

Example 29

trans,trans-2-(4-Methoxyphenyl)-4-(1,4-benzodioxan-6-yl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, substituting 6-(2-nitrovinyl)-1,4-benzodioxane for 5-(2-nitrovinyl)-1,3-benzodioxole and alkylating the pyrrolidine nitrogen with N-methyl-N-propyl bromoacetamide afforded the title compound. m.p. 74-76 °C. Rotational isomers are seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) δ 0.73, 0.83 (2t, J=7Hz, 3H), 1.48 (m, 2H), 2.78 (dd, 1H), 2.85 (2s, 3H), 2.96-3.15 (m, 3H), 3.27-3.42 (m, 3H), 3.52-3.60 (m, 1H), 3.75 (d, 1H), 3.78 (s, 3H), 4.22 (s, 4H), 6.80-6.98 (m, 5H), 7.32 (d, 2H). MS (DCI/NH₃) m/e 469 (M+H)⁺.

25

Example 30

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-butylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. Rotational isomers are seen in the NMR. ¹H NMR (CD₃OD, 300 MHz) δ 0.86 (2t, 3H), 1.04-1.50 (m, 4H), 2.85 (2s, 3H), 2.93-3.20 (m, 4H), 3.40 (m, 2H), 3.52 (dd, 1H), 3.60 (m, 1H), 3.80 (s, 3H), 3.85 (m, 1H), 5.91 (s, 2H), 6.74 (d, 1H), 6.83-6.95 (m, 3H), 7.03 (dd, 1H), 7.35 (dd, 2H).

35

Example 31

trans,trans-2-(4-Methoxy-2-methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-butylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 31AEthyl 2-(4-methoxy-2-methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

Using the procedures described in Examples 1A and 1B and substituting ethyl (4-methoxy-2-methoxymethoxybenzoyl)acetate for ethyl (4-methoxybenzoyl)acetate afforded ethyl 2-(4-methoxy-2-methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-4,5-dihydro-3H-pyrrole-3-carboxylate.

The above dihydro pyrrole carboxylate (3.0 g, 7.0 mmol) was dissolved in 20 mL of methanol, treated with 500 mg of 10% Pd/C and placed under hydrogen atmosphere for 32 hours. The catalyst was removed by filtration and the filtrate was concentrated under reduced pressure and chromatographed on silica gel eluting with ethyl acetate to afford the title compound (1.9 g, 63%) as the *cis-cis* isomer.

Example 31B*trans,trans*-2-(4-Methoxy-2-methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-butylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound resulting from Example 31A was epimerized by the procedure described in Example 6A. The resulting *trans,trans* compound (100 mg, 0.23 mmol) was then reacted by the procedures described in Example 1D substituting N-methyl-N-butyl bromoacetamide for N-propyl bromoacetamide to give the title compound (75 mg, 62%). m.p. 65-67 °C. Rotational isomers are seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) δ 0.64, 0.68 (2t, J=7Hz, 3H), 1.14, 1.12 (2 sextet, J=7Hz, 2H), 1.40-1.48 (m, 2H), 2.86, 2.89 (2s, 3H), 2.95-3.42 (m, 6H), 3.50 (s, 3H), 3.43-3.65 (m, 2H), 3.78 (s, 3H), 4.30 (t, J=7Hz, 1H), 5.09 (q, J=7Hz, 2H), 5.92 (s, 2H), 6.55 (dd, J=3Hz, 1H), 6.68 (s, 1H), 6.72 (s, 1H), 6.85 (2t, J=1Hz, 1H), 7.04 (t, J=1Hz, 1H), 7.42 (dd, J=3Hz, 1H).

Example 32*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-ethoxypropyl)-pyrrolidin-5-one-3-carboxylic acidExample 32AEthyl 2-(4-methoxybenzoyl)-3-carbomethoxy-1,3-benzodioxole-5-propionate

To ethyl (4-methoxybenzoyl)acetate (4.44 g, 0.02 mmol) dissolved in 20 mL of anhydrous THF was added in portions 480 mg of NaH. The mixture was stirred for 30 minutes under nitrogen at ambient temperature. Methyl (1,3-benzodioxol-5-yl) bromoacetate (5.46 g, 0.02 mol) in 5 mL of THF was added. The mixture was stirred overnight at ambient

5

Ethyl 1-(3-ethoxypropyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-4,5-dihydro-5-oxo-1H-pyrrole-3-carboxylate

10

Ethyl 1-(3-ethoxypropyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidin-5-one-3-carboxylate

20

25

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-ethoxypropyl)-pyrrolidin-5-one-3-carboxylic acid

30

35

3.78 (s, 3H), 3.88 (d, J=10Hz, 1H), 4.72 (d, J=10Hz, 1H), 6.02 (s, 2H), 6.74 (dd, J=8Hz, 1H, 1H), 6.87 (d, J=8Hz, 2H), 6.98 (d, J=8Hz, 2H), 7.38 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 442 (M+H)⁺.

Example 33

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-methoxybenzyl)-pyrrolidin-5-one-3-carboxylic acid

Following the procedures described in Example 32 and substituting 3-methoxybenzylamine for 3-ethoxypropylamine afforded the title compound (123 mg, 65%).
m.p. 150-152 °C. ¹H NMR (CD₃OD, 300 MHz) δ 2.96 (dd, J=8Hz, 10Hz, 1H), 3.72 (s, 3H), 3.80 (s, 3H), 4.06 (d, J=10Hz, 1H), 4.58 (d, J=8Hz, 1H), 4.92 (q, J=16Hz, 2H), 5.92 (s, 2H), 6.55-6.63 (m, 2H), 6.82 (d, J=8Hz, 4H), 6.94 (d, J=8Hz, 2H), 7.15-7.22 (m, 3H). MS (DCI/NH₃) m/e 475 (M+H)⁺.

Example 34

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-diisoamylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 0.70 -0.90 (m, 12H), 1.10-1.60 (m, 10H), 2.75 (d, J=13Hz, 1H), 2.90-3.10 (m, 4H), 3.15 - 3.30 (m, 2H), 3.40 (d, J=10Hz, 1H), 3.40 - 3.52 (m, 2H), 3.55 - 3.62 (m, 1H), 3.75 (d, J=12 Hz, 1H), 3.79 (s, 3H), 5.93 (dd, J=1 Hz, 3 Hz, 2H), 6.72 (d, J=8Hz, 1H), 6.82-6.90 (m, 3H), 7.03 (d, J=2Hz, 1H), 7.30 (d, J=9Hz, 2H).

Example 35

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-dipentylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J = 7Hz, 6H), 0.95-1.03 (m, 2H), 1.10-1.30 (m, 8H), 1.40-1.51 (m, 2H), 2.72 (d, J=13Hz, 1H), 2.90-3.08 (m, 4H), 3.25-3.50 (m, 3H), 3.37 (d, J=13Hz, 1H), 3.52-3.60 (m, 1H), 3.70 (J=10Hz, 1H), 3.75 (s, 3H), 5.92 (dd, J=2Hz, 5Hz, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.88 (m, 3H), 7.03 (d, J=2Hz, 1H), 7.30 (d, J=9Hz, 2H).

Example 36*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(2-methoxyethyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 120-122 °C. ¹H NMR (CDCl₃, 300 MHz) δ 2.82 (d, J=13, 1H), 2.94-3.08 (m, 2H), 3.12 (s, 3H), 3.23 (s, 3H), 3.20-3.70 (m, 11H), 3.73 (d, J=10Hz, 1H), 3.79 (s, 3H), 5.92 (dd, J=2Hz, 2H, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.90 (m, 3H), 7.04 (d, J=2Hz, 1H), 7.30 (d, J=9Hz, 2H).

Example 37*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-hexynyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 4, 200 mg. of the pure *trans,trans* isomer, the compound resulting from Example 6A was reacted with 109 mg of 1-bromo-2-hexyne, prepared by the method described in Perkin I, 2004 (1987), for 1 hour at 55 °C, to give 226 mg of the intermediate ester. The ester was hydrolyzed using NaOH in ethanol-water for 3 hours at room temperature to give 175 mg of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ 1.00 (t, J=7Hz, 3H), 1.54 (m, 2H), 2.14-2.22 (m, 2H), 2.96 (dd, J=7Hz, 13Hz, 1H), 3.07 (dd, J=18Hz, 2Hz, 1H), 3.15 (dd, J=9Hz, 2Hz, 1H), 3.26 (t, J=9Hz, 1H), 3.36 (dd, J=18 Hz, 2Hz, 1H), 3.47-3.55 (m, 1H), 3.79 (s, 3H), 3.88 (d, J=9Hz, 1H), 5.95 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.88 (m, 3H), 7.03 (d, J=2Hz, 1H), 7.22 (d, J=9Hz, 2H).

Example 38*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-cyclopropylmethyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 167-169 °C. Rotational isomers were seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) δ -0.1 (m), 0.05 (m), 0.12-0.25 (m), 0.32-0.51 (m), 0.67 and 0.74 (2 triplets, 3H), 0.90-1.00 (m), 1.20-1.55 (m), 2.72 (d, J=13Hz, 1H), 2.85--3.29 (m, 4H), 3.30-3.50 (m, 3H), 3.52-3.62 (m, 1H), 3.65-3.73 (2 doublets, J=10Hz, 2Hz, 1H), 3.78 (s, 3H), 5.95 (2 singlets, 2H), 6.72 (2 doublets, 2H), 6.80-6.90 (m, 3H), 7.00 and 7.05 (2 doublets, J=9Hz, 2H).

Example 39*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-pentylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. Rotational isomers were seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (t, J=7Hz, 3H), 1.00-1.08 (m), 1.13-1.32 (m), 1.35-1.50 (m), 2.72-2.82 (2

doublets, J=13Hz, 1H), 2.83 and 2.86 (2 singlets, 3H), 2.92-3.20 (m, 3H), 3.22-3.45 (m, 3H), 3.52-3.62 (m, 1H), 3.72 (2 doublets, 1H), 3.75 and 3.76 (2 singlets, 3H), 5.92 (2 singlets, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.87 (m, 3H), 7.03 (2 doublets, J=2Hz, 1H), 7.30 (d, J=9Hz, 2H).

Example 40

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-diisobutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 141-143 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.54 (d, J=7Hz, 3H), 0.70-0.90 (3 doublets, J=7Hz, 9H), 1.60-1.75 (m, 1H), 1.90-2.02 (m, 1H), 2.67 (d, J=13Hz, 1H), 2.70 (d, J=13Hz, 1H), 2.84 (dd, J=6Hz, 15Hz, 1H), 2.96-3.06 (m, 2H), 3.20 (dd, J=9Hz, 15Hz, 1H), 3.35 (dd, J=2Hz, 10Hz, 1H), 3.44-3.60 (m, 4H), 3.70 (d, J=9Hz, 1H), 3.79 (s, 3H), 5.94 (dd, J=2Hz, 2Hz, 2H), 6.72 (d, J=9Hz, 1H), 6.82-6.90 (m, 3H), 7.03 (d, J=2Hz, 1H), 7.31 (d, J=9Hz, 2H).

Example 41

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(2-propenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. Rotational isomers were seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) δ 2.09 and 2.32 (2 triplets, J=2Hz, 1H), 2.80-3.10 (m, 3H), 2.90 and 2.99 (2 singlets, 3H), 3.35-3.50 (m, 2H), 3.52-3.62 (m, 1H), 3.78 (s, 3H), 4.03 (d, J=13Hz, 1H), 4.00-4.30 (m, 3H), 5.93 (s, 2H), 6.72 (2 doublets, J=8Hz, 1H), 6.80-6.90 (m, 3H), 7.02 and 7.11 (2 doublets, J=2Hz, 1H), 7.30 (2 doublets, J=9Hz, 2H).

Example 42

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(n-hexyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (2 triplets, J=7Hz, 3H), 1.00-1.50 (m, 8H), 2.72-2.82 (2 doublets, J=13Hz, 1H), 2.81 and 2.86 (2 singlets, 3H), 2.92-3.20 (m, 3H), 3.22-3.45 (m, 3H), 3.52-3.62 (m, 1H), 3.72 (2 doublets, 1H), 3.75 and 3.76 (2 singlets 3H), 5.94 (2 singlets, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.87 (m, 3H), 7.03 (2 doublets, J=2Hz, 1H), 7.30 (d, J=9Hz, 1H).

Example 43

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 123-125 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.79 (t, J=7Hz, 3H), 0.85 (t, J=7Hz, 3H), 1.00-1.50 (m, 8H), 2.74 (d, J=13Hz, 1H), 2.90-3.09 (m, 4H), 3.23-3.50 (m, 3H), 3.38 (d, J=13Hz, 1H), 3.52-3.62 (m, 1H), 3.75 (d, J=10 Hz, 1H), 3.78 (s, 3H), 5.93 (dd, J=2Hz, 4Hz), 6.71 (d, J=8Hz, 1H), 6.81-6.89 (m, 3H), 7.03 (d, J=2Hz, 1H), 7.30 (d, J=9 Hz, 2H). MS (DCI/NH₃) m/e 511 (M+H)⁺. Anal calcd for C₂₉H₃₈N₂O₆: C, 68.21; H, 7.50; N, 5.49. Found: C, 68.07; H, 7.47; N, 5.40.

Example 44

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-diethylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 132-134 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.98 (t, J=7Hz, 3H), 1.06 (t, J=7Hz, 3H), 2.78 (d, J=13 Hz, 1H), 2.95-3.20 (m, 4H), 3.30-3.50 (m, 4H), 3.55-3.65 (m, 1H), 3.76 (d, J=12 Hz, 1H), 3.79 (s, 3H), 5.93 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.90 (m, 3H), 7.02 (d, J=2Hz, 1H), 7.32 (d, J=9Hz, 2H).

Example 45

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N*-methyl-*N*-phenylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. ¹H NMR (CD₃OD, 300 MHz) δ 2.75-2.85 (m, 2H), 3.05-3.13 (m, 1H), 3.18 (s, 3H), 3.40-3.58 (m, 2H), 3.78 (s, 3H), 3.88 (d, J=12Hz, 1H), 5.92 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.75-6.85 (m, 3H), 7.00-7.12 (m, 5H), 7.82-7.92 (m, 3H).

Example 46

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N*-methyl-*N*-cyclohexylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. Rotational isomers were seen in the NMR. ¹H NMR (CD₃OD, 300 MHz) δ 1.00-1.85 (m, 10H), 2.72 and 2.78 (2 singlets, 3H), 2.75-2.82 (2 doublets, J=12Hz, 1H), 2.96-3.22 (m, 3H), 3.40-3.65 (m, 3H), 3.68 and 3.82 (2 doublets, J=10Hz, 1H), 3.77 and 3.78 (2 singlets, 3H), 5.92 (s, 2H), 6.72 (2 doublets, J=8Hz, 1H), 6.82-6.88 (m, 3H), 7.02 (2 doublets, J=2Hz, 1H), 7.30-7.40 (2 doublets, J=9Hz, 2H).

Example 47trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-propyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 170-172 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.69 (t, J=7Hz, 3H), 0.85 (t, J=7Hz, 3H), 1.20-1.55 (m, 4H), 2.72 (d, J=13Hz, 1H), 2.90-3.10 (m, 4H), 3.25-3.47 (m, 4H), 3.35-3.62 (m, 1H), 3.72 (d, J=9Hz, 1H), 3.79 (s, 3H), 5.94 (s, 2H), 6.72 (d, d, J=8Hz, 1H), 6.80-6.90 (m, 3H), 7.02 (d, J=2Hz, 1H), 7.30 (d, J=9Hz, 2H).

Example 48trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-isobutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid using the procedures described in Example 1. Rotational isomers were seen in the NMR. ¹H NMR (CD₃OD, 300 MHz) δ 0.65-0.85 (4 doublets, J=7Hz, 6H), 1.75-1.95 (m, 1H), 2.80 and 2.90 (2 singlets, 3H), 2.90-3.10 (m, 4H), 3.10-3.65 (m, 4H), 3.74 9S, 3H), 3.81 and 3.88 (2 doublets, J=10Hz, 1H), 5.93 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.90 (m, 3H), 7.02 (2 doublets, J=2Hz, 1H), 7.80-7.90 (2 doublets, J=9Hz, 2H).

Example 49Alternate Preparation ofEthyl 2-(4-methoxybenzoyl)-4-nitromethyl-3-(1,3-benzodioxole-5-yl)butyrateExample 49AE-2-(3,4-Methylenedioxyphenyl)-1-nitroethene

To a stirred solution of piperonal (75g, 500 mmol) in methanol (120 mL) at 10 °C was added nitromethane (27.1 mL, 500 mmol, 1 eq) followed by the dropwise addition of sodium hydroxide (21 g, 525 mmol, 1.05 eq) in sufficient water to achieve a total volume of 50 mL while maintaining the temperature between 10-15 °C. The reaction mixture became cloudy, turning to a thick paste. The mixture was stirred for 30 minutes upon completion of the addition, and the mixture was then diluted with ice-water (~350 mL) maintaining the temperature below 5 °C, until solution was achieved. The resultant solution was poured in a narrow stream (such that it just failed to break into drops) into a rapidly stirred solution of 36% hydrochloric acid (100 mL) in water (150 mL). A yellow solid precipitated (nitrostyrene), and this was collected by filtration, washed with water (1.5 L) until the filtrate was neutral. The filter cake was air dried and then recrystallized from hot ethanol (3 L) to

yield E-2-(3,4-methylenedioxy)-nitrostyrene as yellow needles (53 g, 55%). ¹H NMR (300MHz, CDCl₃) δ 7.94 (1H, d, J=13.5Hz), 7.47 (1H, d, J=13.5Hz), 7.09 (1H, dd, J=7.5&2Hz), 7.01 (1H, d, J=2Hz), 6.87 (1H, d, J=7.5Hz), 6.06 (2H, s). MS (DCI/NH₃) m/e 194 (M+H)⁺, 211 (M+H+NH₃)⁺.

Example 49B

Ethyl 2-(4-methoxyphenyl)oxo-4-nitro-3-(3,4-methylenedioxyphenyl)butyrate

To a stirred solution of the nitrostyrene resulting from Example 49A (14.17 g, 73.34 mmol, 1.2 eq) in a mixture of propan-2-ol (75 mL) and tetrahydrofuran (175 mL) at room temperature was added successively a solution of ethyl (4-methoxybenzoyl)acetate (11.5 g, 51.7 mmol) in THF (50 mL) followed by 1,8-diazabicyclo[5,4,0]undec-7-ene (DBU) (0.45 mL, 3.0 mmol, 0.05 eq). The resultant mixture was stirred at room temperature for 1 hour, then additional DBU (0.45 mL, 3.0 mmol, 0.05 eq) was added. The mixture was stirred a further 1 hour, then the volatiles were removed *in vacuo* and the residue purified by flash chromatography on 500 g silica gel, eluting with 20% ethyl acetate-hexanes changing to 25% ethyl acetate-hexanes as the product eluted. The solvents were removed *in vacuo* to yield the nitroketoester (19.36 g, 76%) as a viscous oil. Diastereomers were seen in the NMR. ¹H NMR (300 MHz, CDCl₃) δ 8.06 (2H, d, J=9Hz), 7.89 (2H, d, J=9Hz), 6.96 (2H, d, J=9Hz), 6.91 (2H, d, J=9Hz), 6.77 (1H, dd, J=9Hz,3Hz), 6.73 (1H, d, J=9Hz), 6.65 (1H, d, J=3Hz), 5.95 (2H, s), 5.89 (1H, d, J=4Hz), 5.88 (1H, d, J=4Hz), 4.90-4.60 (3H, m), 4.39 (1H, m), 4.18 (2H, q, J=7Hz), 3.94 (2H, m), 3.80 (3H, s), 3.78 (3H, s), 1.19 (3H, t, J=7Hz), 0.99 (3H, t, J=7Hz), MS (DCI/NH₃) m/e 416 (M+H)⁺, 433 (M+H+NH₃)⁺.

Example 50

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(t-butyl)oxycarbonylmethyl-pyrrolidine-3-carboxylic acid

To a stirred solution of the compound resulting from Example 1C (100 mg, 0.27 mmol) in acetonitrile (2 mL) was added successively diisopropylethylamine (70 μL, 0.40 mmol, 1.5 eq) and t-butyl bromoacetate (48 μL, 0.29 mmol, 1.1 eq). The mixture was stirred 2 hours, then the solvent was removed *in vacuo* to yield the crude diester. To a stirred solution of the diester in ethanol (1 mL) at room temperature was added 50% w/w sodium hydroxide (300 mg, 3.75mmol) in water. The mixture was stirred 2 hours, then the volatiles were removed *in vacuo*. The residue was dissolved in water (5 mL), and the solution was washed with ether. The aqueous phase was acidified with acetic acid (300 μL), and then extracted with ethyl acetate (2x). The combined organic extracts were dried (Na₂SO₄),

filtered, and concentrated to yield the title compound (74 mg, 60%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 7.36 (2H, d, $J=8\text{Hz}$), 7.13 (1H, d, $J=3\text{Hz}$), 6.90 (1H, dt, $J=3\text{Hz}$, 8Hz), 6.88 (2H, d, $J=8\text{Hz}$), 6.76 (1H, d, $J=8\text{Hz}$), 5.96 (2H, s), 3.96 (1H, d, $J=9\text{Hz}$), 3.81 (3H, s), 3.58 (1H, ddd, $J=12, 10\text{Hz}, 3\text{Hz}$), 3.52 (1H, dd, $J=9\text{Hz}, 3\text{Hz}$), 3.32 (1H, d, $J=17\text{Hz}$), 3.08 (1H, t, $J=10\text{Hz}$), 2.92 (1H, dd, $J=9\text{Hz}, 7\text{Hz}$), 2.83 (1H, d, $J=17\text{Hz}$). MS (DCI/NH_3) m/e 456 ($\text{M}+\text{H}$) $^+$.

Anal calc for $\text{C}_{29}\text{H}_{29}\text{NO}_7 \cdot 0.3 \text{H}_2\text{O}$: C, 65.07; H, 6.48; N, 3.04. Found: C, 65.02; H, 6.42; N, 2.93.

Example 51

trans,trans-2-(4-Methoxyphenyl)-4-(1-naphthyl)-1-(N-methyl-N-propyl)aminocarbonylmethyl-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting naphthalene-1-carboxaldehyde for piperonyl in Example 49A. Rotational isomers are seen in the NMR. ^1H NMR (300 MHz, CDCl_3) δ 8.29 (1H, bd, $J=8\text{Hz}$), 7.86 (2H, d, $J=8\text{Hz}$), 7.75 (1H, d, $J=8\text{Hz}$), 7.49 (3H, m), 7.34 (2H, dd, $J=3\text{Hz}, 9\text{Hz}$), 6.83 (2H, dd, $J=9\text{Hz}, 2\text{Hz}$), 4.50 (1H, m), 3.94 (1H, dd, $J=9\text{Hz}, 2\text{Hz}$), 3.78 (3H, s), 3.65 (1H, m), 3.49 (1H, d, $J=14\text{Hz}$), 3.40-2.93 (5H, m), 2.91, 2.83 (3H, s), 1.48 (2H, sept, $J=7\text{Hz}$), 0.83, 0.77 (3H, t, $J=7\text{Hz}$). MS (DCI/NH_3) m/e 461 ($\text{M}+\text{H}$) $^+$. Anal calcd for $\text{C}_{29}\text{H}_{29}\text{NO}_7 \cdot 0.5 \text{HOAc}$: C, 71.00; H, 6.99; N, 5.71. Found: C, 70.95; H, 7.00; N, 5.46.

Example 52

trans,trans-2-(4-Methoxyphenyl)-4-(2,3-dihydrobenzofuran-5-yl)-1-(N-methyl-N-propyl)aminocarbonylmethyl-pyrrolidine-3-carboxylic acid

Example 52A

2,3-Dihydrobenzofuran-5-carboxaldehyde

To a stirred solution of α,α -dichloromethyl methyl ether (2.15 g, 19 mmol, 1.35 eq) in methylene chloride (30 mL) at -40°C was added successively stannic chloride (1.65 g, 17 mmol, 1.2 eq) and 15 minutes later, a solution of 2,3-dihydrobenzofuran (1.68 g, 14 mmol) in CH_2Cl_2 (5 mL) maintaining the temperature at or below -35°C . The mixture was warmed to 0°C , stirred 1 hour, then poured into ice-water, and stirred a further 30 minutes. The mixture was diluted with ether, and the phases separated. The organic phase was concentrated *in vacuo*, and the residue purified by vacuum distillation to yield the title compound (1.25 g, 60%) as a colorless liquid. b.p. $119\text{--}121^\circ\text{C}$ at 0.3 mm Hg.

Example 52B*trans,trans*-2-(4-Methoxyphenyl)-4-(2,3-dihydrobenzofuran-5-yl)-1-(N-methyl-N-propyl)aminocarbonylmethyl-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting the compound resulting from Example 52A for piperonal in Example 49A. Rotational isomers are seen in the NMR. ¹H NMR (300 MHz, CDCl₃) δ 7.33 (1H, d, J=8Hz), 7.28 (1H, m), 7.19 (1H, m), 6.87 (1H, d, J=8Hz), 6.73 (1H, d, J=8Hz), 4.56 (1H, t, J=8Hz), 3.83 (1H, d, J=10Hz), 3.80 (3H, s), 3.63 (1H, m), 3.4-3.0 (9H, m), 2.87, 2.84 (3H, s), 1.51 (2H, septet, J=7Hz), 0.88, 0.78 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 453 (M+H)⁺. Anal calc for C₂₆H₃₂N₂O₅ · 0.25 H₂O: C, 68.33; H, 7.17; N, 6.13. Found: C, 68.60; H, 6.88; N, 5.80.

Example 53*trans,trans*-2,4-Bis(4-methoxyphenyl)-1-(N-methyl-N-propyl)aminocarbonylmethyl-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 4-methoxybenzaldehyde for piperonal in Example 49A. Rotational isomers are seen in the NMR. ¹H NMR (300 MHz, CDCl₃) δ 7.37 (2H, d, J=7.5 Hz), 7.32 (2H, d, J=7.5 Hz), 6.86 (4H, m), 3.83 (1H, m), 3.81 (3H, s), 3.79 (3H, s), 3.64 (1H, m), 3.48-2.97 (6H, m), 2.87, 2.83 (3H, s), 2.85 (1H, m), 1.45 (2H, m), 0.84, 0.74 (3H, t, J=7.5 Hz). MS (DCI/NH₃) m/e 441 (M+H)⁺. Anal calc for C₂₅H₃₂N₂O₅ · 0.5 H₂O: C, 66.80; H, 7.40; N, 6.23. Found: C, 67.15; H, 7.31; N, 6.00.

Example 54*trans,trans*-2-(4-Methoxyphenyl)-4-(3,4-dimethoxyphenyl)-1-(N-methyl-N-propyl)aminocarbonylmethyl-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 3,4-dimethoxybenzaldehyde for piperonal in Example 49A. Rotational isomers are seen in the NMR. ¹H NMR (300 MHz, CDCl₃) δ 7.33 (2H, d, J=7.5 Hz), 7.07 (1H, d, J=2.0 Hz), 6.98 (1H, m), 6.85 (1H, d, 7.5 Hz), 6.82 (2H, d, 7.5 Hz), 3.91 (3H, s), 3.86 (3H, s), 3.83 (1H, m), 3.79 (3H, s), 3.64 (1H, m), 3.50-2.95 (6H, m), 2.87 (1H, m), 2.85, 2.83 (3H, s), 1.45 (2H, m), 0.84, 0.74 (3H, t, J=7.5 Hz). MS (DCI/NH₃) m/e 471 (M+H)⁺. Anal calc for C₂₆H₃₄N₂O₆ · 0.5 H₂O: C, 65.12; H, 7.36; N, 5.84. Found: C, 65.22; H, 7.27; N, 5.59.

Example 55*trans,trans*-2-(4-Methoxyphenyl)-4-(3-methoxyphenyl)-1-(N-methyl-N-propyl)aminocarbonylmethyl-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 3-methoxybenzaldehyde for piperonal in Example 49A. Rotational isomers are seen in the NMR. ¹H NMR (300 MHz, CDCl₃) δ 7.33 (2H, d, J=7.5 Hz), 7.24 (1H, t, J=7.5 Hz), 7.05 (2H, m), 6.85 (2H, dd, J=7.5&2 Hz), 6.76 (1H, m), 3.83 (1H, m), 3.81 (3H, s), 3.79 (3H, s), 3.64 (1H, m), 3.48-2.97 (6H, m), 2.87, 2.83 (3H, s), 2.85 (1H, m), 1.45 (2H, m), 0.84, 0.74 (3H, t, J=7.5 Hz). MS (DCI/NH₃) m/e 441 (M+H)⁺. Anal calc for C₂₅H₃₂N₂O₅ · 0.5 H₂O: C, 66.80; H, 7.40; N, 6.23. Found: C, 66.76; H, 7.36; N, 6.05.

Example 56

trans,trans-2-(4-Methoxyphenyl)-4-(2-naphthyl)-1-(N-methyl-N-propyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting naphthylene-2-carboxaldehyde for piperonal in Example 49A. Rotational isomers are seen in the NMR. ¹H NMR (300 MHz, CDCl₃) δ 7.82 (4H, m), 7.69 (1H, m), 7.47 (2H, m), 7.37 (2H, dd, J=7.5&2 Hz), 6.85 (2H, dd, J=7.5&2 Hz), 3.90 (1H, d, J=8 Hz), 3.78 (3H, s), 3.57 (1H, m), 3.52-2.97 (6H, m), 2.93, 2.85 (3H, s), 2.90 (1H, m), 1.52 (2H, m), 0.86, 0.76 (3H, t, J=7.5 Hz). MS (DCI/NH₃) m/e 461 (M+H)⁺. Anal calc for C₂₈H₃₂N₂O₄ · 0.5 H₂O: C, 71.62; H, 7.08; N, 5.97. Found: C, 71.58; H, 7.11; N, 6.01.

Example 57

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(ethylsulfinyl)ethyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 1C (100 mg, 0.27 mmol) and 2-chloroethyl ethyl sulfide (67.5 mg, 0.5 mmol, 2 equivalents) dissolved in 6 mL of acetonitrile was added 10 mg of KI and 0.5 mL of diisopropylethylamine. The mixture was refluxed for 4 hours and then concentrated *in vacuo*. The residue obtained was purified by flash chromatography on silica gel eluting with 4:1 hexane-ethyl acetate to afford 93 mg (75%) of the ethylthioethyl compound.

To the sulfide (90 mg, 0.2 mmol) dissolved in 5 mL of CH₂Cl₂ in an ice bath was added 68 mg of 3-chloroperoxybenzoic acid. The mixture was stirred for 40 minutes in the ice bath and for 3 hours at room temperature. A 10% solution of sodium hydroxide (2 mL) was added, and the mixture was extracted with EtOAc (2 x 50 mL). The combined organic extracts were washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. The residue obtained was chromatographed on silica gel eluting with EtOAc and 10% MeOH in CH₂Cl₂ to afford the sulfoxide (62 mg, 65%).

The ethyl ester was hydrolyzed by the procedure described in Example 1D to afford the title compound as a diastereomeric mixture. m.p. 61-63 °C. MS (DCI/NH₃) m/e 446 (M+H)⁺. ¹H NMR (CDCl₃, 300 MHz) δ 1.25, 1.32 (t, J=9Hz, 3H), 2.45-2.75 (m, 4H), 2.84-2.96 (m, 3H), 3.02-3.08 (m, 1H), 3.32, 3.36 (d, J=3Hz, 1H), 3.47-3.58 (m, 2H), 3.65, 3.68 (d, J=7.5Hz, 1H), 3.76, 3.80 (s, 3H), 5.94 (s, 2H), 6.72 (d, J=7.5Hz, 1H), 3.84-3.89 (m, 3H), 7.02 (d, J=6Hz, 1H), 7.30, 7.34 (d, J=7.5Hz, 2H).

Example 58

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(isopropylsulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

To 2-bromoethylamine hydrobromide (1 mmol) suspended in anhydrous CH₃CN was added 1 equivalent of Et₃N. The mixture was stirred for 30 minutes and then 1 equivalent of isopropyl sulfonyl chloride and 1 equivalent of Et₃N were added. The resulting mixture was stirred for 2 hours at room temperature and then added to a solution of the compound resulting from Example 1C (185 mg, 0.5 mmol) in 3 mL of CH₃CN. The mixture was warmed at 50-60 °C for 2 hours, cooled to room temperature, treated with water and extracted with EtOAc. The combined organic extracts were washed with water and brine, dried and concentrated *in vacuo*. The residue obtained was chromatographed on silica gel eluting with 3:2 hexane-EtOAc to give 195 mg (75%) of the ethyl ester. The ethyl ester (160 mg, 0.31 mmol) was hydrolyzed by the procedure described in Example 1D to afford the title compound (133 mg, 88%). m.p. 94-96 °C. ¹H NMR (CD₃OD, 300 MHz) δ 1.26 (d, J=6Hz, 6H), 1.97 (s, 1H), 2.38 (m, 1H), 2.77 (m, 1H), 2.88 (t, J=9Hz, 1H), 3.04 (m, 1H), 3.14 (t, J=7.5Hz, 2H), 3.35 (m, 2H), 3.46 (m, 1H), 3.58 (m, 1H), 3.78 (s, 3H), 5.92 (s, 2H), 6.74 (d, J=9Hz, 1H), 6.86 (dd, J=9Hz, 3Hz, 1H), 6.92 (d, J=9Hz, 2H), 7.00 (d, J=3Hz, 1H), 7.36 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e (M+H)⁺.

Example 59

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(isobutoxy)ethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Example 1D from the compound resulting from Example 1C and 2-(isobutoxy)ethyl bromide. m.p. 68-70 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.88 (d, J=6Hz, 6H), 1.82 (quintet, J=6Hz, 1H), 2.22 (m, 2H), 2.72-2.79 (m, 1H), 2.86-2.95 (m, 2H), 3.13 (d, J=6Hz, 2H), 3.45-3.56 (m, 4H), 3.68 (d, J=9Hz, 1H), 3.79 (s, 3H), 5.94 (s, 2H), 6.72 (d, J=7.5Hz, 1H), 6.85 (dd, J=9Hz, 7.5 Hz, 3H), 7.08 (s, 1H), 7.34 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 442 (M+H)⁺.

Example 60*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(butylsulfonyl)-pyrrolidine-3-carboxylic acid

To 100 mg (0.271 mmol) of the compound resulting from Example 1C dissolved in 10 mL of THF was added 1-butanefulfonyl chloride (46.7 mg, 1.1 equivalents) and diisopropylethylamine (53 mg, 1.5 equivalents). The resulting mixture was stirred for 2.5 hours at room temperature and then the solvent evaporated. The crude product was purified by flash chromatography on silica gel eluting with 3:2 hexane-EtOAc to afford 120 mg (90%) of the ethyl ester.

The ester (120 mg, 0.244 mmol) was dissolved in 1 mL of EtOH, and a solution of 100 mg of NaOH in 1 mL of water was added. The mixture was stirred for 3 hours at room temperature and then concentrated under reduced pressure. Water (5 mL) was added and the solution was washed with ether to remove any unhydrolyzed *trans-cis* isomer. The aqueous solution was acidified to pH~6 with acetic acid and then extracted with EtOAc (2 x 50 mL). The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated under reduced pressure to afford the pure title compound (60 mg, 53%) as a white solid. m.p. 67-69 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=7.5Hz, 3H), 1.20-1.33 (m, 2H), 1.58-1.68 (m, 2H), 2.48-2.69 (m, 2H), 3.28 (dd, J=9Hz, 1H), 3.49 (t, J=12Hz, 1H), 3.65 (dd, J=12Hz, 1H), 3.82 (s, 3H), 4.32 (dd, J=12Hz, 1H), 5.17 (d, J=9Hz, 2H), 5.95 (s, 2H), 6.70-6.78 (m, 3H), 6.92 (d, J=9Hz, 2H), 7.35 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 462 (M+H)⁺.

Example 61*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-methyl-N-isopropylcarbonylamino)ethyl)-pyrrolidine-3-carboxylic acidExample 61A*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-bromoethyl)-pyrrolidine-3-carboxylic acid ethyl ester

To the mixture of *cis,trans* and *trans,trans* pyrrolidines resulting from Example 1C (400 mg) dissolved in 9 mL of 1,2-dibromoethane was added 0.7 mL of diisopropylethylamine and 30 mg of sodium iodide. The resultant mixture was heated at 100 °C for 1 hour, and then the solvents were removed *in vacuo*. The residue was taken up in EtOAc and washed sequentially with water and brine, dried and concentrated under reduced pressure. The crude product was purified by flash chromatography on silica gel eluting with 4:1 hexane-EtOAc to give 470 mg of the title product.

Example 61B*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(methylamino)ethyl)-pyrrolidine-3-carboxylic acid ethyl ester

To the compound resulting from Example 61A (450 mg) dissolved in 10 mL of EtOH was added 0.5 mL of 40% aqueous methylamine and 50 mg of sodium iodide. The mixture was heated at 80 °C for 1 hour, and then the solvents were removed *in vacuo*. The residue was taken up in EtOAc and washed sequentially with water and brine, dried and concentrated *in vacuo*. The resultant product was carried on without further purification.

Example 61C*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-methyl-N-isobutrylamino)ethyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 61B (~150 mg) dissolved in 5 mL of 1,2-dichloroethane was added 0.3 mL of diisopropylethylamine. The solution was cooled to -40 °C, isobutryl chloride (0.17 mL) was added, the bath was removed, and the solution was allowed to warm to ambient temperature and stirred for 15 hours. The solvent was removed *in vacuo*; the residue was taken up in EtOAc and washed sequentially with 1:1 sodium bicarbonate solution/water and brine, dried and concentrated *in vacuo*. The product was purified by flash chromatography on silica gel eluting with a gradient 1:1 EtOAc-hexanes going to EtOAc and finally using 10% MeOH-EtOAc.

The ester was dissolved in 1.5 mL of EtOH; 0.75 mL of a 17% aqueous NaOH solution was added, and the resultant mixture was stirred at ambient temperature for 3 hours. The solvents were removed *in vacuo*; the residue was taken up in water and washed with ether. The aqueous phase was acidified with 1 N H₃PO₄ to pH 3 and extracted twice with ether. The combined organic extracts were washed with brine and dried over Na₂SO₄. The solvents were removed *in vacuo* to provide 82 mg of the title compound as a white foam. Rotamers were seen in the NMR. ¹H NMR (CDCl₃, 300 MHz) of the major rotamer δ 1.06 (d, 3H, J=10Hz), 1.12 (d, 3H, J=10Hz), 2.15 (m, 1H), 2.5-3.0 (m, 3H), 2.91 (s, 3H), 3.32 (m, 2H), 3.50 (m, 2H), 3.65 (m, 2H), 3.77 (s, 3H), 5.92 (s, 2H), 6.73 (d, 1H, J=8Hz), 6.75-6.9 (m, 4H), 6.96 (d, 1H, J=2Hz), 7.29 (m, 1H). MS (DCI/NH₃) m/z 469 (M+H)⁺. Analysis calcd for C₂₆H₃₂N₂O₆ · 0.3 TFA: C, 63.55; H, 6.48; N, 5.57. Found: C, 63.44; H, 6.71; N, 5.24.

Example 62*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-methyl-N-propionylamino)ethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Example 61 substituting propionyl chloride for isobutryl chloride in Example 61C. ¹H NMR (CDCl₃,

300 MHz) of the major rotamer δ 1.13 (t, 3H, J=8Hz), 2.19 (m, 1H), 2.30 (m, 2H), 2.65-3.0 (m, 3H), 2.85 (s, 3H), 3.25-3.4 (m, 2H), 3.5-3.7 (m, 3H), 3.79 (s, 3H), 5.92 (s, 2H), 6.74 (d, 1H, J=8Hz), 6.75-6.9 (m, 4H), 7.00 (bd s, 1H), 7.29 (bd s, 1H). MS (DCI/NH₃) m/z 455 (M+H)⁺. Analysis calcd for C₂₅H₃₀N₂O₆ · 1.0 H₂O: C, 63.55; H, 6.83; N, 5.93. Found: C, 63.55; H, 6.52; N, 5.73.

Example 63

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-benzylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) of the major rotamer δ 2.79 (s, 3H), 2.8-3.2 (m, 2H), 3.48 (m, 2H), 3.61 (m, 2H), 3.77 (s, 3H), 3.78 (m, 1H), 4.3-4.5 (m, 2H), 5.95 (d, 2H, J=2Hz), 6.7-6.9 (m, 4H), 7.00 (m, 1H), 7.15-7.35 (m, 7H). MS (FAB/NBA) m/z 503 (M+H)⁺. Anal calcd for C₂₉H₃₀N₂O₆ · 0.5 H₂O: C, 68.36; H, 5.74; N, 5.50. Found: C, 68.41; H, 5.74; N, 5.36.

Example 64

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-butylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) of the major rotamer δ 0.88 (t, 3H, J=7Hz), 1.06 (t, 3H, J=7Hz), 1.27 (m, 2H), 1.45 (m, 2H), 2.8-3.6 (m, 11H), 3.79 (s, 3H), 3.80 (m, 1H), 5.92 (bd s, 2H), 6.75 (d, 1H, J=8Hz), 6.85 (d, 1H, J=8Hz), 6.92 (d, 2H, J=8Hz), 7.03 (s, 1H), 7.33 (d, 1H, J=8Hz). MS (DCI/NH₃) m/z 483 (M+H)⁺. Anal calcd for C₂₇H₃₄N₂O₆ · 0.5 HOAc: C, 65.61; H, 7.08; N, 5.46. Found: C, 65.51; H, 6.70; N, 5.66.

Example 65

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(2,2-dimethylpropyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) of the major rotamer δ 0.90 (s, 9H), 2.8-3.1 (m, 4H), 2.94 (s, 3H), 3.3-3.5 (m, 3H), 3.61 (m, 1H), 3.80 (s, 3H), 3.82 (m, 1H), 5.94 (bd s, 2H), 6.74 (d, 1H, J=8Hz), 6.86 (d, 2H, J=8Hz), 6.87 (m, 1H), 7.03 (d, 1H, J=2Hz), 7.33 (d, 2H, J=8Hz). MS (DCI/NH₃) m/z 483 (M+H)⁺.

Example 66

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-methyl-N-butylsulfonfylamino)ethyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 61B (60 mg, 0.13 mmol) dissolved in 5 mL of CH₃CN was added 0.2 mL of Et₃N and 22 mg (0.143 mmol, 1.1 equivalents) of 1-butanethiol. The mixture was stirred for 1 hour at room temperature and then concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel eluting with 1:1 EtOAc-hexane to yield 64 mg (90%) of the ester. Ester hydrolysis by the procedure described in Example 1D afforded the title compound. m.p. 64-66 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.92 (t, J=7.5Hz, 3H), 1.39 (hexad, J=7.5Hz, 2H), 1.68-1.76 (m, 2H), 2.16-2.25 (m, 1H), 2.72 (s, 3H), 2.75-2.92 (m, 5H), 3.12-3.20 (m, 1H), 3.25-3.34 (m, 1H), 3.46-3.55 (m, 2H), 3.65 (d, J=9Hz, 1H), 3.78 (s, 3H), 5.53 (s, 2H), 6.72 (d, J=7.5Hz, 1H), 6.82 (dd, J=7.5Hz, 3Hz, 1H), 6.86 (d, J=9Hz, 2H), 7.02 (d, J=3Hz, 1H), 7.34 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 519 (M+H)⁺.

Example 67

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-methyl-N-propylsulfonamino)ethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Example 66 substituting 1-propanesulfonyl chloride for 1-butanethiol. m.p. 69-70 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.02 (t, J=7.5Hz, 3H), 1.78 (hexad, J=7.5Hz, 2H), 2.18-2.26 (m, 1H), 2.72 (s, 3H), 2.75-2.95 (m, 6H), 3.13-3.22 (m, 1H), 3.25-3.35 (m, 1H), 3.47-3.58 (m, 2H), 3.66 (d, J=9Hz, 1H), 3.80 (s, 3H), 5.96 (s, 2H), 6.74 (d, J=7.5Hz, 1H), 6.84 (d, J=7.5Hz, 3Hz, 1H), 6.87 (d, J=9Hz, 2H), 7.04 (d, J=3Hz, 1H), 7.43 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 505 (M+H)⁺.

Example 68

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(propylsulfonyl)ethyl)-pyrrolidine-3-carboxylic acid

To 1-propanethiol (3.5 g, 46.05 mmol) dissolved in 10 mL of anhydrous THF was added 632 mg (26.32 mmol) of NaH in portions under a nitrogen atmosphere. The mixture was heated at 60-70 °C for 1 hours. To this mixture was added the compound resulting from Example 61A (180 mg, 0.38 mmol) in 2 mL THF. Heating was continued at 60-70 °C for an additional 2 hours, and then the volatiles were removed under reduced pressure. The crude propylthioethyl adduct was purified by flash chromatography on silica gel eluting with 3:2 hexane-EtOAc to give 170 mg (95%).

To a solution of 170 mg (0.36 mmol) of the sulfide and 93 mg (0.8 mmol) of N-methylmorpholine N-oxide (NMO) in a mixture of 20 mL of acetone and 5 mL of H₂O was added a solution of osmium tetroxide (10 mg) in 0.3 mL of t-butanol. The resulting mixture

was stirred overnight at room temperature and then concentrated under reduced pressure. The residue was partitioned between EtOAc and H₂O. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. Flash chromatography afforded 177 mg (98%) of the ethyl ester which was hydrolyzed by the procedures described in Example 1D to afford the title compound. m.p. 73-75 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.04 (t, J=7.5Hz, 3H), 1.78 (hexad, J=7.5Hz, 2H), 2.59-2.66 (m, 1H), 2.84-3.08 (m, 7H), 3.43 (dd, J=9Hz, 3Hz, 1H), 3.53-3.60 (m, 1H), 3.68 (d, J=9Hz, 1H), 3.82 (s, 3H), 5.96 (s, 2H), 6.75 (d, J=7.5Hz, 1H), 6.82 (dd, J=7.5Hz, 3Hz, 1H), 6.88 (d, J=9Hz, 2H), 6.99 (d, J=3Hz, 1H), 7.32 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 476 (M+H)⁺.

Example 69

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-N-(*trans*-5-methylhex-2-enyl)-pyrrolidine-3-carboxylic acid

Example 69A

trans-5-Methylhex-2-enoic acid ethyl ester

Oil dispersion sodium hydride (0.85 g) was washed with hexanes and suspended in THF (20 mL), and the mixture was cooled in an ice bath to 0 °C. Diisopropyl(ethoxycarbonylmethyl) phosphonate (5.0 mL) was added slowly and the mixture stirred for 20 minutes at 0 °C. Isovaleraldehyde (2.0 mL) in THF (5 mL) was added dropwise over five minutes. The ice bath was removed and the mixture stirred for 18 hours at ambient temperature. Saturated ammonium chloride solution (50 mL) was added and the mixture extracted with diethyl ether (3 x 50 mL). The ether extracts were combined, dried with Na₂SO₄, and evaporated to give a colorless oil which was purified by flash chromatography on silica gel eluting with hexanes. The title compound was isolated as a colorless oil (2.1 g).

Example 69B

trans-5-Methylhex-2-en-1-ol

The compound resulting from Example 69A (2.0 g) was dissolved in toluene and cooled to 0 °C in an ice bath. Diisobutylaluminum hydride (1.5 N in toluene, 20 mL) was added dropwise and the solution stirred at 0 °C for two hours. Citric acid solution (25 mL) was added very slowly to the cooled solution. The resulting mixture was stirred for 18 hours at ambient temperature. Diethyl ether (50 mL) was added, the solids removed by filtration and washed with additional ether (2 x 25 mL). The filtrate was extracted with ether (2 x 25 mL). The ether extractions and washings were combined, dried, and evaporated to give a

colorless oil which was purified by flash chromatography on silica gel eluting with 25% EtOAc-hexanes. The title compound was isolated as a colorless oil (1.25 g).

Example 69C

trans-1-Bromo-5-methylhex-2-ene

The compound resulting from Example 69B (1.0 g) was dissolved in diethyl ether and cooled to 0 °C in an ice bath. Phosphorus tribromide (2.5 g, 0.87 mL) was added dropwise and the solution stirred at 0 °C for two hours. The solution was poured onto ice, the layers separated, and the aqueous layer extracted with additional ether (3 x 25 mL). The ether layers were combined, dried, and evaporated to give a colorless oil which was used without further purification (0.95 g).

Example 69D

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-N-(trans-5-methylhex-2-enyl)-pyrrolidine-3-carboxylic acid

The title compound was synthesized using the methods detailed in Example 1D but substituting the compound resulting from Example 69C for N-propyl bromoacetamide. ¹H NMR (CDCl₃, 300 MHz) δ 0.84 (d, 6H, J=8Hz), 1.57 (heptet, 1H, J=8Hz), 1.87 (t, 2H, J=6Hz), 2.60 (dd, 1H, J=8Hz,14Hz), 2.86 (t, 1H, J=10Hz), 2.96 (dd, 1H, J=8Hz,10Hz), 3.20 (dd, 1H, J= 5Hz,14Hz), 3.29 (dd, 1H, J=3Hz,10Hz), 3.50 (m, 1H), 3.70 (d, 1H, J=10Hz), 3.78 (s, 3H), 5.47 (m, 2H), 5.93 (s, 2H), 6.71 (d, 1H, J=8Hz), 6.83 (d, 3H, J=9Hz), 7.05 (s, 1H), 7.32 (d, 2H, J=9Hz). MS (DCI/NH₃) m/e 438 (M+H)⁺. Anal calcd for C₂₆H₃₁NO₅: C, 71.37; H, 7.14; N, 3.20. Found: C, 71.16; H, 7.24; N, 3.17.

Example 70

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-N-(trans-3,5-dimethylhex-2-enyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Example 69 but substituting 4-methyl-2-pentanone for isovaleraldehyde in Example 69A, which gave ~7:1 mixture of trans/cis olefins. The crude product was purified by preparative HPLC (Vydac μC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The desired fractions were lyophilized to give the product (and its diastereomer) as a white solid. ¹H NMR of the major (trans) isomer: (CDCl₃, 300 MHz) δ 0.83 (d, 6H, J=8Hz), 1.56 (s, 3H), 1.74 (m, 1H), 1.92 (d, 2H, J=6Hz), 3.3-3.5 (m, 3H), 3.6-3.8 (m, 4H), 3.78 (s, 3H), 3.9-4.0 (m, 1H), 5.22 (m, 1H), 5.90 (d, 2H, J=12Hz), 6.63 (m, 1H), 6.78 (m, 3H), 6.95 (s, 1H), 7.45 (d, 3H, J=8Hz). MS (DCI/NH₃) m/e 438 (M+H)⁺. Anal calcd for C₂₇H₃₃NO₅ · 1.0 TFA: C, 61.59; H, 6.06; N, 2.48. Found: C, 61.36; H, 6.10; N, 2.34.

Example 71

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-heptylcarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 71A1-Chloro-3-propyl-2-hexanone

To 2-propylpentanoic acid (156.6 μ L, 1.00 mmol) dissolved in anhydrous dichloromethane (2 mL) was added DMF (3 μ L, 4 mole %), and the solution was cooled to 0 °C under a nitrogen atmosphere. To the solution was added oxalyl chloride (94.3 μ L, 1.08 mmol) dropwise over a few minutes. The reaction was stirred 18 hours while warming to ambient temperature. The mixture was cooled to 0 °C and excess ~0.3 M ethereal diazomethane solution was added. The reaction mixture was stirred 18 hours while warming to ambient temperature. The reaction mixture was washed with 1 M aqueous sodium carbonate solution (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was dissolved in ether (2 mL) and cooled to 0 °C under a nitrogen atmosphere. Hydrogen chloride as a 4 N solution in dioxane (275 μ L, 1.10 mmol) was added dropwise over a few minutes. The reaction was stirred 18 hours while warming to ambient temperature. The reaction mixture was concentrated under reduced pressure and the residual oil was used in the next step without further purification.

Example 71B

trans,trans-Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-heptylcarbonylmethyl)-pyrrolidine-3-carboxylate

To the compound resulting from Example 71A (1.00 mmol, maximum theoretical yield) was added a solution of the *trans,trans* ethyl carboxylate from Example 1C (295 mg, 0.80 mmol as a 50 % solution in toluene), diisopropylethylamine (700 μ L, 4.00 mmol) and acetonitrile (4 mL). To the resulting solution was added sodium iodide (12 mg, 10 mole %), and the reaction mixture was stirred 18 hours under a nitrogen atmosphere at ambient temperature. Additional sodium iodide (24 mg, 20 mole %) and acetonitrile (4 mL) were added, and the reaction mixture was heated at 45-50 °C with stirring for 18 hours. The reaction mixture was concentrated under reduced pressure, and the residue was chromatographed on silica gel eluting with 1:9 ethyl acetate-hexane to give 237 mg (46%) of the title compound as a yellow oil.

Example 71C

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-heptylcarbonylmethyl)-pyrrolidine-3-carboxylic acid

To the compound resulting from Example 71B (231 mg, 0.4532 mmol) dissolved in ethanol (10 mL) was added a solution of lithium hydroxide (38 mg, 0.9065 mmol) in water (2.5 mL). The solution was stirred for 18 hours under a nitrogen atmosphere, additional lithium hydroxide (19 mg, 0.4532 mmol) in water (0.5 mL) was added, and stirring was continued 24 hours. The reaction mixture was concentrated under reduced pressure to remove the ethanol, and the aqueous residue was diluted with water (45 mL) and washed with ether (50 mL). The aqueous layer was neutralized with 1 N hydrochloric acid to cloudiness and then 10% aqueous citric acid was added to adjust the pH to ~5. This solution was then extracted with 10% ethanol in chloroform (4 x 25 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC on silica gel eluted with 1:1 ethyl acetate-hexane to give 86 mg (39%) of the title compound as an off white powder. ¹H NMR (CDCl₃, 300 MHz) δ 0.73-0.97 (m, 6H), 1.03-1.33 (m, 6H), 1.36-1.58 (m, 2H), 2.46 (m, 1H), 2.80-2.98 (m, 3H), 3.38-3.64 (m, 3H), 3.75-3.90 (m, 1H), 3.79 (s, 3H), 5.94 (s, 2H), 6.75 (d, 1H), 6.86 (d, 2H), 6.92 (d, 1H), 7.12 (s, 1H), 7.32 (d, 2H). MS (FAB) m/e 482 (M+H)⁺. Anal calcd for C₂₈H₃₅NO₆: C, 69.83; H, 7.32; N, 2.91. Found: C, 69.57; H, 7.41; N, 2.73.

Example 72

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(valeryl)methyl)-pyrrolidine-3-carboxylic acid

Example 72A

1-Chloro-2-hexanone

Using the procedure described in Example 71A and substituting pentanoic acid for 2-propylpentanoic acid afforded the title compound as an oil which was used in the next step without further purification.

Example 72B

trans,trans-Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxole-5-yl)-1-(valeryl)methyl)-pyrrolidine-3-carboxylate

Substituting the compound resulting from Example 72A for 1-chloro-3-propyl-2-hexanone and using the procedure described in Example 71B, except deleting the first addition of sodium iodide, stirring 18 hours at ambient temperature and purifying by silica gel chromatography eluting with 3:17 ethyl acetate-hexane, the title compound 305 mg (65%) was obtained as a yellow oil.

Example 72Ctrans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(valeryl-methyl)-pyrrolidine-3-carboxylic acid

By substituting the compound resulting from Example 72B for *trans,trans*-Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-heptylcarbonylmethyl)-pyrrolidine-3-carboxylate and using the procedure described in Example 71C, except only one solution of lithium hydroxide (81.5 mg, 1.942 mmol) in water (3.5 mL) was added followed by stirring for 18 hours, the title compound 130 mg (46%) was obtained as an off white powder. ¹H NMR (CDCl₃, 300 MHz) δ 0.87 (t, 3H), 1.26 (m, 2H), 1.49 (m, 2H), 2.37 (m, 2H), 2.79-2.98 (m, 3H), 3.31-3.49 (m, 2H), 3.56 (m, 1H), 3.77, 3.79 (d,s, 4H), 5.94 (s, 2H), 6.75 (d, 1H), 6.81-6.93 (m, 3H), 7.09 (d, 1H), 7.33 (d, 2H). MS (FAB) m/e 440 (M+H)⁺. Anal. calcd for C₂₅H₂₉NO₆: C, 68.32; H, 6.65; N, 3.19. Found: C, 67.95; H, 6.64; N, 3.05.

Example 73trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,4-dimethoxybenzyl)-N-methylaminocarbonylmethyl)pyrrolidine-3-carboxylic acidExample 73Atrans,trans- and cis,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((3,4-dimethoxybenzyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid ethyl ester

Using the procedure of Example 1D, paragraph 1, substituting 3,4-dimethoxybenzyl bromoacetamide for dipropyl bromoacetamide, the desired product mixture was obtained as a white foam in 81% yield.

Example 73Btrans,trans- and cis,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,4-dimethoxybenzyl)-N-methylaminocarbonylmethyl)pyrrolidine-3-carboxylic acid ethyl ester

The resultant product from Example 73A (220 mg, 0.404 mmol) was dissolved in 2 mL dry THF and added dropwise to a stirred, cooled (0 °C) suspension of sodium hydride (23 mg of a 60% by weight mineral oil suspension, 16.5 mg, 0.69 mmol) in 0.2 mL THF, under an argon atmosphere. The resulting mixture was stirred at 0 °C for 1 hour, then methyl iodide (28 μL, 64 mg, 0.45 mmol) was added. The reaction mixture was stirred at 0 °C for 45 minutes. TLC (Et₂O) indicated incomplete reaction. An additional portion of methyl iodide

(28 μ L, 64 mg, 0.45 mmol) and dry 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)pyrimidinone (50 μ L, 0.41 mmol) were added. The reaction mixture was stirred at ambient temperature for 2 days. The reaction was poured into 25 mL of 0.5 M aqueous citric acid and extracted with 2 x 25 mL EtOAc. The combined organic extracts were washed sequentially with 30 mL water and 30 mL brine, then dried (Na_2SO_4), filtered and concentrated under reduced pressure to produce 270 mg of crude material. Flash chromatography on silica gel eluting with Et_2O gave the title compounds as an inseparable mixture in 43% yield. ^1H NMR (CDCl_3 , 300 MHz) δ 2.79 (s) and 2.81 (s), for the N-CH_3 signals. MS m/z 591 ($\text{M}+\text{H}$) $^+$.

Example 73C

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,4-dimethoxybenzyl)-N-methylaminocarbonylmethyl)pyrrolidine-3-carboxylic acid

To the resultant compound from Example 73B (98 mg, 0.17 mmol) dissolved in 1 mL EtOH and cooled to 0 $^\circ\text{C}$ was added a solution of lithium hydroxide monohydroxide (17 mg, 0.41 mmol) in 0.5 mL H_2O . The resulting solution was stirred under a nitrogen atmosphere for 16 hours. The solution was concentrated *in vacuo*, and the residue was partitioned between 15 mL H_2O and 15 mL Et_2O . The aqueous phase was extracted with 5 mL Et_2O , then the aqueous phase was acidified with 10% aqueous citric acid. The acidic aqueous phase was saturated with NaCl and extracted with 3 x 15 mL EtOAc. The EtOAc extracts were combined, dried (Na_2SO_4), then filtered and concentrated *in vacuo* to give 40 mg (42%) of the title compound as a white foam. ^1H NMR (CD_3OD , 300 MHz, two rotameric forms) δ 2.85 (s, 3H), 2.94-3.25 (br m, 3H), 3.35-3.70 (br m) and 3.64 (s, 4 H total), 3.70-3.97 (br m), 3.74 (s), 3.76 (s), 3.78 (s), 3.79 (s), 3.81 (s), and 4.03 (br d, $J=14$ Hz, 8H total), 4.43 (AB, 1H), 5.91 (s) and 5.93 (s, 2H total), 6.50-6.60 (m, 1H), 6.67-7.02 (br m, 6H), 7.29 (br d) and 7.35 (br d, 2H total). HRMS calcd for $\text{C}_{31}\text{H}_{35}\text{N}_2\text{O}_8$ ($\text{M}+\text{H}$) $^+$: 563.2393. Found: 563.2385.

Example 74

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,4-dimethoxybenzyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The procedure of Example 73C was used, with the substitution of the resultant compound from Example 73A for the resultant compound from Example 73B, to provide the title compound. ^1H NMR (CD_3OD , 300 MHz) δ 2.85 (d, $J=16$ Hz, 1H), 2.92 (br t, $J=9$ Hz, 1H), 2.98 (br t, $J=10$ Hz, 1H), 3.32-3.39 (br m, 2H), 3.54-3.65 (br m, 1H), 3.67 (s, 3H), 3.78 (s, 3H), 3.80 (s, 3H), 3.85 (d, $J=10$ Hz, 1H), 4.21 (d, $J=15$ Hz, 1H), 4.41 (d, $J = 15$ Hz, 1H), 5.91 (s, 2H), 6.67 (d, $J=8$ Hz, 1H), 6.75-6.95 (m, 7H), 7.33-7.40 (m, 2H). HRMS calcd for $\text{C}_{30}\text{H}_{32}\text{N}_2\text{O}_8$ ($\text{M}+\text{H}$) $^+$: 549.2237. Found: 549.2224.

Example 75

(2R,3R,4R)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1R)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid

Example 75A

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1R)-1-(benzyloxycarbonyl)butyl)pyrrolidine-3-carboxylic acid ethyl ester

The procedure of Fung, et. al., J. Med. Chem., 35(10): 1722-34 (1992) was adapted. The resultant compound from Example 6A (103 mg, 0.279 mmol) was dissolved in 0.7 mL of nitromethane and 0.7 mL of H₂O, and ammonium carbonate (34 mg, 0.35 mmol) and (2S)-benzyl 2-bromopentanoate (78 mg, 0.30 mmol) were added. The reaction was refluxed for 24 hours. The reaction was partitioned between 15 mL of 1 M aqueous Na₂CO₃ and 25 mL of CH₂Cl₂. The aqueous phase was extracted with 2 x 10 mL CH₂Cl₂, and the combined organic phases were washed with 15 mL brine, dried (Na₂SO₄), then filtered and concentrated under reduced pressure to a brown oil (169 mg). The crude product was purified by silica gel chromatography eluting with 3:1 CH₂Cl₂-hexane to produce 106 mg (68%) of the title compound as a waxy solid. ¹H NMR indicated the presence of two diastereomeric products.

Example 75B

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1R)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid ethyl ester

The resultant compound from Example 75A (101 mg, 0.180 mmol) and 30 mg of 10% palladium on charcoal were stirred in 2 mL EtOAc under 1 atmosphere of H₂ for 4 hours. The reaction mixture was filtered through a plug of Celite, using 15 mL MeOH to wash the catalyst. The combined filtrate and wash were concentrated *in vacuo* to give 81.4 mg (96%) of the crude acid as a white solid.

The above crude acid was combined with HOBt hydrate (41 mg, 0.27 mmol), dipropylamine (26 mg, 0.26 mmol), and 4-methylmorpholine (37 mg, 0.37 mmol) in 2 mL dry DMF. The solution was cooled to -15 °C, then 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (44 mg, 0.23 mmol) was added. The mixture was stirred at -15 °C and allowed to warm slowly to room temperature overnight. The solvent was removed by distillation under reduced pressure, and the residue was partitioned between 20 mL EtOAc and 10 mL of 1 M aqueous Na₂CO₃. The organic phase was washed with 10 mL of brine, dried (Na₂SO₄), then filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel, eluting with 1:2 Et₂O-hexane. Further purification of overlap fractions by preparative TLC eluting with 1:2 Et₂O-

hexane yielded 32 mg (34%) of a less polar product, and 44 mg (46%) of a more polar product.

Example 75C

(2R,3R,4R)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1R)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid

The procedure of Example 73C was followed, with the substitution of the less polar isomer from Example 75B for the resultant product from Example 73B, to provide the title compound in 94% yield. $[\alpha]_D = -52^\circ$ ($c=0.235$, CH_3OH). $^1\text{H NMR}$ (CD_3OD , 300 MHz)

δ 0.55 (t, $J=7\text{Hz}$, 3H), 0.87 (t, $J=7\text{Hz}$) and 0.87-0.94 (m, 6H total), 1.03-1.25 (br m, 2H), 1.25-1.68 (br m, 4H), 1.90-2.07 (br m, 1H), 2.75-2.94 (br m, 2H), 2.94-3.02 (br m, 2H), 3.20-3.40 (m, overlapping with CD_2HOD signal), 3.40-3.60 (br m, 2H), 3.79 (s, 3H), 4.04 (br d, $J=9\text{ Hz}$, 1H), 5.92 (dd, $J=3,5\text{ Hz}$, 2H), 6.72 (d, $J=8\text{ Hz}$, 1H), 6.79 (dd, $J=1.5,8\text{ Hz}$, 1H), 6.92-6.98 (br m, 3H), 7.29-7.39 (m, 2H). MS m/z 525 ($\text{M}+\text{H}$) $^+$.

Example 76

(2S,3S,4S)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1R)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid

The procedure of Example 73C was followed, with the substitution of the more polar isomer from Example 75B for the resultant product from Example 73B, to provide the title compound in 88% yield. $[\alpha]_D = +58^\circ$ ($c=0.37$, CH_3OH). $^1\text{H NMR}$ (CD_3OD , 300 MHz)

δ 0.57 (br t, $J=7\text{Hz}$, 3H), 0.88-0.98 (m, 6H), 1.08-1.35 (br m, 2H), 1.35-1.68 (br m, 4H), 1.75-1.90 (br m, 1H), 2.75-2.86 (br m, 2H), 3.10-3.30 (br m, 2H), 3.51-3.65 (br m, 2H), 3.69 (s, 3H), 4.03-4.16 (br m, 2H), 5.91 (s, 2H), 6.71-6.83 (m, 2H), 6.86-6.97 (m, 3H), 7.32 (br d, $J=9\text{Hz}$, 2H). MS m/z 525 ($\text{M}+\text{H}$) $^+$.

Example 77

(2S,3S,4S)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1S)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid

Example 77A

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1S)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid ethyl ester

(2R)-N,N-dipropyl 2-hydroxypentanamide (106 mg, 0.528 mmol, made by standard procedure) was dissolved in 2 mL THF under an argon atmosphere, diisopropylethylamine (75 mg, 0.58 mmol) was added, then the solution was cooled to -20°C .

Trifluoromethanesulfonic anhydride (95 μL , 159 mg, 0.565 mmol) was added to the cooled

solution over 1 minute, and the reaction mixture was stirred at -20 °C for 1 hour, and at room temperature for an additional 1 hour. The resulting slurry was recooled to 0 °C, and a solution of the resultant compound from Example 6A (195 mg, 0.528 mmol) and diisopropylethylamine (101 μ L, 75 mg, 0.58 mmol) in 3 mL of CH₂Cl₂ was added. The reaction was stirred at 0 °C for 3 hours and for an additional 2 days at room temperature. TLC (Et₂O-hexane 1:2) indicated starting materials remained, so the mixture was warmed to reflux for 4 hours. The reaction was cooled, then partitioned between 30 mL EtOAc and 15 mL of 1 M aqueous Na₂CO₃. The aqueous phase was extracted with 15 mL EtOAc, then the combined organic phases were washed with 20 mL brine, dried (Na₂SO₄), filtered and concentrated *in vacuo* to a yellowish oil. Purification by flash chromatography on silica gel eluting with 1:2 Et₂O-hexane gave 19.9 mg (7%) of a less polar product and 20.1 mg (7%) of a more polar product. ¹H NMR spectra and MS were the same as those of Example 76B.

Example 77B

(2S,3S,4S)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1S)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid

The procedure of Example 73C was followed, with the substitution of the less polar isomer from Example 77A for the resultant product from Example 73B, to provide the title compound in 100% yield. ¹H NMR (CD₃OD, 300 MHz) and MS identical to those of Example 75C.

Example 78

(2R,3R,4R)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((1S)-1-(N,N-dipropylaminocarbonyl)-1-butyl)pyrrolidine-3-carboxylic acid

The procedure of Example 73C was followed, with the substitution of the more polar isomer from Example 77A for the resultant product from Example 73B, to provide the title compound in 88% yield. ¹H NMR (CD₃OD, 300 MHz) and MS identical to those of Example 76.

Example 79

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-3-(5-tetrazolyl)pyrrolidine

Carbonyldiimidazole (510 mg, 3.148 mmol) was added to 1.020 g (2.00 mmol) of the compound resulting from Example 43 in 2.7 mL THF, and the mixture was heated for 40 minutes at 50 °C. The reaction mixture was cooled in an ice bath, and 25% solution of ammonia in methanol was added. After 30 minutes, the solid which had formed was filtered,

washed with ethanol and finally with ether to yield 850 mg (83%) of the 3-carboxamide compound. m.p. 194-196 °C.

Phosphorus oxychloride (1.06 g) was added to this amide in 7 mL of pyridine, and the mixture was stirred 1 hour at room temperature. Dichloromethane was added, and the solution was washed with potassium bicarbonate solution, dried over sodium sulfate, and concentrated. The residue was chromatographed on silica gel eluting with 2:1 hexane-ethyl acetate to give 790 mg (96%) of the 3-carbonitrile compound.

To this nitrile in 5 mL toluene was added 385 mg of trimethyl tin chloride and 126 mg sodium azide. The mixture was heated 20 hours at 125 °C (bath temp). After cooling, methanol (5 mL) was added, and the solution was concentrated *in vacuo*. To the resulting residue was added 6 mL of methanol and 6 mL of water containing 0.2 g phosphoric acid. After stirring 1 hour at room temperature, water was added and the mixture extracted with dichloromethane. The combined organic extracts were dried and concentrated, and the resulting residue was crystallized from ether to give a solid. The solid was dissolved in sodium hydroxide solution, filtered from insoluble material and acidified with acetic acid to get 532 mg (62%) of the title compound. m.p. 165-167 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (t, J=7Hz, 3H), 0.87 (t, J=7Hz, 3H), 1.10-1.50 (m, 8H), 3.0-3.6 (m, 8H), 3.70 (s, 3H), 3.7-3.8 (m, 1H), 3.90 (t, J=9Hz, 1H), 4.37 (d, J=9Hz, 1H), 5.86 (s, 2H), 6.62 (d, J=8Hz, 1H), 6.65-6.73 (m, 3H), 6.95 (d, J=2Hz, 1H), 7.11 (d, J=9Hz, 2H).

Example 80

trans,trans-2-(4-Fluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound was prepared as an amorphous solid from methyl (4-fluorobenzoyl) acetate and 5-(2-nitrovinyl)-1,3-benzodioxole using the procedures described in Examples 1 and 43. ¹H NMR (CDCl₃, 300 MHz) δ 0.81 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.0-1.55 (m, 8H), 2.81 (d, J=13 Hz, 1H), 2.90-3.10 (m, 4H), 3.15-3.30 (m, 1H), 3.32-3.45 (m, 3H), 3.55-3.65 (m, 1H), 3.86 (d, J=10Hz, 1H), 5.94 (dd, J=2Hz, 4Hz, 2H), 6.72 (d, J=8 Hz, 1H), 6.86 (d, J= 8 Hz, 1H), 6.95-7.07 (m, 3H), 7.32-7.45 (m, 2H).

Example 81

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

N,N-Dibutyl glycine (150 mg, 0.813 mmol), prepared by the method of Bowman, R.E., J. Chem. Soc. 1346 (1950), in 0.7 mL of THF was treated with 138 mg (0.852 mmol) carbonyldiimidazole and heated for 30 minutes at 50 °C. After cooling to room temperature, 250 mg (0.678 mmol) of ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-

pyrrolidine-3-carboxylate, the compound resulting from Example 6A, was added, and the mixture was heated at 45 °C for 30 minutes. The product was chromatographed on silica gel, eluting with 1:1 hexane-ethyl acetate to give 306 mg of the intermediate ethyl ester.

The ester was hydrolyzed with sodium hydroxide in water and ethanol to give 265 mg of the title compound as a white powder. ¹H NMR (CDCl₃, 300 MHz) δ rotational isomers - 0.75 and 0.85 (2 t, J=7Hz, 3H), 1.05-1.5 (m, 8H), 2.65-3.20 (m, 6H) 3.43-3.70 (m, 3H), 3.72 (s, 3H), 3.87 (d, J=15Hz, 1H), 4.49 (dd, J=12Hz, 6Hz) and 5.23 (dd, J=12Hz, 8Hz) 2H, 5.90 (dd, J=2Hz, 4Hz, 2H), 6.63-6.78 (m, 3H), 6.86 and 7.04 (d, J=9Hz, 2H), 7.22 (d, J=9Hz, 2H).

Example 82

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-n-butyl)-N-(n-propyl)aminocarbonylmethylpyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Example 1. m.p. 160-162 °C. ¹H NMR (CDCl₃, 300 MHz) rotational isomers δ 0.69, 0.80, 0.84, 0.87 (four triplets, J=7Hz, 6H), 1.00-1.52 (m, 6H), 2.63 and 2.66 (two doublets, J=13Hz, 1H), 2.90-3.10 (m, 4H), 3.23-3.61 (m, 5H), 3.71 and 3.75 (two doublets, J=10Hz, 1H), 3.78 (s, 3H), 5.92-5.96 (m, 2H), 6.72 (d, J=8Hz, 1H), 6.83-6.89 (m, 3H), 7.03 (d, J=2Hz, 1H), 7.81 (d, J=9Hz, 2H).

Example 83

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N,N-di(n-propyl)aminocarbonyl)ethyl]pyrrolidine-3-carboxylic acid

The compound resulting from Example 6A (250 mg, 0.677 mmol), 205 mg (1.36 mmol) diallyl acrylamide (Polysciences, Inc.), and 10 mg acetic acid were heated at 85 °C in 0.75 mL of methoxyethanol for one hour. Toluene was added, and the solution was washed with bicarbonate solution, dried, and concentrated. Chromatography on silica gel eluting with 3:1 hexane-ethyl acetate gave 283 mg (80%) of the diallyl compound.

The diallyl compound was hydrogenated using 10% Pd/C catalyst (27 mg) in ethyl acetate (25 mL) under a hydrogen atmosphere. The catalyst was removed by filtration, and the filtrate was concentrated to afford the dipropyl amide ethyl ester in 100% yield.

The ester was hydrolyzed to the title compound by the method of Example 1D in 83% yield. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 and 0.83 (two triplets, J=7Hz, 6H), 1.39-1.54 (m, 4H), 2.35-2.60 (m, 3H), 2.80-3.07 (m, 5H), 3.14-3.21 (m, 2H), 3.31-3.38 (m, 1H), 3.51-3.61 (m, 1H), 3.73 (d, J=12Hz, 1H), 3.75 (s, 3H), 5.94 (s, 2H), 6.71 (d, J=9Hz, 1H), 6.79-6.85 (m, 3H), 7.04 (d, J=2Hz, 1H) < 7.32 (d, J=9Hz, 2H).

Example 84*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Example 8 using
 5 dibutyl carbamoyl chloride, prepared by the method of Hoshino *et al.*, Syn. Comm., 17: 1887-1892 (1987), as a starting material. ¹H NMR (CDCl₃, 300 MHz) δ 0.86 (t, J=7Hz, 6H), 1.14-1.28 (m, 4H), 1.35-1.48 (m, 4H), 2.81-2.94 (m, 2H), 3.11 (t, J=12Hz, 1H), 3.30-3.41 (m, 2H), 3.59-3.68 (m, 2H), 3.76 (s, 3H), 3.78-3.85 (m, 1H), 5.81 (d, J=9Hz, 1H), 5.94 (s, 2H), 6.73-6.86 (m, 5H), 7.24 (d, J=9Hz, 2H).

Example 85*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid sodium salt

Sodium hydroxide (48.2 mg of 98.3% pure, 1.184 mmol) in 2 mL of MeOH was
 15 added to the compound resulting from Example 43 (610 mg, 1.196 mmol.) in 5 mL MeOH. The solution was concentrated to dryness, and the resulting powder was stirred with heptane. The heptane was removed *in vacuo* to give a powder which was dried in the vacuum oven for 2 hours at 60 °C to yield 627.5 mg of the title compound.

Example 86*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(*N,N*-di(*n*-butyl)amino)ethyl]pyrrolidine-3-carboxylic acid

A solution of the bromoethyl compound resulting from Example 61A (150 mg),
 dibutylamine (150 mg) and sodium iodide (18 mg) in 0.75 mL ethanol was heated at 80 °C for
 25 1 hour. After cooling, toluene was added, and the solution was washed with potassium bicarbonate solution, dried over Na₂SO₄ and concentrated. More toluene was added, and the solution was again concentrated to get rid of excess dibutylamine. The residue was dissolved in warm heptane and filtered from a small amount of insoluble material. The heptane was removed *in vacuo* to give 143 mg (87%) of the intermediate ethyl ester.

The ester was hydrolyzed by the method of Example 1D to give the title compound as
 30 a white powder. ¹H NMR (CD₃OD, 300 MHz) δ 0.89 (t, J=7Hz, 6H), 1.16-1.30 (m, 4H), 1.44-1.56 (m, 4H), 2.48-2.57 (m, 1H), 2.80-3.08 (m, 8H), 3.14-3.25 (m, 1H), 3.31-3.38 (m, 1H), 3.59-3.60 (m, 1H), 3.74 (s, 3H), 3.75 (d, J=10Hz, 1H), 5.89 (s, 2H), 6.71 (d, J=9Hz, 1H), 6.81 (dd, J=9Hz, 2Hz, 1H), 6.90 (d, J=10Hz, 2H), 6.96 (d, J=2Hz, 1H), 7.37 (d, J=10Hz, 2H).

Example 87

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-[N-(N,N-di(n-butyl)aminocarbonyl)-N-methylamino]ethyl]pyrrolidine-3-carboxylic acid

Dibutyl carbamoyl chloride (135 mg) was added to the compound resulting from Example 61B (250 mg) and 150 mg triethylamine in 1 mL dichloromethane. After stirring 1 hour at room temperature, toluene was added, and the solution was washed with potassium bicarbonate solution, dried over Na₂SO₄ and concentrated. The residue was chromatographed on silica gel, eluting with a mixture of 38% EtOAc and 62% hexane to give 194 mg of the ethyl ester intermediate.

The ester was hydrolyzed by the method of Example 1D to afford 141 mg of the title compound. ¹H NMR (CD₃OD, 300 MHz) δ 0.92 (t, J=7Hz, 6H), 1.21-1.32 (m, 4H), 1.42-1.53 (m, 4H), 2.62 (s, 3H), 2.65-2.76 (m, 1H), 3.00-3.20 (m, 8H), 3.44-3.55 (m, 1H), 3.62-3.78 (m, 2H), 3.80 (s, 3H), 4.07 (d, J=12 Hz, 1H), 5.93 (s, 2H), 6.75 (d, J=9Hz, 1H), 6.87 (dd, J=9Hz, 2Hz, 1H), 6.94 (d, J=10 Hz, 2H), 7.04 (d, J=2Hz, 1H), 7.40 (d, J=10Hz, 2H).

Example 88

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-(N-methanesulfonyl)carboxamide

Carbonyldiimidazole (75 mg, 0.463 mmol) was added to 150 mg (0.294 mmol) of the compound resulting from Example 43 in 0.4 mL of tetrahydrofuran, and the solution was stirred at 60 °C for 2 hours. After cooling, 50 mg (0.526 mmol) of methanesulfonamide and 68 mg (0.447 mmol) of DBU in 0.3 mL of THF were added. The mixture was stirred at 45 °C for 2 hours. The solvents were removed *in vacuo*, and the residue was dissolved in water. A few drops of acetic acid were added, and the solution was lyophilized to give 121 mg (70%) of the title compound. m.p. 170-173 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.05-1.51 (m, 8H), 2.75-2.86 (m, 2H), 2.83-3.25 (m, 4H), 3.17 (s, 3H), 3.32-3.50 (m, 3H), 3.70-3.78 (m, 1H), 3.80 (s, 3H), 3.87 (d, J=10Hz, 1H), 5.96 (dd, J=2Hz, 4Hz, 2H), 6.74 (d, J=9Hz, 1H), 6.84 (dd, J=9Hz, 2Hz, 1H), 6.90 (d, J=10 Hz, 2H), 7.01 (d, J=2Hz, 1H), 7.34 (d, J=10Hz, 2H).

Example 89

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-(N-benzenesulfonyl)carboxamide

The compound resulting from Example 43 was converted to the title compound by the method of Example 88 substituting benzenesulfonamide for methanesulfonamide. m.p. 169-171 °C for a sample recrystallized from acetonitrile. ¹H NMR (CDCl₃, 300 MHz) δ 0.81(t, J=7 Hz, 3H), 0.89 (t, J=7Hz, 3H), 1.02-1.50 (m, 8H), 2.65-2.80 (m, 2H), 2.90-3.25 (m, 4H), 3.80-3.95 (m, 3H), 3.50-3.60 (m, 1H), 3.65 (d, J=10Hz, 1H), 3.81 (s, 3H), 5.94 (s, 2H), 6.70

(s, 2H), 6.81-6.90 (m, 3H), 7.17 (d, J=10Hz, 2H), 7.55 (t, J=7 Hz, 2H), 7.66 (t, J=7Hz, 1H), 8.95 (d, J=7Hz, 2H).

Example 90

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminosulfonylmethyl]-pyrrolidine-3-carboxylic acid

Chloromethyl sulfonyl chloride, prepared by the method of Brintzinger et. al., Chem. Ber. 85: 455-457 (1952), is reacted with dibutylamine by the method of E. Vilsmaier described in Liebigs Ann. Chem. 1055-1063 (1980) to give N,N-dibutyl chloromethyl sulfonyl chloride. Alternatively dimethyl(methylthio)sulfonium tetrafluoroborate is reacted with dibutylamine to give N,N-dibutyl methylsulfonyl chloride which is chlorinated with N-chlorosuccinimide to give chloromethyl sulfonyl chloride by the method of E. Vilsmaier, described in the above reference.

The N,N-dibutyl chloromethyl sulfonyl chloride is reacted with the compound resulting from Example 6A to give ethyl trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminosulfonylmethyl]-pyrrolidine-3-carboxylate. This is oxidized with osmium tetroxide and N-methyl morpholine N-oxide by the method of S. Kaldor and M. Hammond, Tet. Lett. 32: 5043-5045 (1991) to give the title compound after hydrolysis of the ethyl ester.

Example 91

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminocarbonyl-1-(RS)-ethyl]pyrrolidine-3-carboxylic acid

Example 91A

(±)-Dibutyl 2-bromopropanamide

2-Bromopropanoic acid (510 mg, 3.33 mmol) and 4-methylmorpholine (0.74 mL, 6.73 mmol) were dissolved in 10 mL of CH₂Cl₂, the solution was cooled to 0 °C under a N₂ atmosphere, and then treated dropwise with isobutyl chloroformate (0.45 mL, 3.5 mmol). After 10 minutes at 0 °C, dibutylamine (0.57 mL, 3.4 mmol) was added. The reaction was stirred at 0 °C for 1 hour and for an additional 16 hours at room temperature. The mixture was partitioned with 25 mL of 1.0 M aqueous Na₂CO₃ solution, then the organic phase was washed sequentially with 25 mL of 1 M aqueous NaHSO₄ and 25 mL brine, dried (Na₂SO₄), filtered, and concentrated under reduced pressure to afford 698 mg (2.64 mmol, 79 %) of the crude bromoamide as a colorless oil. ¹H NMR (CDCl₃, 300 MHz) δ 0.93 (t, J=7Hz) and 0.97 (t, J=7.5Hz, 6H

total), 1.26-1.60 (m, 7H), 1.60-1.78 (m, 1H), 1.82 (d, J=6Hz, 3H), 3.04-3.27 (m, 2H), 3.42-3.64 (m, 2H), 4.54 (q, J=7H, 1H). MS (DCI/NH₃) m/e 264 and 266 (M+H)⁺.

Example 91B

5 *trans,trans*- and *cis,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N,N-di(n-butyl)amino)carbonyl-1-(*RS*)-ethyl)pyrrolidine-3-carboxylic acid ethyl ester

A solution of the resultant mixture of *trans,trans* and *cis,trans* compounds from Example 1C (232 mg, 0.628 mmol) and the resultant compound from Example 91A (183 mg, 0.693 mmol) in 2 mL of CH₃CN was treated with diisopropylethylamine (0.22 mL, 1.3 mmol). The solution was stirred at 60-80 °C under a N₂ atmosphere for 16 hours. The reaction was concentrated under reduced pressure, then the residue was partitioned between 30 mL Et₂O and 10 mL of 1 M aqueous Na₂CO₃ solution. The organic phase was washed with 20 mL water and 20 mL brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude amino amide as a brown oil (339 mg, 98% crude). The product was obtained by flash chromatography on silica gel eluting with 20% EtOAc-hexane to provide 224 mg (70%) of the title compounds as a mixture of 4 diastereomers. ¹H NMR (CDCl₃, 300 MHz) δ 0.66-1.55 (several m, 19H), 2.63-3.00 (m, 3H), 3.05-3.39 (m, 2H), 3.40-3.76 (m, 4H), 3.78-3.80 (4 s, 3H), 3.84-4.25 (m, 2,6H), 4.38 (d, J=10.5Hz, 0,2H) and 4.58 (d, J=10.5Hz, 0,2H), 5.90-5.97 (m, 2H), 6.68-6.96 (m, 5H), 7.38-7.43 (m, 2H). MS (DCI/NH₃) m/e 553 (M+H)⁺.

Example 91C

25 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N,N-dibutylamino)carbonyl-1-(*RS*)-ethyl)pyrrolidine-3-carboxylic acid

The procedure of Example 73C was used, substituting the resultant compound from Example 91B for the resultant compound from Example 73B to give the title compound in 61% yield. ¹H NMR (CD₃OD, 300 MHz) δ 0.70-1.05 (several m, 8H), 1.14 (d, J=6Hz, 2H), 1.17-1.55 (m, 6H), 2.79-3.03 (m, 3,5H), 3.20-3.65 (br m, 4,6H plus CD₂HOD), 3.70-3.78 (m, 0,4H), 3.79 (s, 3H), 3.98 (d, J=8Hz, 0,6H), 4.06 (t, J=7.5Hz, 0,4H), 4.25 (d, J=8Hz, 0,4H), 5.92 (s) and 5.94 (s, 2H total 6H), 6.73 (d, J=2.5Hz) and 6.75 (d, J=3Hz, 1H total), 6.78-6.85 (m, 1H), 6.91-7.00 (m, 3H), 7.30-7.38 (m, 2H). MS (DCI/NH₃) m/e 525 (M+H)⁺. Anal calcd for C₃₀H₄₀N₂O₆·0.5H₂O: C, 67.52; H, 7.74; N, 5.25. Found: C, 67.63; H, 7.65; N, 5.21.

Example 92

trans,trans-2-(Pentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Example 92A

Methyl 2-(4-hexenoyl)-4-nitro-3-(1,3-benzodioxole-5-yl)butyrate

A solution of methyl 3-oxo-6-octenoate (502 mg, 2.95 mmol) in 10 mL of isopropanol was added to a solution of 5-(2-nitrovinyl)-1,3-benzodioxole (712 mg, 3.69 mmol) in 10 mL THF, then DBU (22 μ L, 0.15 mmol) was added. The resulting reddish solution was stirred at room temperature for 20 minutes. TLC (ethyl acetate-hexane, 1:3) indicated complete consumption of ketoester. The solution was concentrated *in vacuo* and flash chromatographed on silica gel eluting with 18% ethyl acetate in hexane to produce 879 mg (2.42 mmol, 82%) of the title compound as a mixture of diastereomers in a 1:1 ratio. ^1H NMR (CDCl_3 , 300 MHz) δ 1.55-1.66 (m, 3H), 2.02-2.17 (br m, 1H), 2.20-2.37 (m, 1.5H), 2.49-2.76 (m, 1.5H), 3.57 (s, 1.5H), 3.74 (s, 1.5H), 3.97 (d, $J=7.5\text{H}$, 0.5H) and 4.05 (d, $J=8\text{Hz}$, 0.5H), 4.10-4.20 (m, 1H), 4.68-4.82 (m, 2H), 5.06-5.52 (m, 2H), 5.95 (2s, 2H), 6.65 (m, 1H), 6.68 (br s, 1H), 6.75 (d, 7.5Hz, 1H). MS (DCI/NH_3) m/e 381 ($\text{M}+\text{NH}_4$) $^+$. Anal calcd for $\text{C}_{18}\text{H}_{21}\text{NO}_7$: C, 59.50; H, 5.82; N, 3.85. Found: C, 59.32; H, 5.71; N, 3.72.

Example 92B

Methyl *trans,trans*-2-(pentyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylate

The procedures of Example 1B and Example 1C were followed, with the substitution of the resultant compound from Example 92A for the resultant compound from Example 1A, and the substitution of the this resultant compound for the resultant compound from Example 1B, to provide the title compound in crude form as a yellow oil. This crude compound was epimerized under the following conditions. A solution of the crude compound (660 mg, 2.07 mmol) in 3 mL methanol was treated with a solution of sodium methoxide (made by the addition of sodium metal (14 mg, 0.61 mmol) to 1 mL of methanol). The resultant solution was heated at reflux for 18 hours. The reaction was concentrated under reduced pressure, and the residue was partitioned between 25 mL saturated NaHCO_3 diluted with 10 mL water and 30 mL of CH_2Cl_2 . The aqueous phase was extracted (2 x 30 mL CH_2Cl_2), then the combined organic phases were washed with 20 mL brine, dried over Na_2SO_4 , filtered and the filtrate concentrated under reduced pressure to afford the crude product. Purification by flash chromatography on silica gel eluting with 3.5% methanol in CH_2Cl_2 gave 336 mg (57%) the title compound as a yellow oil. ^1H NMR (CDCl_3 , 300 MHz) δ 0.90 (br t, 3H), 1.25-1.70 (br m, 8H), 1.83-2.02 (br s, 2H), 2.58 (dd, $J=8.9\text{Hz}$, 1H), 2.99 (dd,

J=8,14Hz, 1H), 3.34-3.45 (m, 2H), 3.53 (q, J=9Hz, 1H), 3.66 (s, 3H), 5.94 (s, 2H), 6.65-6.75 (m, 3H). MS (DCI/NH₃) m/e 320 (M+H)⁺. Anal calcd for C₁₈H₂₅N₄O₄: C, 67.69; H, 7.89; N, 4.39. Found: C, 67.39; H, 7.84; N, 4.37.

5

Example 92C

trans,trans-2-(Pentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The procedures of Example 1B-1D were used, with the substitution of the resultant compound from Example 92A for the resultant compound from Example 1B, to provide the title compound as a white foam. ¹H NMR (CDCl₃, 300 MHz) δ 0.87 (br t) and 0.89 (br t, 6H total), 0.97 (t, J=7.5Hz, 3H), 1.21-1.42 (br m, 10), 1.43-1.78 (br m, 6H), 2.76 (t, J=7Hz, 1H), 3.02-3.30 (br m, 6H), 3.40-3.60 (m, 3H), 3.73 (d, J=14Hz, 1H), 5.98 (AB, 2H), 6.70 (d, J=7Hz, 1H), 6.77 (dd, J=1.5,7Hz, 1H), 6.89 (d, J=1.5Hz, 1H). MS (DCI/NH₃) m/e 475 (M+H)⁺. Anal calcd for C₂₇H₄₂N₂O₅·0.5H₂O: C, 67.05; H, 8.96; N, 5.79. Found: C, 67.30; H, 8.77; N, 5.68.

15

Example 93

trans,trans-2-(Pentyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-propylsulfonylamino)ethyl]pyrrolidine-3-carboxylic acid

20

Example 93A

Methyl trans,trans-2-(pentyl)-4-(1,3-benzodioxol-5-yl)-1-(2-bromoethyl)pyrrolidine-3-carboxylate

The procedure of Example 61A was used, with the substitution of the resultant compound from Example 92B for the resultant compound from Example 1C, to provide the title compound as a yellow oil. ¹H NMR (CDCl₃, 300 MHz) δ 0.89 (br t, J=7Hz, 3H), 1.24-1.40 (br m, 6H), 1.60-1.80 (br m, 2H), 2.61-2.75 (m, 2H), 2.76-2.91 (m, 2H), 3.10-3.22 (m, 2H), 3.36-3.47 (m, 2H), 3.68 (s, 3H), 5.92 (s, 2H), 6.69-6.77 (m, 2H), 6.90-6.94 (m, 1H). MS (DCI/NH₃) m/e 426, 428 (M+H)⁺.

30

Example 93B

Methyl trans,trans-2-(Pentyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-propylsulfonylamino)ethyl]pyrrolidine-3-carboxylate

A solution of the resultant compound from Example 93A (102 mg, 0.24 mmol) and tetrabutylammonium iodide (6 mg, 16 μmol) in 1 mL EtOH was treated with propylamine (60 μL, 0.73 mmol). The solution was warmed to 80 °C for 4 hours. The reaction was concentrated under reduced pressure, then the residue was dissolved

35

in 35 mL ethyl acetate and extracted with 2 x 15 mL of 1 M aqueous Na₂CO₃. The organic phase was washed with 15 mL brine, then dried over Na₂SO₄, filtered and concentrated under reduced pressure to provide the crude secondary amine as a yellow oil (94.2 mg). The crude amine was dissolved in 1 mL of CH₂Cl₂,

diisopropylethylamine (65 μ L, 0.373 mmol) was added, followed by propylsulfonyl chloride (29 μ L, 0.26 mmol). The solution was stirred at room temperature for 4 hours. The reaction was quenched with 10% aqueous citric acid (to pH 4), and the mixture was extracted with 2 x 3 mL CH₂Cl₂. The combined organic extracts were washed with 2 mL brine, then dried over Na₂SO₄, filtered, concentrated *in vacuo*.

Purification by flash chromatography eluting with 20% ethyl acetate in hexane provided 65.0 mg (53%) of the title compound as a waxy solid. R_f = 0.17 (20%EtOAc-hexane). MS (DCI/NH₃) m/e 511 (M+H)⁺.

Example 93C

trans,trans-2-(Pentyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-propylsulfonylamino)ethyl]pyrrolidine-3-carboxylic acid

The procedure of Example 71C was followed, with the substitution of the resultant compound from Example 93B for the resultant compound from Example 71B, to provide the title compound as a white foam (47 mg, 80%), R_f = 0.14 (5%MeOH-CH₂Cl₂). ¹H NMR (CDCl₃, 300 MHz) δ 0.88 (br t) and 0.92 (t, J=7Hz, 6H total), 1.22-1.52 (br m, 6H), 1.63 (sextet, J=8Hz, 2H), 1.75-2.10 (br m, 4H), 2.89-2.98 (m, 2H), 3.05 (br t, J=9Hz, 1H), 3.10-3.30 (m, 3H), 3.30-3.80 (br m, 7H), 5.94 (s, 2H), 6.71 (t, J=8Hz, 1H), 6.77 (dd, J=1.5,8Hz, 1H), 6.89 (d, J=1.5Hz, 1H). MS (DCI/NH₃) m/e 497 (M+H)⁺.

Example 94

trans,trans-2-(Propyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Example 94A

Ethyl 2-(4-butanoyl)-4-nitro-3-(1,3-benzodioxole-5-yl)butyrate

The procedure of Example 92A was followed, with the substitution of ethyl butyryl acetate for methyl 3-oxo-6-octenoate, to provide the title compound as a mixture of *trans* and *cis* isomers (47 mg, 80%), R_f = 0.28 (25%EtOAc-hexane). ¹H NMR (CDCl₃, 300 MHz) δ 0.74 (t, J=7.5Hz) and 0.91 (t, J=7.5Hz, 3H total), 1.08 (t, J=7Hz) and 1.28 (t, J=7Hz, 3H total), 1.45 (sextet, J=7Hz, 1.5H), 1.63 (sextet, J=7Hz, approx. 1.5H), 2.17 (t, J=7Hz) and 2.24 (t, J=7Hz, 0.5H total) 2.40-2.54 (m, 1H), 2.60

(t, J=7.5Hz) and 2.67 (t, J=7.5Hz, 0.5H total), 3.93-4.09 (m, 2H), 4.10-4.20 (br m, 1H), 4.23 (q, J=7Hz, 1H), 4.67-4.85 9m, 2H), 5.94 (s, 2H), 6.62-6.75 (m, 3H). MS (DCI/NH₃) m/e 369 (M+NH₄)⁺. Anal calcd for C₁₇H₂₁N₃O₇: C, 58.11; H, 6.02; N, 3.99. Found: C, 58.21; H, 5.98; N, 3.81.

5

Example 94B

Ethyl *trans,trans*-2-(propyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylate

The procedure of Example 92B was followed, with the substitution of the resultant compound from Example 94A for the resultant compound from Example 92A, to afford the title compound. MS (DCI/NH₃) m/e 306 (M+H)⁺.

10

Example 94C

trans,trans-2-(Propyl)-4-(1,3-benzodioxol-5-yl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The procedure of Example 92C was followed, with the substitution of the resultant product from Example 94B for the resultant product from Example 92B, to give the title compound. ¹H NMR (CDCl₃, 300 MHz) δ 0.89 (t, J=7.5Hz), 0.92 (t, J=7.5Hz), and 0.97 (t, J=7.5H, 9H total), 1.22-1.80 (br m, 12H), 2.83 (t, J=7.5Hz, 1H), 3.40-3.55 (br m, 2H), 3.55-3.68 (m, 1H), 3.78 (d, J=15Hz, 1H), 5.92 (q, J=1Hz, 2H), 6.70 (d, J=8Hz, 1H), 6.79 (dd, J=1Hz, 8Hz, 1H), 6.90 (d, J=1Hz, 1H). MS (DCI/NH₃) m/e 447 (M+H)⁺. Anal calcd for C₂₅H₃₈N₂O₅·0.5 H₂O: C, 65.91; H, 8.63; N, 6.15. Found: C, 65.91; H, 8.68; N, 5.94.

15

20

Example 95

(2R,3R,4S)-(+)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(tert-butyloxycarbonyl-aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

25

Example 95A

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((tert-butyloxycarbonylaminocarbonylmethyl)pyrrolidine-3-carboxylic acid

30

The resulting mixture of 64% *trans,trans*- and *cis,trans*- pyrrolidines resulting from Example 1C (3.01 g, 8.15 mmol) was dissolved in 50 mL of methylene chloride. To this was added dropwise a solution of di-tert-butyl dicarbonate (1.96 g, 8.97 mmol) in 20 mL methylene chloride under a nitrogen atmosphere, and the resulting solution was stirred 30 minutes at which point TLC (ethyl acetate:hexane, 1:1) indicated that all of the starting material was consumed. The reaction mixture was concentrated and dried under high vacuum to give 3.94 g of the ethyl ester as a yellow-brown oil. ¹H NMR (CDCl₃, 300 MHz) δ 0.99,

35

1.07 (br t, br t, J=7 Hz, 3H), 1.11-1.62 (several br m, 9H), 3.05 (br m, 1H), 3.44-3.95 (m, 3H), 3.81 (s, 3H), 4.04 (q, J=7 Hz, 1H), 4.14-4.28 (br m, 1H), 4.89-5.24 (br m, 1H), 5.94 (d, J=3 Hz, 2H), 6.69-6.90 (m, 5H), 7.06-7.20 (m, 2H). MS (DCI/NH₃) m/e 470 (M+H)⁺.

To the ethyl ester dissolved in 170 mL of ethanol was added a solution of lithium hydroxide (1.06 g, 25.17 mmol) in 60 mL of water. The reaction mixture was vigorously stirred for 18 hours under a nitrogen atmosphere. The reaction mixture was concentrated to remove ethanol, diluted with 250 mL of water and extracted three times with 250 mL of ether. The organic phase acidified to slight cloudiness (pH ~7) with 1 N hydrochloric acid, then to pH 4 with 10 % citric acid and extracted with 5 % ethanol in methylene chloride (3 x 100 mL). The combined organic layers dried (Na₂SO₄), filtered, concentrated and dried on high vacuum to give the title compound as a white foam (2.19 g, 60 %). ¹H NMR (CDCl₃, 300 MHz) δ 1.16 (v br s, 9H), 3.11 (br m, 1H), 3.50-3.64 (m, 2H), 3.81 (s, 3H), 4.24 (br m, 1H), 4.96 (br m, 1H), 5.94 (s, 2H), 6.71-6.79 (m, 3H), 6.84-6.91 (m, 2H), 7.19 (d, J=9 Hz, 2H). MS (DCI/NH₃) m/e 442 (M+H)⁺.

Example 95B

(2R,3R,4S)-(+)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(tert-butyloxycarbonylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound resulting from Example 95A (2.15 g, 4.86 mmol) and (+)-cinchonine (1.43 g, 4.86 mmol) were added to 100 mL of methylene chloride; this suspension was swirled with warming as necessary to get all solids to dissolve. The solution was then concentrated and dried on high vacuum to a white foam. This material was crystallized from a mixture of refluxing chloroform (64 mL) and hexane (360 mL). The resulting crystals were isolated by filtration and recrystallized under the same conditions seven additional times. Each time the resulting crystals and filtrate were monitored by ¹H NMR and chiral HPLC. The amount of (2S,3S,4R)-(-)- enantiomer decreased first in the crystals and then in the filtrate with the predetermined endpoint achieved when the (2S,3S,4R)-(-)- enantiomer could no longer be detected in the filtrate. The pure (2R,3R,4S)-(+)- enantiomer thus obtained was partitioned between 100 mL of 10% citric acid and 100 mL of ether. The aqueous layer was further extracted twice with 100 mL of ether. The combined ether layers were washed with brine, dried (Na₂SO₄), filtered, concentrated and dried on high vacuum to a white powder (550 mg, 55 % of theoretical 50 % maximum, >99.5 ee). ¹H NMR (CDCl₃, 300 MHz) δ 1.05-1.50 (br m, 9H), 3.12 (br m, 1H), 3.50-3.65 (m, 2H), 3.81 (s, 3H), 4.24 (m, 1H), 4.96 (br m, 1H), 5.95 (s, 2H), 6.70-6.79 (m, 3H), 6.86 (d, J=9 Hz, 2H), 7.19 (d, J=9 Hz, 2H). MS (DCI/NH₃) m/e 442 (M+H)⁺.

Example 95C

(2R,3R,4S)-(+)-Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The compound resulting from Example 95B (251 mg, 0.568 mmol) was dissolved in 20 mL of a saturated solution of anhydrous HCl(g) in anhydrous ethanol. The resulting solution was heated at 50 °C. with stirring for 18 hours at which point all of the precipitated solid had dissolved. The reaction mixture was concentrated to a solid which was partitioned between 0.8 M aqueous sodium carbonate (50 mL) and methylene chloride (50 mL). The aqueous layer was further extracted with methylene chloride (2 x 50 mL). The combined organic layers were dried (Na₂SO₄), filtered, concentrated and dried under high vacuum to give the title compound as an almost colorless oil (158 mg, 69%). ¹H NMR (CDCl₃, 300MHz) δ 1.11 (t, J=7 Hz, 3H), 2.18 (v br s, 1H), 2.93 (t, J= 9 Hz, 1H), 3.19,3.22 (dd, J=7 Hz, 1H), 3.50-3.69 (m, 2H), 3.80 (s, 3H), 4.07 (q, J=7 Hz, 2H), 4.49 (d, J=9 Hz, 1H), 5.94 (s, 2H), 6.73 (d, J=2 Hz, 2H), 6.81-6.92 (m, 3H), 7.34-7.41 (m, 2H). MS (DCI/NH₃) m/e 370 (M+H)⁺.

Example 95D

(2R,3R,4S)-(+)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(tert-butyloxycarbonyl-aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

To the resulting compound from Example 95C (131 mg, 0.355 mmol) was added, diisopropylethylamine (137 mg, 185 μL, 1.06 mmol), acetonitrile (2 mL), N,N-di-(n-butyl)bromoacetamide (133 mg, 0.531 mmol), and the mixture was heated at 50 °C. for 1.5 hours. The reaction mixture was concentrated to a solid, dried under high vacuum, and purified by chromatography on silica gel eluting with 1:3 ethyl acetate-hexane to give pure ester as a colorless oil. ¹H NMR (CDCl₃, 300MHz) δ 0.81 (t, J=7 Hz, 3H), 0.88 (t, J=7 Hz, 3H), 1.10 (t, J=7 Hz, 3H), 1.00-1.52 (m, 8H), 2.78 (d, J=14 Hz, 1H), 2.89-3.10 (m, 4H), 3.23-3.61 (m, 5H), 3.71 (d, J=9 Hz, 1H), 3.80 (s, 3H), 4.04 (q, J=7 Hz, 2H), 5.94 (dd, J=1.5 Hz, 2H), 6.74 (d, J=9 Hz, 1H), 6.83-6.90 (m, 3H), 7.03 (d, J=2 Hz, 1H), 7.30 (d, J=9 Hz, 2H). MS (DCI/NH₃) m/e 539 (M+H)⁺.

To the ethyl ester dissolved in 7 mL of ethanol was added a solution of lithium hydroxide (45 mg, 1.06 mmol) in water (2.5 mL). The mixture was stirred for 1 hour at ambient temperature and then warmed slowly to 40 °C. over 2.5 hours at which point all of the starting material had been consumed. The reaction mixture was concentrated to remove the ethanol, diluted with 60 mL water and extracted with ether (3 x 40 mL). The aqueous solution was treated with 1 N aqueous hydrochloric acid until cloudy, and the pH was then adjusted to ~4-5 with 10% aqueous citric acid. This mixture was extracted with 1:19 ethanol-methylene chloride (3 x 50 mL). The combined extracts were dried (Na₂SO₄), filtered, concentrated and dried under high vacuum to give the title compound as a white foam (150

mg, 83%). ¹H NMR (CDCl₃, 300MHz) δ 0.80 (t, J=7 Hz, 3H), 0.88 (t, J=7 Hz, 3H), 1.08 (m, 2H), 1.28 (m, 3H), 1.44 (m, 3H), 2.70-3.77 (svr br m, 12H), 3.79 (s, 3H), 5.95 (m, 2H), 6.75 (d, J=8 Hz, 1H), 6.87 (br d, J=8 Hz, 3H), 7.05 (br s, 1H), 7.33 (v br s, 2H). MS (DCI/NH₃) m/e 511 (M+H)⁺. [α]_D²² = +74.42°. Anal calcd for C₂₉H₃₈N₂O₆ · 0.5 H₂O: C ,67.03; H, 7.56; N, 5.39. Found: C, 67.03; H, 7.59; N, 5.33.

Example 95E

Alternate Preparation of (2R,3R,4S)-(+)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(tert-butyloxycarbonylamino)carbonylmethyl-pyrrolidine-3-carboxylic acid

The product of Example 95A (2.858 g) was suspended in 10 mL of EtOAc. 0.7833 g of R (+) alpha methyl benzylamine in 3 mL ethyl acetate was added. On swirling all of the solids were dissolved. The ethyl acetate was removed in vacuum. Ether (13 ml) was added to the residue. When all of the residue had dissolved, 5 mg of seed crystals were added and these crystals were crushed with a metal spatula while cooling in ice. The product crystallized very slowly. After 1 hour the solid was filtered and washed with ether giving 1.4213 g, m.p. 163-167°. The filtrate was concentrated, cooled and scratched with a spatula to give a second crop 0.1313 g, m.p. 164-168°. The filtrate was concentrated again and put in the refrigerator and let stand overnight giving 1.6906 g, m.p. 102-110°. (HPLC of this showed 20% of the desired enantiomer and 80% of the unwanted enantiomer.)

The first two batches of crystallized material were combined and suspended in 20 mL dichloromethane (Note: the unwanted isomer is more soluble in dichloromethane) and stirred for 2 minutes. The mixture was concentrated, but not to dryness, and ether (10 mL) was added. After stirring for a few minutes the crystals were filtered. Yield: 1.401 g, m.p. 164-172°.

Treatment of the crystalline product with 10% citric acid and ether according to the method described in Example 95B provided the title compound.

Example 96

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-butyrylamino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and butyryl chloride for isobutyryl chloride in Example 61C. The product was purified by preparative HPLC (Vydac μC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (m, 3H), 0.90 (t, 3H, J=8Hz), 1.42 (m, 2H), 1.58 (heptet, 2H, J=8Hz), 2.20 (t, 3H, J=8Hz), 2.94 (br m,

2H), 3.10 (br m, 2H), 3.48 (br m, 4H), 3.76 (br m, 2H), 3.78 (s, 3H), 4.30 (br s, 1H), 5.95 (s, 2H), 6.75 (d, 1H, J=8Hz), 6.84 (m, 1H), 6.85 (d, 2H, J=8Hz), 7.04 (d, 1H, J=1Hz), 7.40 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 497 (M+H)⁺. Anal calcd for C₂₈H₃₆N₂O₆ · 1.0 TFA: C, 58.82; H, 6.42; N, 4.57. Found: C, 58.77; H, 6.30; N, 4.42.

Example 97

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(ethylaminocarbonyl)amino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and ethyl isocyanate for isobutyl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) mixture of rotamers δ 0.80 (t, J=8Hz) and 1.05 (t, J=8Hz) and 1.20 (m) and 1.42 (m) total of 8H for the four peaks, 2.35 (br s, 1H), 2.70 (m, 1H), 3.0 (m, 3H), 3.2 (m, 3H), 3.25 (dq, 1H, J=1,8Hz), 3.42 (m, 1H), 3.6 (m, 1H), 3.75 (m, 1H), 3.78 (s, 3H), 4.8 (br s, 1H), 5.95 (s, 2H), 6.74 (d, 1H, J=8Hz), 6.85 (m, 3H), 7.00 (s, 1H), 7.30 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 498 (M+H)⁺. Anal calcd for C₂₇H₃₅N₃O₆ · 0.75 H₂O: C, 63.45; H, 7.20; N, 8.22. Found: C, 63.38; H, 7.29; N, 8.44.

Example 98

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-butyl-N-butyrylamino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting butylamine for methylamine in Example 61B and butyryl chloride for isobutyl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (m, 3H), 0.90 (t, 3H, J=8Hz), 1.45 (m, 4H), 1.6 (m, 2H), 2.20 (t, 3H, J=8Hz), 2.94 (br m, 2H), 3.10 (br m, 2H), 3.5 (br m, 4H), 3.80 (br m, 2H), 3.82 (s, 3H), 4.30 (br s, 1H), 5.95 (s, 2H), 6.75 (d, 1H, J=8Hz), 6.84 (m, 1H), 6.85 (d, 2H, J=8Hz), 7.04 (d, 1H, J=1Hz), 7.40 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 511 (M+H)⁺. HRMS calcd for C₂₉H₃₈N₂O₆: 511.2808. Found: 511.2809

Example 99

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-ethoxycarbonylamino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and ethyl chloroformate for isobutyl chloride in Example 61C. The crude product was purified by trituration with 1:1

diethyl ether-hexane. The resulting solid was dissolved in CH_3CN and water and lyophilized to give the product as a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 0.80 (t, 3H, $J=8\text{Hz}$), 1.05 (m, 2H), 1.22 (m, 3H), 1.45 (m, 3H), 2.08 (br s, 1H), 2.75 (m, 1H), 2.88 (br q, 2H, $J=8\text{Hz}$), 3.08 (br m, 2H), 3.27 (br m, 2H), 3.44 (m, 1H), 3.54 (dt, 1H, $J=1.8\text{Hz}$), 3.63 (d, 1H, $J=8\text{Hz}$), 3.78 (s, 3H), 4.02 (br d, 2H), 5.93 (s, 2H), 6.72 (d, 1H, $J=8\text{Hz}$), 6.81 (dd, 1H, $J=1.8\text{Hz}$), 6.85 (d, 2H, $J=8\text{Hz}$), 7.00 (s, 1H), 7.30 (d, 2H, $J=8\text{Hz}$). MS (DCI/NH_3) m/e 499 ($\text{M}+\text{H}$) $^+$. Anal calcd for $\text{C}_{27}\text{H}_{34}\text{N}_2\text{O}_7 \cdot 0.5 \text{H}_2\text{O}$: C, 63.89; H, 6.95; N, 5.52. Found: C, 64.03; H, 6.71; N, 5.30.

5

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000

Example 100*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-methyl-N-(2-ethylbutyryl)amino)ethyl]pyrrolidine-3-carboxylic acid

To the compound resulting from Example 61B (190 mg) dissolved in THF (2 mL) was added HOBt (60 mg), EDCI (85 mg), N-methylmorpholine (50 μ L), and DMF (2 mL). 2-Ethylbutyric acid was added and the solution stirred overnight at ambient temperature. Water (10 mL) was added, and the mixture was extracted with EtOAc (2 x 25 mL). The combined organic extracts were washed with saturated sodium bicarbonate solution, 1 N H_3PO_4 , and brine, dried with Na_2SO_4 , and evaporated to give an oil which was purified by flash chromatography on silica gel eluting with 1:3 EtOAc-hexane. The resulting ethyl ester was saponified by the procedure described in Example 61C. The crude product was dissolved in CH_3CN and water and lyophilized to give the product as a white solid. ^1H NMR (CDCl_3 , 300 MHz) (mixture of rotamers) δ 0.66, 0.74, 0.80, 0.88 (all triplets, total of 6H, $J=8\text{Hz}$), 1.05 (m, 2H), 1.25-1.75 (m, 5H), 2.16 (m, 1H), 2.32 (m, 1H), 2.45 (m, 1H), 2.70 (m, 1H), 2.86, 2.94 (s, total 3H), 2.95 (m, 1H), 3.35 (m, 1H), 3.52 (m, 2H), 3.65 (m, 1H), 3.80 (s, 3H), 5.94, 5.96 (s, total 2H), 6.73 (m, 1H), 6.84 (m, 3H), 6.97 (m, 1H), 7.30 (m, 2H). MS (DCI/NH_3) m/e 497 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{28}\text{H}_{36}\text{N}_2\text{O}_6 \cdot 0.25 \text{H}_2\text{O}$: C, 67.11; H, 7.34; N, 5.59. Found: C, 67.13; H, 7.24; N, 5.56.

Example 101*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-methyl-N-(2-propylvaleryl)amino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedure described in Example 100, but substituting 2-propylpentanoic acid for 2-ethylbutyric acid. The crude product was purified by preparative HPLC (Vydac $\mu\text{C}18$) eluting with a 10-70% gradient of CH_3CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 0.79 (t, 3H, $J=8\text{Hz}$), 0.82 (t, 3H, $J=8\text{Hz}$), 1.10 (m, 4H), 1.2-1.5 (m, 4H), 2.55 (m, 1H), 2.96 (s, 3H), 3.15 (br m, 1H), 3.32 (br m, 1H), 3.56 (m, 2H), 3.68 (m, 1H), 3.68 (s, 3H), 3.70 (m, 1H), 3.80 (m, 2H), 4.65 (br d, 1H), 5.92 (s, 2H), 6.75 (d, 1H, $J=8\text{Hz}$), 6.84 (m, 1H), 6.85 (d, 2H, $J=8\text{Hz}$), 7.05 (s, 1H), 7.42 (d, 2H, $J=8\text{Hz}$). MS (DCI/NH_3) m/e 525 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{30}\text{H}_{40}\text{N}_2\text{O}_6 \cdot 1.25 \text{TFA}$: C, 58.51; H, 6.23; N, 4.20. Found: C, 58.52; H, 6.28; N, 4.33.

Example 102*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(tert-butyloxycarbonylmethyl)amino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and t-butyl bromoacetate for isobutyryl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, 3H, J=8Hz), 1.18 (m, 2H), 1.19 (s, 9H), 2.12 (m, 1H), 2.46 (m, 2H), 2.70 (m, 3H), 2.85 (m, 2H), 3.20 (s, 2H), 3.40 (dd, 1H, J=2,8Hz), 3.50 (dt, 1H, J=2,8Hz), 3.62 (d, 1H, J=8Hz), 3.78 (s, 3H), 5.95 (s, 2H), 6.72 (d, 1H, J=8Hz), 6.84 (m, 1H), 6.85 (d, 2H, J=8Hz), 7.05 (s, 1H), 7.16 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 541 (M+H)⁺. Anal calcd for C₃₀H₄₀N₂O₇ · 1.0 H₂O: C, 64.50; H, 7.58; N, 5.01. Found: C, 64.75; H, 7.35; N, 4.86.

Example 103

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(n-propylaminocarbonylmethyl)amino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and N-propyl bromoacetamide for isobutyryl chloride in Example 61C. The crude product was purified by preparative HPLC (Vydac μC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.78 (t, 3H, J=8Hz), 0.88 (t, 3H, J=8Hz), 1.45 (m, 2H), 1.48 (m, 3H, J=8Hz), 2.55-2.7 (m, 2H), 2.90 (m, 1H), 3.04 (m, 1H), 3.15 (m, 3H), 3.28 (t, 1H, J=8Hz), 3.45 (t, 1H, J=8Hz), 3.60 (m, 2H), 3.70 (d, 2H, J=8Hz), 3.75 (m, 1H), 3.80 (s, 3H), 4.25 (d, 1H, J=8Hz), 5.95 (s, 2H), 6.75(d, 1H, J=8Hz), 6.86 (dt, 1H, J=1,8Hz), 6.88 (d, 2H, J=8Hz), 7.04 (d, 1H, J=1Hz), 7.40 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 526 (M+H)⁺. Anal calcd for C₂₉H₃₉N₃O₆ · 1.85 TFA: C, 53.32; H, 5.59; N, 5.70. Found: C, 53.45; H, 5.62; N, 5.63.

Example 104

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(4-methoxyphenoxy carbonyl)amino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and 4-methoxyphenylchloroformate for isobutyryl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CD₃OD, 300 MHz) mixture of rotamers δ 0.88 (m, 3H), 1.57 (m, 2H), 2.45 (br s) and 2.60 (br s, total of 1H), 2.90-3.15 (m, 4H), 3.42-3.7 (m, 5H), 3.78 (s, 3H), 3.80 (s, 3H), 3.85 (m) and 4.0 (m, total of 1H), 5.95 (s) and 5.98 (s, total of 2H), 6.63(m, 1H), 6.72 (d, 1H, J=8Hz), 6.81 (m,

2H), 6.93 (m, 5H), 7.40 (m, 2H). MS (DCI/NH₃) m/e 577 (M+H)⁺. Anal calcd for C₃₂H₃₆N₂O₈ · 1.0 H₂O: C, 64.63; H, 6.44; N, 4.71. Found: C, 64.70; H, 6.38; N, 4.63.

Example 105

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(4-methoxybenzoyl)amino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and anisoyl chloride for isobutyryl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) mixture of rotamers δ 0.78 (m) and 0.98 (t, J=8Hz) total of 3H, 1.47 (m) and 1.52 (q, J=8Hz) total of 2H, 2.25 (br s, 1H), 2.78 (br s, 1H), 2.90 (br t, 2H), 3.12-3.68 (m, 7H), 3.80 (s, 3H), 3.82 (s, 3H), 5.94 (s, 2H), 6.75(d, 1H, J=8Hz), 6.83 (m, 5H), 6.94 (m, 1H), 7.22 (m, 4H). MS (FAB) m/e 561 (M+H)⁺. Anal calcd for C₃₂H₃₆N₂O₇ · 0.75 H₂O: C, 66.94; H, 6.58; N, 4.88. Found: C, 67.00; H, 6.38; N, 4.59.

Example 106

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-benzoylamino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and benzoyl chloride for isobutyryl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) mixture of rotamers δ 0.65 and 0.9 (m, total of 3H), 1.4 and 1.55 (m, total of 2H), 2.05 and 2.15 (m, total of 1H), 2.6 - 3.6 (m, 8H), 5.92 (s, 2H), 6.70(d, 1H, J=8Hz), 6.82 (m, 4H), 7.2 - 7.4 (m, 6H). MS (DCI/NH₃) m/e 531 (M+H)⁺. Anal calcd for C₃₁H₃₄N₂O₆ · 0.3 H₂O: C, 69.46; H, 6.51; N, 5.23. Found: C, 69.48; H, 6.19; N, 4.84.

Example 107

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-benzyloxycarbonylamino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and benzyl chloroformate for isobutyryl chloride in Example 61C. The crude product was purified by preparative HPLC (Vydac μC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.8 (m,

3H) 1.45 (m, 2H), 2.20 (br m, 1H), 2.75 (m, 1H), 2.93 (m, 1H), 3.15 (m, 2H), 3.32 (m, 3H), 3.52 (m, 2H), 3.66 (m, 1H), 3.78 (s, 3H), 5.00 (m, 2H), 5.94 (s, 2H), 6.72(d, 1H, J=8Hz), 6.82 (m, 3H), 7.0 (br d, 1H, J= 15Hz), 7.2 (s, 4H), 7.30 (m, 3H). MS (FAB) m/e 561 (M+H)⁺.

Anal calcd for C₃₂H₃₆N₂O₇ · 1.0 TFA: C, 60.53; H, 5.53; N, 4.15. Found: C, 60.66; H, 5.34;

5 N, 4.28.

Example 108

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(4-methoxybenzoyloxycarbonyl)amino)ethyl]pyrrolidine-3-carboxylic acid

10 The title compound is prepared by the methods described in Example 61, substituting propylamine for methylamine in Example 61B and 4-methoxybenzyl chloroformate for isobutryl chloride in Example 61C.

Example 109

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-butyl-N-ethoxycarbonylamino)ethyl]pyrrolidine-3-carboxylic acid

15 The title compound was prepared by the methods described in Example 61, but substituting butylamine for methylamine in Example 61B and ethyl chloroformate for isobutryl chloride in Example 61C. The crude product was purified by preparative HPLC (Vydac μC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, 3H, J=8Hz), 1.20 (m, 5H), 1.34 (m, 2H), 3.08 (m, 2H), 3.17 (m, 2H), 3.52 (m, 2H), 3.75 (m, 2H), 3.78 (s, 3H), 4.06 (q, 2H, J=8Hz), 4.35 (br s, 1H), 5.94 (s, 2H), 6.76 (d, 1H, J=8Hz), 6.92 (d, 2H, J=8Hz), 7.03 (br s, 1H), 7.17 (br s, 1H), 7.7 (br s, 2H). MS (FAB) m/e 513 (M+H)⁺.

20 25 Anal calcd for C₂₈H₃₆N₂O₇ · 0.5 TFA: C, 61.15; H, 6.46; N, 4.92. Found: C, 60.99; H, 6.80; N, 4.93.

Example 110

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-butyl-N-propoxycarbonylamino)ethyl]pyrrolidine-3-carboxylic acid

30 The title compound was prepared by the methods described in Example 61, but substituting butylamine for methylamine in Example 61B and propyl chloroformate for isobutryl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (br s, 1H), 0.85 (t, 3H, J=8Hz), 0.92 (br s, 1H), 1.22 (m, 3H), 1.40 (m, 3H), 1.62 (br m, 1H), 2.15 (br s, 1H), 2.72 (m, 1H), 2.87 (m, 1H), 3.1-3.45 (m, 5H), 3.55 (m, 1H), 3.64 (d, 1H, J=8Hz), 3.79 (s, 3H), 3.88

35

(br s, 1H), 3.97 (br s, 1H), 5.95 (s, 2H), 6.73(d, 1H, J=8Hz), 6.85 (m, 3H, 7.0 (s, 1H), 7.30 (d, 2H, J=8Hz). MS (FAB) m/e 527 (M+H)⁺. Anal calcd for C₂₉H₃₈N₂O₇ · 0.15 H₂O: C, 65.80; H, 7.29; N, 5.29. Found: C, 65.79; H, 7.30; N, 5.21.

5

Example 111

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-propoxycarbonylamino)ethyl]pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods described in Example 61, but substituting propylamine for methylamine in Example 61B and propyl chloroformate for isobutyryl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether-hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (t, 3H, J=8Hz), 0.93 (m, 3H), 1.43 (m, 3H), 1.62 (m, 1H), 2.15 (br s, 1H), 2.68-3.45 (m, 8H), 3.54 (m, 1H), 3.66 (m, 1H), 3.78 (s, 3H), 3.94 (m, 2H), 5.94 (s, 2H), 6.72 (d, 1H, J=8Hz), 6.82 (m, 1H), 6.84 (d, 2H, J=8Hz), 7.00 (br s, 1H), 7.33 (m, 2H). MS (DCI/NH₃) m/e 513 (M+H)⁺. Anal calcd for C₂₈H₃₆N₂O₇ · 0.15 H₂O: C, 65.26; H, 7.10; N, 5.44. Found: C, 65.22; H, 6.74; N, 5.06.

15

Example 112

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2,4-di(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

20

Ethyl (3,4-methylenedioxybenzoyl)acetate, prepared by the method of Krapcho *et al.*, Org. Syn. 47, 20 (1967) starting with 3,4-methylenedioxyacetophenone instead of 4-methoxyacetophenone, was reacted by the procedures described in Example 1 to give the title compound as a white solid. m.p. 58-60 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.87 (quintet, J=6Hz, 6H), 1.12 (sextet, J=6Hz, 2H), 1.24-1.51 (m, 6H), 2.80 (d, J=13Hz, 1H), 2.94-3.12 (m, 4H), 3.28-3.50 (m, 4H), 3.58-3.62 (m, 1H), 3.78 (d, J=9Hz, 1H), 5.95 (s, 4H), 6.73 (dd, J=8Hz, 3Hz, 2H), 6.84-6.89 (m, 2H), 6.92 (d, J=1Hz, 1H), 7.01 (d, H=1Hz, 1H). MS (DCI/NH₃) m/e 525 (M+H)⁺.

25

30

Example 113

trans,trans-1-(2-(N-(n-Butyl)-N-propylsulfonylamino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 64-65 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.83 (t, J=7Hz, 3H), 0.98 (t, J=7Hz, 3H), 1.12-1.25 (m, 2H), 1.32-1.41 (m, 2H), 1.75 (sextet, J=7Hz, 2H), 2.23-2.31 (m, 2H), 2.72-3.32 (m, 8H), 3.43 (dd, J=9Hz, 3Hz, 1H), 3.53-3.59 (m, 1H), 3.65 (d, J=9Hz, 1H),

35

3.80 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.83 (dd, J=8Hz, 1Hz, 1H), 6.88 (d, J=9Hz, 2H), 7.02 (d, J=1Hz, 1H), 7.33 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 547 (M+H)⁺.

Example 114

5 *trans,trans*-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Examples 28 and 43, the title compound was prepared as a white solid. m.p. 74-76 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (t, J=6Hz, 3H), 0.88 (t, J=8Hz, 3H), 1.08 (sextet, J=8Hz, 2H), 1.21-1.48 (m, 6H), 2.75 (d, J=12Hz, 1H), 2.95-3.09 (m, 4H), 3.26-3.59 (m, 5H), 3.75 (d, J=9Hz, 1H), 3.79 (s, 3H), 4.28 (s, 4H), 6.78 (d, J=9Hz, 1H), 6.85 (d, J=9Hz, 2H), 6.91 (d,d, J=3Hz, 9Hz, 1H), 6.98 (d, J=3Hz, 1H), 7.32 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 525 (M+H)⁺.

Example 115

15 *trans,trans*-1-(2-(N-Propyl-N-propylsulfonylamino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 72-73 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.79 (t, J=8Hz, 3H), 0.98 (t, J=8Hz, 3H), 1.43 (sextet, J=8Hz, 2H), 1.75 (sextet, J=8Hz, 2H), 2.22-2.32 (m, 1H), 2.69-3.32 (m, 9H), 3.42 (dd, J=3Hz, 12Hz, 1H), 3.52-3.58 (m, 1H), 3.64 (d, J=12Hz, 1H), 3.80 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=11Hz, 1H), 6.83 (dd, J=1Hz, 11Hz, 1H), 6.87 (d, J=11Hz, 2H), 7.0 (d, J=2Hz, 1H), 7.32 (d, J=11Hz, 2H). MS (DCI/NH₃) m/e 533 (M+H)⁺.

Example 116

25 *trans,trans*-1-(2-(N-Butyl-N-butylsulfonylamino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 62-63 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=6Hz, 3H), 0.91 (t, J=6Hz, 3H), 1.20 (sextet, J=6Hz, 2H), 1.33-1.42 (m, 4H), 1.68 (quintet, J=6Hz, 3H), 2.23-2.32 (m, 1H), 2.70-3.28 (m, 9H), 3.41 (d, J=8Hz, 1H), 3.52-3.58 (m, 1H), 3.65 (d, J=8Hz, 1H), 3.79 (s, 3H), 5.95 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.82 (d, J=8Hz, 1H), 6.87 (d, J=8Hz, 2H), 7.01 (s, 1H), 7.32 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 561 (M+H)⁺.

Example 117

35 *trans,trans*-1-(2-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

4-Hydroxyacetophenone was treated with chloromethyl methyl ether and triethylamine in THF at room temperature to give ethyl 4-methoxymethoxybenzoylacetate which was treated by the procedures described in Example 1 to afford the title compound as a white solid. m.p. 48-49 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.81 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.06 (sextet, J=7Hz, 2H), 1.20-1.35 (m, 4H), 1.44 (quintet, J=7Hz, 2H), 2.75 (d, J=12Hz, 1H), 2.94-3.10 (m, 4H), 3.25-3.35 (m, 1H), 3.40 (d, J=12Hz, 1H), 3.43-3.52 (m, 2H), 3.47 (s, 3H), 3.55-3.62 (m, 1H), 3.77 (d, J=9Hz, 1H), 5.15 (s, 2H), 5.94 (m, 2H), 6.73 (d, J=8Hz, 1H), 6.86 (dd, J=1Hz, 8Hz, 1H), 7.0 (d, J=8Hz, 2H), 7.04 (d, J=1Hz, 1H), 7.32 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 541 (M+H)⁺.

Example 118

trans,trans-1-(2-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-hydroxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid hydrochloride salt

The compound resulting from Example 116 was treated with concentrated HCl in 1:1 THF-isopropanol to give the title compound as a white solid. m.p. 211-212 °C. ¹H NMR (CD₃OD, 300 MHz) δ 0.90 (t, J=8Hz, 6H), 1.12-1.27 (m, 6H), 1.36-1.45 (m, 2H), 3.04 (bs, 1H), 3.14-3.35 (t, J=9Hz, 1H), 3.90 (bs, 3H), 4.17 (d, J=15Hz, 1H), 5.96 (s, 2H), 6.82-6.93 (m, 4H), 7.03 (d, J=1Hz, 1H), 7.42 (bs, 2H). MS (DCI/NH₃) m/e 497 (M+H)⁺.

Example 119

trans,trans-1-(2-(N-Isobutyl-N-propylsulfonylamino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 73-74 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (d, J=6Hz, 6H), 0.98 (t, J=8Hz, 3H), 1.62 (sextet, J=6Hz, 1H), 1.74 (sextet, J=8Hz, 2H), 2.23-2.34 (m, 1H), 2.68-2.98 (m, 7H), 3.08-3.18 (m, 1H), 3.26-3.42 (m, 2H), 3.52-3.58 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.80 (s, 3H), 5.90 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.82 (d, J=8Hz, 1H), 6.86 (d, J=8Hz, 2H), 6.98 (d, J=1Hz, 1H), 7.33 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 547 (M+H)⁺.

Example 120

trans,trans-1-(2-(N-Benzenesulfonyl-N-propylamino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 89-91 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.74 (t, J=6Hz, 3H), 1.33 (sextet, J=6Hz, 2H), 2.20-2.30 (m, 1H), 2.62-2.72 (m, 1H), 2.85-3.05 (m, 4H), 3.12-3.22 (m, 1H), 3.38 (dd, J=3Hz, 9Hz, 1H), 3.49-3.57 (m, 1H), 3.62 (d, J=9Hz, 1H), 3.82 (s, 3H), 5.96 (s, 2H),

6.73 (d, J=8Hz, 1H), 6.84 (dd, J=1Hz, 8Hz, 1H), 6.85 (d, J=9Hz, 2H), 7.02 (d, J=1Hz, 1H), 7.28 (d, J=9Hz, 2H), 7.39-7.54 (m, 3H), 7.70 (d, J=7Hz, 2H). MS (DCI/NH₃) m/e 567 (M+H)⁺.

5

Example 121

trans,trans-1-(2-(N-(4-Methoxybenzenesulfonyl)-N-propylamino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 96-97 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.73 (t, J=7Hz, 3H), 1.34 (sextet, J=7Hz, 2H), 2.20-2.30 (m, 1H), 2.62-2.71 (m, 1H), 2.82-3.03 (m, 4H), 3.08-3.18 (m, 2H), 3.38 (dd, J=3Hz, 9Hz, 1H), 3.48-3.56 (m, 1H), 3.62 (d, J=9Hz, 1H), 3.81 (s, 3H), 3.86 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.81-6.89 (m, 5H), 7.01 (d, J=1Hz, 1H), 7.28 (d, J=8Hz, 2H), 7.62 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 597 (M+H)⁺.

15

Example 122

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(2-methoxyethoxy-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

2-Hydroxy-5-methoxyacetophenone was treated with sodium hydride and bromoethyl methyl ether in THF at 70 °C to provide ethyl 2-methoxyethoxy-4-methoxybenzoyleacetate which was treated by the procedures described in Example 1 to provide the title compound as a white solid. m.p. 63-65 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.84 (t, J=7Hz, 3H), 0.89 (t, J=7Hz, 3H), 1.16 (sextet, J=7Hz, 2H), 1.28 (sextet, J=7Hz, 2H), 1.45-1.52 (m, 4H), 2.87-2.94 (m, 2H), 3.00-3.16 (m, 3H), 3.26-3.36 (m, 2H), 3.43 (s, 3H), 3.47-3.54 (m, 3H), 3.66-3.72 (m, 2H), 3.78 (s, 3H), 3.76-3.84 (m, 1H), 4.02-4.10 (m, 2H), 4.25 (d, J=9Hz, 1H), 5.92 (s, 2H), 6.40 (d, J=2Hz, 1H), 6.52 (dd, J=2Hz, 9Hz, 1H), 6.70 (d, J=8Hz, 1H), 6.83 (dd, J=1Hz, 8Hz, 1H), 5.98 (d, J=2Hz, 1H), 7.53 (d, J=9Hz, 1H). MS (DCI/NH₃) m/e 585 (M+H)⁺.

25

Example 123

trans,trans-1-(2-(N-Propyl-N-(2,4-dimethylbenzenesulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 88-90 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.69 (t, J=7Hz, 3H), 1.32 (sextet, J=7Hz, 2H), 2.12-2.20 (m, 1H), 2.32 (s, 3H), 2.47 (s, 3H), 2.62-2.69 (m, 1H), 2.78 (t, J=9Hz, 1H), 2.89 (dd, J=8Hz, 1H), 3.02 (sextet, J=9Hz, 2H), 3.15-3.32 (m, 3H), 3.46-3.55 (m, 1H), 3.60 (d, J=9Hz, 1H), 3.82 (s, 3H), 5.96 (s, 2H), 6.72 (d, J=7Hz, 1H), 6.80 (dd, J=1Hz, 9Hz,

35

1H), 6.86 (d, J=9Hz, 2H), 6.97 (d, J=1Hz, 1H), 7.03 (bs, 2H), 7.29 (d, J=9Hz, 1H). MS (DCI/NH₃) m/e 595 (M+H)⁺.

Example 124

trans,trans-1-(2-(N-Propyl-N-(3-chloropropylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 75-76 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (t, J=7Hz, 3H), 1.45 (sextet, J=7Hz, 2H), 2.15-2.31 (m, 3H), 2.70-2.80 (m, 1H), 2.85-3.10 (m, 6H), 3.23-3.31 (m, 2H), 3.43 (bd, J=9Hz, 1H), 3.55-3.66 (m, 4H), 3.81 (s, 3H), 5.94 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.82 (d, J=8Hz, 1H), 6.86 (d, J=8Hz, 2H), 7.00 (s, 1H), 7.33 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 567 (M+H)⁺.

Example 125

trans,trans-1-(2-(N-Propyl-N-(2-methoxyethylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, *trans,trans*-1-(2-(N-Propyl-N-(vinylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid was prepared. Ester hydrolysis using aqueous sodium hydroxide in methanol afforded the title compound as a white solid. m.p. 62-64 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.78 (t, J=7Hz, 3H), 1.42 (sextet, J=7Hz, 2H), 2.23-2.32 (m, 1H), 2.72-2.79 (m, 1H), 2.86-3.05 (m, 4H), 3.10-3.27 (m, 4H), 3.32 (s, 3H), 3.43 (dd, J=3Hz, 9Hz, 1H), 3.53-3.58 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.69 (t, J=6Hz, 2H), 3.80 (s, 3H), 5.94 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.82 (dd, J=1Hz, 8Hz, 1H), 6.87 (d, J=8Hz, 2H), 7.02 (d, J=1Hz, 1H), 7.33 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 549 (M+H)⁺.

Example 126

trans,trans-1-(2-(N-Propyl-N-(2-ethoxyethylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 58-60 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.78 (t, J=7Hz, 3H), 1.18 (t, J=7Hz, 3H), 1.43 (sextet, J=7Hz, 2H), 2.24-2.33 (m, 1H), 2.70-2.80 (m, 1H), 2.87-3.05 (m, 4H), 3.13-3.20 (m, 2H), 3.22-3.32 (m, 2H), 3.42 (dd, J=2Hz, 9Hz, 1H), 3.46 (q, J=7Hz, 2H), 3.52-3.58 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.72 (t, J=6Hz, 2H), 3.80 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=7Hz, 1H), 6.83 (dd, J=1Hz, 7Hz, 1H), 6.87 (d, J=8Hz, 2H), 7.00 (d, J=1Hz, 1H), 7.32 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 563 (M+H)⁺.

Example 127*trans,trans*-1-(2-(N-Propyl-N-(5-dimethylamino-1-naphthylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

- 5 Using the procedures described in Example 66, the title compound was prepared as a yellow solid. m.p. 102-104 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.62 (t, J=7Hz, 3H), 1.28 (sextet, J=7Hz, 2H), 2.12-2.20 (m, 1H), 2.78 (t, J=9Hz, 1H), 2.88 (s, 6H), 2.72-2.89 (m, 1H), 3.05-3.12 (m, 2H), 3.26-3.45 (m, 3H), 3.45-3.52 (m, 1H), 3.58 (d, J=9Hz, 1H), 6.97 (d, J=1Hz, 1H), 7.13 (d, J=7Hz, 1H), 7.26 (d, J=8Hz, 1H), 7.42-7.50 (m, 2H), 8.08 (dd, J=1Hz, 7Hz, 1H), 8.20 (d, J=8Hz, 1H), 8.48 (d, J=8Hz, 1H). MS (DCI/NH₃) m/e 660 (M+H)⁺.

Example 128*trans,trans*-1-(2-(N-Propyl-N-(ethylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

- 15 Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 70-72 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.79 (t, J=8Hz, 3H), 1.28 (t, J=7Hz, 3H), 1.43 (q, J=8Hz, 2H), 2.22-2.30 (m, 1H), 2.71-2.80 (m, 1H), 2.82-3.10 (m, 6H), 3.18-3.32 (m, 2H), 3.43 (dd, J=3Hz, 9Hz, 1H), 3.53-3.60 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.80 (s, 3H), 5.96 (s, 2H), 6.73 (d, J=7Hz, 1H), 6.82 (dd, J=1Hz, 7Hz, 1H), 6.88 (d, J=8Hz, 2H), 7.00 (d, J=1Hz, 1H), 7.32 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 519 (M+H)⁺.

Example 129*trans,trans*-1-(2-(N-Propyl-N-(4-methylbenzenesulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

- 25 Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 78-79 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.73 (t, J=7Hz, 3H), 1.33 (sextet, J=7Hz, 2H), 2.20-2.30 (m, 1H), 2.40 (s, 3H), 2.61-2.72 (m, 1H), 2.83-3.05 (m, 4H), 3.08-3.19 (m, 2H), 3.48 (dd, J=3Hz, 9Hz, 1H), 3.49-3.57 (m, 1H), 3.62 (d, J=9Hz, 1H), 3.81 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.82 (d, J=8Hz, 1H), 6.87 (d, J=8Hz, 2H), 7.00 (s, 1H), 7.21 (d, J=8Hz, 2H), 7.29 (d, J=8Hz, 2H), 7.57 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 581 (M+H)⁺.

Example 130*trans,trans*-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(3-pyridyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

- 35 Methyl nicotinoyl acetate was prepared by the method of Wenkert, et al., J. Org. Chem. 48: 5006 (1983) and treated by the procedures described in Example 1 to provide the

title compound as a white solid. m.p. 167-168 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=7Hz, 3H), 0.89 (t, J=7Hz, 3H), 1.14 (sextet, J=7Hz, 2H), 1.23-1.48 (m, 6H), 2.86-3.20 (m, 6H), 3.34-3.43 (m, 2H), 3.57 (dd, J=3Hz, 9Hz, 1H), 3.75-3.83 (m, 1H), 4.08 (d, J=9Hz, 1H), 5.93 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.90 (dd, J=2Hz, 8Hz, 1H), 7.03 (d, J=2Hz, 1H), 7.38 (dd, J=4Hz, 8Hz, 1H), 8.04 (d, J=8Hz, 1H), 8.48 (dd, J=2Hz, 4Hz, 2H). MS (DCI/NH₃) m/e 482 (M+H)⁺.

Example 131

trans,trans-1-(2-(N-Propyl-N-(n-butylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 65-66 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.78 (t, J=7Hz, 3H), 0.92 (t, J=7Hz, 3H), 1.31-1.46 (m, 4H), 1.68 (quintet, J=7Hz, 2H), 2.21-2.32 (m, 1H), 2.70-3.08 (m, 7H), 3.12-3.23 (m, 2H), 3.42 (dd, J=2Hz, 9Hz, 1H), 3.52-3.58 (m, 1H), 3.64 (d, J=9Hz, 1H), 3.80 (s, 3H), 5.96 (s, 2H), 6.72 (d, J=7Hz, 1H), 6.83 (dd, J=1Hz, 7Hz, 1H), 6.86 (d, J=8Hz, 2H), 7.00 (d, J=1Hz, 1H), 7.32 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 547 (M+H)⁺.

Example 132

trans,trans-1-(2-(N-Propyl-N-(4-chlorobenzenesulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 105-106 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.72 (t, J=7Hz, 3H), 1.34 (sextet, J=7Hz, 2H), 2.56-2.62 (m, 1H), 2.78-2.86 (m, 1H), 2.96-3.03 (m, 3H), 3.13-3.26 (m, 3H), 3.51 (dd, J=5Hz, 9Hz, 1H), 3.62-3.68 (m, 1H), 3.80 (s, 3H), 3.94 (d, J=9Hz, 1H), 5.92 (s, 2H), 6.75 (d, J=8Hz, 1H), 6.84 (dd, J=2Hz, 8Hz, 1H), 6.94 (d, J=8Hz, 2H), 6.98 (d, J=2Hz, 1H), 7.36 (d, J=8Hz, 1H), 7.49 (d, J=8Hz, 1H), 7.68 (d, J=8Hz, 1H). MS (DCI/NH₃) m/e 601 (M+H)⁺.

Example 133

trans,trans-1-(2-(N-Propyl-N-(benzylsulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 88-89 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.72 (t, J=7Hz, 3H), 1.32 (sextet, J=7Hz, 2H), 2.06-2.16 (m, 1H), 2.56-2.67 (m, 1H), 2.75-3.10 (m, 6H), 3.30 (dd, J=2Hz, 9Hz, 1H), 5.95 (s, 2H), 6.73 (d, J=7Hz, 1H), 6.80 (dd, J=1Hz, 7Hz, 1H), 6.86 (d, J=8Hz, 2H), 6.97 (d, J=1Hz, 1H), 7.27-7.35 (m, 7H). MS (DCI/NH₃) m/e 581 (M+H)⁺.

Example 134

trans,trans-1-(2-(N-Propyl-N-(4-fluorobenzenesulfonyl)amino)ethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 91-93 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.73 (t, J=7Hz, 3H), 1.44 (sextet, J=7Hz, 2H), 2.18-2.27 (m, 1H), 2.56-2.67 (m, 1H), 2.78-2.87 (m, 2H), 2.97 (septet, J=8Hz, 2H), 3.11-3.16 (m, 2H), 3.33 (dd, J=2Hz, 9Hz, 1H), 3.43-3.50 (m, 1H), 3.57 (d, J=9Hz, 1H), 3.78 (s, 3H), 7.08 (t, J=8Hz, 2H), 7.24 (d, J=8Hz, 2H), 7.69 (dd, J=5Hz, 8Hz, 2H). MS (DCI/NH₃) m/e 585 (M+H)⁺.

Example 135

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-benzofuranyl)pyrrolidine-3-carboxylic acid

Example 135A

Benzofuran-4-carboxaldehyde

To a suspension of 60% sodium hydride in mineral oil (4.00 g, 100 mmol, 1.25 eq) in DMF (60 mL) at 0 °C was added a solution of 3-bromophenol (13.8 g, 80 mmol) in DMF (5 mL). After 10 minutes, bromoacetaldehyde diethyl acetal (14.9 mL, 96.6 mmol, 1.24 eq) was added, and the resultant mixture then heated at 120 °C for 2.5 hours. The mixture was cooled to room temperature and was poured into water, and extracted once with ether. The organic solution was dried over MgSO₄, filtered, evaporated and vacuum distilled to yield a colorless liquid (17.1 g, 74%). b.p. 160-163 °C at 0.4 mm Hg.

To warm polyphosphoric acid (15.3 g) was added a solution of the above compound (17.1 g, 59.3 mmol) in benzene (50 mL). The resultant mixture was heated under reflux with vigorous stirring for 4 hours, after which time the benzene layer was carefully decanted off, and the lower layer washed once with hexanes. The combined organic solutions were concentrated *in vacuo*, and then vacuum distilled to yield a colorless liquid (8.13 g, 70%). b.p. 62-72 °C at 0.6 mm Hg.

To a solution of the above compounds (8.11 g, 41.5 mmol) in ether (80 mL) at -78 °C was added 1.7 M *t*-butyllithium (48.8 mL, 83 mmol, 2 eq) such that the temperature did not exceed -70 °C. After stirring for 15 minutes, a solution of DMF (6.5 mL, 83 mmol, 2 eq) in ether (20 mL) was added, and the mixture allowed to warm to room temperature over 2 hours. The mixture was poured into water and the phases separated. The organic solution was dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash chromatography on

silica gel eluting with 10% ether in hexanes to yield benzofuran-6-carboxaldehyde (1.22 g) and benzofuran-4-carboxaldehyde (1.86 g), both as colorless oils.

Example 135B

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-benzofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Examples 1 and 49 substituting the compound resulting from Example 135A in Example 49A for piperonal. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 7.59 (1H, t, J=3Hz), 7.4-7.2 (6H, m), 6.8 (2H, d, J=8Hz), 4.03 (1H, m), 3.94 (1H, dd, J=8Hz, 3Hz), 3.77 (3H, s), 3.61 (1H, dd, J=8Hz, 7.3Hz), 3.42 (1H, dd, J=11Hz, 5Hz), 3.40-2.90 (5H, m), 2.82 (2.81) (3H, s), 1.50 (2H, septet, J=7Hz), 0.82 (0.75) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 451 (M+H)⁺. Anal.calc. for C₂₆H₃₀N₂O₅ · AcOH: C, 65.87; H, 6.71; N, 5.49. Found: C, 66.04; H, 6.42; N, 5.60. s

Example 136

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(6-benzofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in Examples 1 and 49 substituting benzofuran-6-carboxaldehyde, prepared as described in Example 135A, in Example 49A for piperonal. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 7.65 (1H, bd), 7.60 (1H, d, J=2Hz), 7.55 (1H, d, J=8Hz), 7.35 (3H, m), 6.85 (2H, dd, J=8Hz, 3Hz), 6.75 (1H, dd, J=3Hz, 2Hz), 3.83 (2H, m), 3.79 (3H, s), 3.60-3.0 (7H, m), 2.91 (2.83) (s, 3H), 1.51 (2H, septet, J=7Hz), 0.83 (0.78) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 451 (M+H)⁺. Anal.calc. for C₂₆H₃₀N₂O₅ · 0.5 H₂O: C, 67.96; H, 6.80; N, 6.10. Found: C, 67.90; H, 6.71; N, 6.07.

Example 137

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(6-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by catalytic hydrogenation (4 atmospheres of H₂ in AcOH, followed by preparative hplc) of the compound resulting from Example 136. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 7.49 (7.47) (2H, d, J=8Hz), 7.19 (1H, d, J=8Hz), 7.00 (1H, m), 7.82 (3H, m), 5.40 (1H, dd, J=11Hz, 7Hz), 4.58 (2H, t, J=8Hz), 4.18 (1H, m), 4.10 (1H, m), 3.88 (1H, m), 3.79 (3H, s), 3.60 (1H, m), 3.35 (1H, m), 3.19 (2H, t, J=8Hz), 3.00 (4H, m), 2.91 (2.78) (s, 3H), 1.53 (1.40) (2H, septet, J=7Hz), 0.88 (0.78) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 453 (M+H)⁺. Anal.calc. for C₂₆H₃₂N₂O₅ · 1.25 TFA: C, 57.53; H, 5.63; N, 4.71. Found: C, 57.68; H, 5.68; N, 4.70.

Example 138*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-benzofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting benzofuran-4-carboxaldehyde in Example 49A for piperonal and substituting N,N-dibutyl bromoacetamide for N-methyl-N-propyl bromoacetamide. ¹H NMR (300 MHz, CDCl₃) δ 7.62 (1H, d, J=3Hz), 7.39 (1H, dt, J=8Hz, 2Hz), 7.34 (3H, m), 7.26 (1H, d, J=2Hz), 7.23 (1H, d, J=8Hz), 6.84 (2H, d, J=8Hz), 4.02 (1H, ddd, J=8, 6Hz, 4Hz), 3.89 (1H, d, J=9Hz), 3.79 (3H, s), 3.67 (1H, dd, J=10Hz, 3Hz), 3.44 (2H, m), 3.35-3.15 (3H, m), 3.00 (2H, m), 2.84 (1H, d, J=14Hz), 1.43 (3H, m), 1.23 (3H, m), 1.08 (2H, m), 0.87 (3H, t, J=7Hz), 0.82 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 507 (M+H)⁺. Anal.calc. for C₃₀H₃₈N₂O₅: C, 71.12; H, 7.56; N, 5.53. Found: C, 70.86; H, 7.45; N, 5.24.

Example 139*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-benzofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting benzofuran-5-carboxaldehyde, prepared by the procedures described in Example 135A substituted 4-bromophenol for 3-bromophenol, in Example 49A for piperonal and substituting N,N-dibutyl bromoacetamide for N-methyl-N-propyl bromoacetamide. ¹H NMR (300 MHz, CDCl₃) δ 7.64 (1H, bd), 7.59 (1H, d, J=2Hz), 7.43 (2H, m), 7.33 (2H, d, J=8Hz), 6.85 (2H, d, J=8Hz), 6.73 (1H, dd, J=3Hz, 1Hz), 3.82 (1H, d, J=11Hz), 3.89 (1H, d, J=9Hz), 3.79 (3H, s), 3.53 (1H, dd, J=10Hz, 3Hz), 3.44 (2H, m), 3.30 (1H, m), 3.20-2.95 (5H, m), 2.82 (1H, d, J=14Hz), 1.43 (3H, m), 1.23 (3H, m), 1.08 (2H, m), 0.87 (3H, t, J=7Hz), 0.82 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 507 (M+H)⁺. Anal.calc. for C₃₀H₃₈N₂O₅: C, 71.12; H, 7.56; N, 5.53. Found: C, 70.73; H, 7.45; N, 5.29.

Example 140*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(6-benzofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting benzofuran-6-carboxaldehyde in Example 49A for piperonal and substituting N,N-dibutyl bromoacetamide for N-methyl-N-propyl bromoacetamide. ¹H NMR (300 MHz, CDCl₃) δ 7.63 (1H, bd), 7.59 (1H, d, J=2Hz), 7.53 (1H, d, J=8Hz), 7.36 (3H, m), 6.85 (2H, d, J=8Hz), 6.73 (1H, dd, J=3Hz, 1Hz), 3.82 (1H, d, J=11Hz), 3.89 (1H, d, J=9Hz), 3.79 (3H,

s), 3.53 (1H, dd, J=10Hz, 3Hz), 3.44 (2H, m), 3.30 (1H, m), 3.20-2.95 (5H, m), 2.80 (1H, d, J=14Hz), 1.43 (3H, m), 1.23 (3H, m), 1.08 (2H, m), 0.87 (3H, t, J=7Hz), 0.82 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 507 (M+H)⁺. Anal.calc. for C₃₀H₃₈N₂O₅ · 0.75 H₂O: C, 69.28; H, 7.65; N, 5.39. Found: C, 69.11; H, 7.33; N, 5.32.

Example 141

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(6-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by catalytic hydrogenation of the compound resulting from Example 140 (4 atmospheres of H₂ in AcOH, followed by preparative hplc). ¹H NMR (300 MHz, CDCl₃) δ 7.40 (2H, d, J=8Hz), 7.16 (1H, d, J=8Hz), 6.97 (1H, dd, J=8Hz, 2Hz), 6.89 (3H, m), 5.90 (1H, bs) 4.57 (2H, t, J=9Hz), 4.93 (2H, m), 3.80 (3H, s), 3.70-3.58 (2H, m), 3.40 (1H, m), 3.30-2.90 (8H, m), 1.40 (2H, m), 1.29 (3H, m), 1.08 (2H, m), 0.92 (3H, t, J=7Hz), 0.82 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 509 (M+H)⁺. Anal.calc. for C₃₀H₄₀N₂O₅ · 0.85 TFA: C, 62.88; H, 6.80; N, 4.63. Found: C, 63.04; H, 6.66; N, 4.60.

Example 142

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(5-indanyl)pyrrolidine-3-carboxylic acid

Example 142A

Indane-5-carboxaldehyde

Indane-5-carboxaldehyde was prepared by formylation of indane under the conditions described for 2,3-dihydrobenzofuran in Example 52A. The resultant mixture of 4- and 5-carboxaldehydes was purified as follows: to a 6:1 mixture of indane-4-carboxaldehyde and indane-5-carboxaldehyde (3.46 g, 23 mmol) was added aniline (2.20 g, 23 mmol, 1 eq). The resultant solution slowly solidified to a mixture of imines which was recrystallized from hot acetonitrile to yield the 5-aldimine as a white solid. The aldimine (2.65 g) was suspended in water (6 mL), and treated with 4 N hydrochloric dioxane (10 mL). The mixture was boiled for 1 hour, cooled to room temperature, and poured into ether. The organic solution was dried over MgSO₄, filtered, and concentrated *in vacuo*. Vacuum distillation of the residue afforded indane-5-carboxaldehyde (1.54 g, 88%) as a colorless liquid. b.p. 88-90 °C at 0.9 mm Hg.

Example 142B

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(5-indanyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting indane-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 7.25-7.1 (5H, m), 6.78 (2H, d, J=8Hz), 3.89 (1H, d, J=8Hz), 3.75 (3H, s), 3.50-2.90 (6H, m), 2.88 (6H, t, J=6Hz), 2.82 (2.80) (3H, s), 2.04 (2H, t, J=8Hz), 1.48 (2H, septet, J=7Hz), 0.83 (0.73) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 451 (M+H)⁺, 473 (M+Na)⁺. Anal.calc. for C₂₇H₃₄N₂O₄ · 2.5 H₂O : C, 65.44; H, 7.93; N, 5.65. Found: C, 65.36; H, 7.45; N, 5.53.

Example 143

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(6-indolyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting indole-6-carboxaldehyde, prepared by the method of Rapoport, J. Org. Chem. 51: 5106 (1986), for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 8.43 (1H, brs), 7.57 (1H, d, J=8Hz), 7.43 (1H, s), 7.31 (2H, dd, J=6Hz, 3Hz), 7.22 (1H, d, J=8Hz), 7.1 (1H, t, J=3Hz), 6.78 (2H, dd, J=6Hz, 3Hz), 6.45 (1H, m), 3.93 (1H, dd, J=6Hz, 3Hz), 3.80 (1H, m), 3.73 (3H, s), 3.60-2.90 (6H, m), 2.86 (2.82) (3H, s), 1.47 (2H, septet, J=7Hz), 0.83 (0.73) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 450 (M+H)⁺. Anal.calc. for C₂₆H₃₁N₃O₄ · 0.75 H₂O : C, 67.44; H, 7.07; N, 9.07. Found: C, 67.42; H, 7.09; N, 8.91.

Example 144

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(3,4-difluorophenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 3,4-difluorobenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 7.60-7.3 (4H, m), 7.13 (1H, q, J=9Hz), 6.90 (2H, d, J=8Hz), 3.90 (1H, m), 3.79 (3H, s), 3.60-2.95 (6H, m), 2.92 (2.78) (3H, s), 1.55 (2H, septet, J=7Hz), 0.88 (0.73) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 447 (M+H)⁺. Anal.calc. for C₂₄H₂₈F₂N₂O₄ · 1.80 H₂O : C, 60.19; H, 6.65; N, 5.85. Found: C, 60.13; H, 6.34; N, 5.84.

Example 145

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(phenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting benzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) (minor rotamer) δ 7.53 (4H, d, J=6Hz), 7.40-7.20 (3H, m), 6.88 (2H, d, J=8Hz), 3.90 (1H, m),

3.79 (3H, s), 3.70-2.95 (8H, m), 2.90 (2.79) (3H, s), 1.50 (2H, sept, J=7Hz), 0.87 (0.72) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 411 (M+H)⁺. Anal.calc. for C₂₄H₃₀N₂O₄ · 2.00 H₂O: C, 64.55; H, 7.67; N, 6.27. Found: C, 64.37; H, 7.43; N, 6.29.

5

Example 146

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-hydroxyphenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 4-hydroxybenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃-CD₃OD) (minor rotamer) δ 7.35 (2H, d, J=8Hz), 7.28 (2H, dd, J=7Hz, 3Hz), 6.90 (2H, dd, J=7Hz, 3Hz), 6.89 (2H, d, J=8Hz), 3.81 (3H, s), 3.65 (1H, d, J=8Hz), 3.70-3.00 (8H, m), 2.92 (2.83) (3H, s), 1.50 (2H, septet, J=7Hz), 0.87 (0.77) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 427 (M+H)⁺. Anal.calc. for C₂₄H₃₀N₂O₅ · 1.00 H₂O: C, 64.85; H, 7.26; N, 6.30. Found: C, 64.82; H, 7.39; N, 6.46.

15

Example 147

trans,trans-1-(N-Methyl-N-propylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(2,4-dimethoxyphenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 2,4-dimethoxybenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃-CD₃OD) (minor rotamer) δ 7.61 (1H, d, J=8Hz), 7.30 (2H, d, J=8Hz), 6.82 (2H, d, J=8Hz), 6.55 (1H, d, J=8Hz), 6.45 (1H, d, J=3Hz), 3.90 (1H, m), 3.81 (3H, s), 3.79 (3H, s), 3.77 (3H, s), 3.70-2.90 (8H, m), 2.85 (3H, s), 1.50 (2H, sept, J=7Hz), 0.87 (0.77) (3H, t, J=7Hz). MS (DCI/NH₃) m/e 471 (M+H)⁺. Anal.calc. for C₂₆H₃₄N₂O₆ · 0.75 H₂O: C, 64.51; H, 7.39; N, 5.79. Found: C, 64.65; H, 7.07; N, 5.75.

25

Example 148

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 2,3-dihydrobenzofuran-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.31 (2H, d, J=8Hz), 7.27 (1H, d, J=2Hz), 7.18 (1H, dd, J=7Hz, 3Hz), 6.86 (2H, d, J=8Hz), 6.72 (1H, d, J=8Hz), 4.56 (2H, t, J=7Hz), 3.78 (3H, s), 3.62 (1H, m), 3.50-3.25 (4H, m), 3.17 (2H, t, J=7Hz), 3.15-2.90 (5H, m), 2.79 (1H, d, J=14Hz), 1.43 (3H, m), 1.26 (3H, m), 1.08 (2H, m), 0.87 (3H, t, J=7Hz), 0.81 (3H, t, J=7Hz). MS

35

(DCI/NH₃) m/e 509 (M+H)⁺. Anal.calc. for C₃₀H₄₀N₂O₅ · 0.25 H₂O: C, 70.22; H, 7.95; N, 5.46. Found: C, 70.21; H, 7.92; N, 5.36.

Example 149

5 trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-methoxyphenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 4-methoxybenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.38 (2H, d, J=8Hz), 7.30 (2H, d, J=8Hz), 6.87 (4H, dd, J=7Hz, 3Hz), 3.78 (3H, s), 3.76 (3H, s), 3.63 (1H, m), 3.50-3.20 (4H, m), 3.15-2.90 (5H, m), 2.78 (1H, d, J=14Hz), 1.43 (3H, m), 1.27 (3H, m), 1.09 (2H, m), 0.87 (3H, t, J=7Hz), 0.81 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 497 (M+H)⁺. Anal.calc. for C₂₉H₄₀N₂O₅: C, 70.13; H, 8.12; N, 5.64. Found: C, 69.78; H, 8.10; N, 5.54.

15 Example 150

20 trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(3,4-difluorophenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 3,4-difluorobenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.35 (1H, m), 7.30 (2H, d, J=8Hz), 7.20-7.00 (2H, m), 6.87 (2H, d, J=8Hz), 3.78 (3H, s), 3.79 (1H, m), 3.62 (1H, m), 3.50-3.30 (3H, m), 3.23 (1H, m), 3.15-2.90 (4H, m), 2.78 (1H, d, J=14Hz), 1.43 (2H, m), 1.27 (4H, m), 1.08 (2H, m), 0.85 (3H, t, J=7Hz), 0.80 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 503 (M+H)⁺. Anal.calc. for C₂₈H₃₆F₂N₂O₄ · 1 H₂O: C, 64.60; H, 7.36; N, 5.38. Found: C, 64.59; H, 7.20; N, 5.35.

25 Example 151

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(2,4-dimethoxyphenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 2,4-dimethoxybenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.37 (2H, d, J=8Hz), 7.20 (1H, d, J=8Hz), 6.92 (2H, d, J=8Hz), 6.60 (1H, d, J=3Hz), 6.49 (1H, dd, J=6Hz, 2Hz), 5.35 (1H, d, J=8Hz), 4.20 (3H, m), 4.10 (3H, s), 3.83 (3H, s), 3.81 (3H, s), 3.75 (3H, m), 3.17 (2H, hep, J=7Hz), 3.05 (2H, t, J=7Hz), 1.30 (4H, m), 1.07 (4H, m), 0.87 (3H, t, J=7Hz), 0.80 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 527 (M+H)⁺. Anal.calc. for C₃₀H₄₂N₂O₆ · 1.30 TFA: C, 58.02; H, 6.47; N, 4.15. Found: C, 57.92; H, 6.43; N, 4.07.

Example 152*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-phenyl-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl benzoylacetate in Example 49B. ¹H NMR (300 MHz, CDCl₃) δ 7.50-7.25 (5H, m), 7.04 (1H, d, J=3Hz), 6.87 (1H, dd, J=7Hz, 3Hz), 6.74 (1H, d, J=8Hz), 5.94 (1H, d, J=4Hz), 5.92 (1H, d, J=4Hz), 3.85 (1H, d, J=8Hz), 3.64 (1H, m), 3.42 (3H, m), 3.27 (2H, m), 3.20-2.90 (5H, m), 2.81 (1H, d, J=14Hz), 1.43 (2H, m), 1.27 (4H, m), 1.05 (2H, m), 0.85 (3H, t, J=7Hz), 0.80 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 481 (M+H)⁺. Anal.calc. for C₂₈H₃₆N₂O₅: C, 69.98; H, 7.55; N, 5.83. Found: C, 69.69; H, 7.63; N, 5.71.

Example 153*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-phenyl-4-(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl benzoylacetate in Example 49B and 2,3-dihydrobenzofuran-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.53 (2H, m), 7.40 (4H, m), 7.13 (1H, dd, J=7Hz, 3Hz), 6.72 (1H, d, J=8Hz), 5.40 (1H, d, J=10Hz), 4.56 (2H, t, J=8Hz), 4.18 (1H, d, J=14Hz), 4.07 (2H, m), 3.79 (2H, m), 3.48 (1H, d, J=14Hz), 3.35 (1H, m), 3.28 (3H, m), 2.95 (2H, m), 1.47 (2H, m), 1.28 (4H, m), 1.10 (2H, m), 0.93 (3H, t, J=7Hz), 0.78 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 479 (M+H)⁺. Anal.calc. for C₂₉H₃₈N₂O₄ · 1.10 TFA: C, 62.04; H, 6.52; N, 4.64. Found: C, 61.89; H, 6.44; N, 4.57.

Example 154*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-*t*-butylphenyl)-4-(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting *t*-butyl benzoylacetate, prepared by the method of Krapcho et al., Org. Syn. 47:20 (1967) starting from 4-*t*-butylacetophenone, in Example 49B and 2,3-dihydrobenzofuran-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.60-7.30 (6H, m), 6.90 (1H, m), 4.50 (2H, m), 3.95 (1H, m), 3.85-2.95 (11H, m), 2.90 (1H, d, J=14Hz), 1.58 (2H, m), 1.50 (7H, m), 1.41 (6H, s), 1.10 (2H, m), 1.00 (3H, t, J=7Hz), 0.90 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 535 (M+H)⁺. Anal.calc. for C₃₃H₄₆N₂O₄ · 0.25 H₂O: C, 73.50; H, 8.69; N, 5.19. Found: C, 73.57; H, 8.58; N, 5.14.

Example 155*trans,trans*-2-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(4-fluorophenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 4-fluorobenzaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.50 (1H, m), 7.42 (1H, dd, J=7Hz, 3Hz), 7.36 (2H, d, J=8Hz), 7.01 (3H, t, J=8Hz), 6.87 (1H, d, J=8Hz), 3.83 (1H, m), 3.8 (3H, s), 3.67 (1H, m), 3.47 (3H, m), 3.30-2.90 (5H, m), 2.82 (1H, d, J=14Hz), 1.43 (2H, m), 1.28 (4H, m), 1.08 (2H, m), 0.90 (3H, t, J=7Hz), 0.82 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 485 (M+H)⁺. Anal.calc. for C₂₈H₃₇FN₂O₄: C, 69.40; H, 7.70; N, 5.78. Found: C, 69.03; H, 8.00; N, 5.74.

Example 156*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(3-furyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting β-oxo-3-furanpropionate in Example 49B. ¹H NMR (300 MHz, CDCl₃) δ 7.41 (2H, m), 6.97 (1H, d, J=3Hz), 6.85 (1H, dd, J=7Hz, 3Hz), 6.72 (1H, d, J=8Hz), 6.42 (1H, s), 5.94 (1H, d, J=4Hz), 5.92 (1H, d, J=4Hz), 3.90 (1H, m), 3.70-3.25 (5H, m), 3.20-2.90 (4H, m), 2.85 (1H, d, J=14Hz), 1.43 (2H, m), 1.40-1.05 (6H, m), 0.90 (6H, m). MS (DCI/NH₃) m/e 471 (M+H)⁺. Anal.calc. for C₂₆H₃₄N₂O₆: C, 66.36; H, 7.28; N, 5.95. Found: C, 66.09; H, 7.24; N, 5.87.

Example 157*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(isopropyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl isobutyrylacetate in Example 49B. ¹H NMR (300 MHz, CDCl₃) δ 6.85 (1H, d, J=2Hz), 6.76 (1H, dd, J=6Hz, 2Hz), 6.71 (1H, d, J=8Hz), 5.92 (2H, s), 3.75 (1H, d, J=14Hz), 3.66 (1H, q, J=7Hz), 3.42 (3H, m), 3.25 (3H, m), 3.11 (2H, m), 2.83 (1H, t, J=7Hz), 1.88 (1H, m), 1.55 (4H, m), 1.32 (4H, m), 0.92 (12H, m). MS (DCI/NH₃) m/e 447 (M+H)⁺. Anal.calc. for C₂₅H₃₈N₂O₅ · 0.50 H₂O: C, 65.91; H, 8.63; N, 6.15. Found: C, 66.07; H, 8.10; N, 6.03.

Example 158*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-*t*-butylphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl 4-*t*-butylbenzoyleacetate, prepared by the method of Krapcho et al., Org. Syn. 47: 20 (1967) starting with 4-*t*-butylacetophenone), in Example 49B. ¹H NMR (300 MHz, CDCl₃) δ 7.32 (4H, d, J=3Hz), 7.04 (1H, d, J=2Hz), 6.87 (1H, dd, J=8Hz, 3Hz), 6.74 (1H, d, J=9Hz), 5.94 (1H, d, J=4Hz), 5.92 (1H, d, J=4Hz), 3.77 (1H, d, J=14Hz), 3.65-3.25 (5H, m), 3.15-2.85 (4H, m), 2.73 (1H, d, J=14Hz), 1.45 (2H, m), 1.29 (13H, s), 1.00 (2H, m), 0.86 (3H, t, J=7Hz), 0.76 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 537 (M+H)⁺. Anal.calc. for C₃₂H₄₄N₂O₅: C, 71.61; H, 8.26; N, 5.22. Found: C, 71.43; H, 8.09; N, 5.11.

Example 159

trans,trans-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-*t*-butylphenyl)-4-(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl isobutyrylacetate in Example 49B and 2,3-dihydrobenzofuran-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.30 (1H, s), 7.13 (1H, dd, J=7Hz, 2Hz), 6.82 (1H, d, J=8Hz), 4.68 (2H, t, J=8Hz), 4.48 (1H, s), 3.19 (3H, m), 3.80 (3H, m), 3.48 (2H, m), 3.3 (5H, m), 2.41 (1H, m), 1.65 (4H, m), 1.44 (4H, m), 1.21 (3H, d, J=5Hz), 1.17 (3H, d, J=5Hz), 1.05 (6H, m). MS (DCI/NH₃) m/e 445 (M+H)⁺. Anal.calc. for C₂₆H₄₀N₂O₄ · 1.2 TFA: C, 58.67; H, 7.14; N, 4.82. Found: C, 58.54; H, 7.25; N, 4.74.

Example 160

trans,trans-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(anti-4-methoxycyclohexyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Example 160A

syn and *anti* Ethyl 4-methoxycyclohexanoylacetate

Syn, anti-4-Methoxycyclohexane carboxylic acid (5.00 g, 31.6 mmol) and carbonyldiimidazole (6.15 g, 37.9 mmol, 1.2 eq) were stirred in anhydrous tetrahydrofuran (50 mL) for 6 hours at room temperature. At the same time, magnesium chloride (3.01 g, 31.6 mmol) and ethyl malonate potassium salt (7.52 g, 44.2 mmol, 1.4 equivalents) were stirred in anhydrous tetrahydrofuran (75 mL) for 6 hours at 50 °C. The mixture was cooled to room temperature, and the imidazole-acid mixture added to it. The reaction stirred overnight at room temperature. The solvents were removed under reduced pressure, and the residue was taken up in chloroform/water. The organic phase washed with 5% potassium bisulfate, water, and brine, dried with magnesium sulfate, filtered, and concentrated under reduced pressure.

The residue was purified by flash chromatography on 175 g silica gel, eluting with 20% ethyl acetate in hexanes. Pure fractions of the syn and anti methoxycyclohexyl β -keto esters were obtained. The solvents were removed under reduced pressure to yield the trans-4-methoxycyclohexyl β -keto ester (914 mg) as a colorless oil and the cis 4-methoxycyclohexyl β keto ester (1.07 g) as a colorless oil.

Example 160B

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(anti-4-methoxycyclohexyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting the anti-compound resulting from Example 160A in Example 49B. ^1H NMR (300 MHz, CDCl_3) δ 6.84 (1H, d, $J=2\text{Hz}$), 6.76 (1H, dd, $J=7\text{Hz}$, 2Hz), 6.61 (1H, d, $J=8\text{Hz}$), 5.92 (2H, s), 3.69 (2H, m), 3.50-3.27 (5H, m), 3.26 (3H, s), 3.25-3.00 (3H, m), 2.88 (1H, m), 1.95 (2H, m), 1.62 (7H, m), 1.33 (9H, m), 0.97 (3H, t, $J=7\text{Hz}$), 0.92 (3H, t, $J=7\text{Hz}$). MS (DCI/ NH_3) m/e 517 ($\text{M}+\text{H}$) $^+$. Anal.calc. for $\text{C}_{29}\text{H}_{44}\text{N}_2\text{O}_6 \cdot 0.50 \text{H}_2\text{O}$: C, 66.26; H, 8.63; N, 5.33. Found: C, 66.27; H, 8.50; N, 5.13.

Example 161

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2-(syn-4-methoxycyclohexyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting the syn-compound resulting from Example 160A in Example 49B. ^1H NMR (300 MHz, CDCl_3) δ 6.84 (1H, d, $J=2\text{Hz}$), 6.77 (1H, dd, $J=6\text{Hz}$, 2Hz), 6.61 (1H, d, $J=8\text{Hz}$), 5.92 (2H, s), 3.65 (2H, m), 3.42 (2H, m), 3.32 (3H, s), 3.30-3.00 (6H, m), 2.82 (1H, m), 2.10 (2H, m), 1.83 (2H, m), 1.52 (6H, m), 1.33 (4H, m), 1.20-1.00 (4H, m), 0.96 (3H, t, $J=7\text{Hz}$), 0.91 (3H, t, $J=7\text{Hz}$). MS (DCI/ NH_3) m/e 517 ($\text{M}+\text{H}$) $^+$. Anal.calc. for $\text{C}_{29}\text{H}_{44}\text{N}_2\text{O}_6 \cdot 0.30 \text{H}_2\text{O}$: C, 66.72; H, 8.61; N, 5.37. Found: C, 66.76; H, 8.65; N, 5.28.

Example 162

trans,trans-1-(N,N-Di(n-butyl)aminocarbonylmethyl)-2,4-di(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

Example 162A

5-Acetyl-2,3-dihydrobenzofuran

To a 0 °C solution of acetyl chloride (1.64 mL, 23.0 mmol, 1.3 equivalents) in methylene chloride (30 mL) was added stannic chloride (2.49 mL, 21.3 mmol, 1.2 equivalents), maintaining the temperature below 5 °C. The solution was stirred 15 minutes at 0 °C, and then a solution of 2,3-dihydrofuran (2.00 mL, 17.7 mmol) in methylene chloride (5 mL) was added dropwise while maintaining the temperature below 8 °C. The dark red solution was stirred 1 hour at 2 °C and then poured into 50 mL of ice water. The reaction was stirred an additional 30 minutes, and the layers were separated. The organic layer was washed with water and aqueous sodium bicarbonate, dried over magnesium sulfate, filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography on 150 g silica gel, eluting with 18% ethyl acetate in hexanes. The solvents were removed under reduced pressure to yield the title compound (2.68 g, 93%) as a yellow solid.

Example 162B

trans,trans-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2,4-di(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting the compound resulting from Example 162A in Example 49B and 2,3-dihydrobenzofuran-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.43 (1H, s), 7.38 (1H, s), 7.06 (2H, m), 6.75 (1H, d, J=6Hz), 6.70 (1H, d, J=6Hz), 5.40 (1H, d, J=9Hz), 4.58 (4H, q, J=7Hz), 4.16 (1H, d, J=14Hz), 4.09 (2H, m), 3.82 (2H, m), 3.57 (1H, d, J=14Hz), 3.38 (1H, m), 3.30-3.05 (6H, m), 2.95 (2H, q, J=6Hz), 1.50 (2H, m), 1.30 (4H, m), 1.15 (2H, m), 0.94 (3H, t, J=7Hz), 0.83 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 521 (M+H)⁺. Anal.calc. for C₃₁H₄₀N₂O₅ · 1.25 TFA: C, 60.67; H, 6.27; N, 4.22. Found: C, 60.49; H, 6.18; N, 4.13.

Example 163

trans,trans-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(3-furyl)-4-(5-benzo-2,3-dihydrofuranyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl β-oxo-3-furanpropionate in Example 49B and 2,3-dihydrobenzofuran-5-carboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.42 (1H, m), 7.38 (1H, m), 7.13 (1H, s), 7.16 (1H, dd, J=7Hz, 3Hz), 6.70 (1H, d, J=8Hz), 6.41 (1H, m), 4.57 (2H, t, J=7Hz), 3.95 (1H, d, J=8Hz), 3.63 (1H, m), 3.55 (1H, d, J=14), 3.50-3.25 (4H, m), 3.18 (2H, t, J=6Hz), 3.15-2.95 (3H, m), 2.87 (1H, d, J=14Hz), 1.45 (4H, m), 1.35-1.10 (4H, m), 0.85 (6H, m). MS (DCI/NH₃) m/e 469 (M+H)⁺. Anal.calc. for C₂₇H₃₆N₂O₅ · 0.25 H₂O: C, 68.55; H, 7.78; N, 5.92. Found: C, 68.62; H, 7.68; N, 5.82.

Example 164*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(3-fluorophenyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 3-fluorobenzenecarboxaldehyde for piperonal in Example 49A. ¹H NMR (300 MHz, CDCl₃) δ 7.30 (2H, d, J=8Hz), 7.22 (2H, m), 6.91 (1H, m), 6.86 (2H, d, J=8Hz), 3.79 (1H, m), 3.78 (3H, s), 3.68 (1H, m), 3.55-3.37 (3H, m), 3.29 (1H, m), 3.15-2.90 (5H, m), 2.78 (1H, d, J=14Hz), 1.43 (2H, m), 1.25 (4H, m), 1.07 (2H, m), 0.87 (3H, t, J=7Hz), 0.80 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 485 (M+H)⁺. Anal.calc. for C₂₈H₃₇FN₂O₄ · 0.25 H₂O: C, 68.76; H, 7.73; N, 5.73. Found: C, 68.87; H, 7.69; N, 5.67.

Example 165*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(3-pyridyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting 3-pyridinecarboxaldehyde for piperonal in Example 49A. The nitro styrene was prepared by the method of Bourguignon et al., Can. J. Chem. 63: 2354 (1985). ¹H NMR (300 MHz, CDCl₃) δ 8.82 (1H, bs), 8.73 (1H, bd, J=9Hz), 8.62 (1H, bd, J=7Hz), 7.78 (1H, bdd, J=9Hz, 3Hz), 7.38 (2H, d, J=10Hz), 6.90 (2H, d, J=10Hz), 4.39 (1H, d, J=12Hz), 3.95 (1H, m), 3.80 (3H, s), 3.79 (1H, m), 3.68 (1H, d, J=18Hz), 3.50-3.30 (3H, m), 3.25-2.90 (6H, m), 1.47 (2H, m), 1.31 (4H, m), 1.20 (2H, m), 0.92 (3H, t, J=7Hz), 0.83 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 468 (M+H)⁺. Anal.calc. for C₂₇H₃₇N₃O₄ · 1.65 TFA: C, 55.50; H, 5.94; N, 6.41. Found: C, 55.53; H, 5.90; N, 6.27.

Example 166*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(2-fluorophenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl 2-fluorobenzoylacetate in Example 49B. ¹H NMR (300 MHz, CDCl₃) δ 7.52 (1H, dt, J=7Hz, 3Hz), 7.25 (1H, m), 7.13 (1H, dt, J=7Hz, 3Hz), 7.02 (2H, m), 6.88 (1H, dd, J=7Hz, 3Hz), 6.73 (1H, d, J=8Hz), 5.93 (1H, d, J=4Hz), 5.92 (1H, d, J=4Hz), 4.25 (1H, d, J=9Hz), 3.68 (1H, m), 3.42 (3H, m), 3.39 (1H, m), 3.20-2.95 (4H, m), 2.91 (1H, d, J=14Hz), 1.45 (3H, m), 1.26 (3H, m), 1.08 (2H, m), 0.87 (3H, t, J=7Hz), 0.81 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 499 (M+H)⁺. Anal.calc. for C₂₈H₃₅FN₂O₅ · 0.25 H₂O: C, 66.85; H, 7.11; N, 5.57. Found: C, 66.51; H, 6.67; N, 5.18.

Example 167*trans,trans*-1-(N,N-Di(*n*-butyl)aminocarbonylmethyl)-2-(3-fluorophenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 1 and 49 substituting ethyl 3-fluorobenzoylacetate in Example 49B. ¹H NMR (300 MHz, CDCl₃) δ 7.38 (1H, m), 7.18 (1H, d, J=7Hz), 7.15 (1H, m), 7.00 (1H, d, J=2Hz), 6.95 (1H, m), 6.86 (1H, dd, J=7Hz, 2Hz), 6.75 (1H, d, J=8Hz), 5.93 (1H, d, J=4Hz), 5.92 (1H, d, J=4Hz), 3.94 (1H, d, J=14Hz), 3.63 (1H, m), 3.42 (3H, m), 3.35-2.95 (5H, m), 2.87 (1H, d, J=14Hz), 1.44 (3H, m), 1.27 (3H, m), 1.10 (2H, m), 0.88 (3H, t, J=7Hz), 0.81 (3H, t, J=7Hz). MS (DCI/NH₃) m/e 499 (M+H)⁺. Anal.calc. for C₂₈H₃₅FN₂O₅: C, 67.45; H, 7.08; N, 5.62. Found: C, 67.32; H, 7.05; N, 5.40.

Example 168*trans,trans*-1-(4-N,N-Di(*n*-butyl)aminophenyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

4-Nitro-1-fluorobenzene, ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate (the compound resulting from Example 6A), and diisopropylethylamine are heated in dioxane to give ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-nitrophenyl)-pyrrolidine-3-carboxylate. The nitro compound is hydrogenated to give the corresponding aminophenyl compound. The aminophenyl compound is reacted with butyraldehyde and sodium cyanoborohydride according to the method of Borch, J. Am Chem. Soc. 93: 2897 (1971) to give the corresponding N,N-dibutylaminophenyl compound. Hydrolysis with sodium hydroxide using the method of Example 1D affords the title compound.

Example 169*trans,trans*-1-(2-N,N-Dibutylaminopyrimidin-4-yl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

2-(Dibutylamino)-4-chloropyrimidine is prepared from 2,4-dichloropyrimidine according to the method of Gershon, J. Heterocyclic Chem. 24: 205 (1987) and reacted with ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate (the compound resulting from Example 6A) and diisopropylethylamine in dioxane with heating to give the intermediate ethyl ester, which is hydrolyzed with sodium hydroxide using the method of Example 1D to the title compound.

Examples 170-266

Using the procedures described in Examples 1, 4, 5, 7, 8 and 9 and Scheme X, the following compounds can be prepared.

Ex. No.	Name
170	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(isopropylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
171	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(ethylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
172	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(1-methylpropylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
173	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(phenylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
174	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(piperidinylcarbonylmethyl)-pyrrolidine-3-carboxylic acid;
175	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(1-(propylaminocarbonyl)ethyl)-pyrrolidine-3-carboxylic acid;
176	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(α -(propylaminocarbonyl)benzyl)-pyrrolidine-3-carboxylic acid;
177	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(bis-(propylaminocarbonyl)methyl)-pyrrolidine-3-carboxylic acid;
178	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(propylaminocarbonyl)ethyl)-pyrrolidine-3-carboxylic acid;
179	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminosulfonylmethyl)-pyrrolidine-3-carboxylic acid;
180	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-phenethyl)-pyrrolidine-3-carboxylic acid;
181	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(pentanoylmethyl)-pyrrolidine-3-carboxylic acid;
182	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(benzoylmethyl)-pyrrolidine-3-carboxylic acid;
183	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(hexyl)-pyrrolidine-3-carboxylic acid;
184	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-hexynyl)-pyrrolidine-3-carboxylic acid;
185	<i>trans,trans</i> -2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propoxymethylcarbonyl)-pyrrolidine-3-carboxylic acid;

- 186 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(phenylacetyl)-pyrrolidine-3-carboxylic acid;
- 187 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(anilinylicarbonyl)-pyrrolidine-3-carboxylic acid;
- 188 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-acetylaminoethyl)-pyrrolidine-3-carboxylic acid;
- 189 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-phenoxyethyl)-pyrrolidine-3-carboxylic acid;
- 190 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-benzodioxanylethyl)-pyrrolidine-3-carboxylic acid;
- 191 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-tetrahydrofuranylmethyl)-pyrrolidine-3-carboxylic acid;
- 192 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(propylaminocarbonylamino)ethyl)-pyrrolidine-3-carboxylic acid;
- 193 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(propylaminocarbonylamino)ethyl)-pyrrolidine-3-carboxylic acid;
- 194 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-oxohex-1-enyl)-pyrrolidine-3-carboxylic acid;
- 195 *trans,trans*-2-(2,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 196 *trans,trans*-2-(2-Carboxy-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 197 *trans,trans*-2-(2-Aminocarbonyl-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 198 *trans,trans*-2-(2-Methanesulfonamido-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 199 *trans,trans*-2-(2-Aminocarbonylmethoxy-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 200 *trans,trans*-2-(2-Methoxyethoxy-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

- 201 *trans,trans*-2-(2-Carboxymethoxy-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 202 *trans,trans*-2-(4-Methoxy-2-tetrazolylmethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 203 *trans,trans*-2-(2-Allyloxy-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 204 *trans,trans* 2,4-Bis(4-methoxyphenyl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 205 *trans,trans* 2,4-Bis(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 206 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 207 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxole-5-yl)-1-(N-methyl-N-butylaminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 208 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(4-methoxyphenyl)aminocarbonyl)-3-pyrrolidine-3-carboxylic acid;
- 209 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-phenylaminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 210 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-allylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 211 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 212 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-isobutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 213 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-cyclopentylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

- 214 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(2-methoxyethyl)aminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 215 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-butoxyethylaminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 216 *trans,trans*-2-(1,3-Benzodioxol-5-yl)-4-(4-methoxyphenyl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 217 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,4-benzodioxan-6-yl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 218 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-isopropylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 219 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-ethylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 220 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(1-methylpropyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 221 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-phenylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 222 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(1-(N-methyl-N-propylaminocarbonyl)ethyl)-pyrrolidine-3-carboxylic acid;
- 223 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(α -(N-methyl-N-propylaminocarbonyl)benzyl)-pyrrolidine-3-carboxylic acid;
- 224 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 225 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxole-5-yl)-1-(N-ethyl-N-butylaminocarbonyl)-pyrrolidine-3-carboxylic acid;

- 226 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-(4-methoxyphenyl)aminocarbonyl)-3-pyrrolidine-3-carboxylic acid;
- 227 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-phenylaminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 228 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-allylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 229 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-isobutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 230 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-cyclopentylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 231 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-methoxyethylaminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 232 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-butoxyethylaminocarbonyl)-pyrrolidine-3-carboxylic acid;
- 233 *trans,trans*-2-(1,3-Benzodioxol-5-yl)-4-(4-methoxyphenyl)-1-(N-ethyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 234 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,4-benzodioxan-6-yl)-1-(N-ethyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 235 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-isopropylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 236 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-diethylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 237 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-(1-methylpropyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 238 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-phenylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

- 239 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(1-(N-ethyl-N-propylaminocarbonyl)ethyl)-pyrrolidine-3-carboxylic acid;
- 240 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(α -(N-ethyl-N-propylaminocarbonyl)benzyl)-pyrrolidine-3-carboxylic acid;
- 241 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-isobutylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 242 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-cyclohexylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 243 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-dipropylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 244 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(isobutyloxyethyl)-pyrrolidine-3-carboxylic acid;
- 245 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(butylsulfonyl)-pyrrolidine-3-carboxylic acid;
- 246 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(isopropylsulfonylaminoethyl)-pyrrolidine-3-carboxylic acid;
- 247 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(ethoxymethylcarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 248 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-ethylbutyrylmethyl)-pyrrolidine-3-carboxylic acid;
- 249 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(3,4-dimethoxybenzyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 250 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[(1R)-1-(N-methyl-N-propylaminocarbonyl)butyl]-pyrrolidine-3-carboxylic acid;
- 251 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[(1S)-1-(N-methyl-N-propylaminocarbonyl)butyl]-pyrrolidine-3-carboxylic acid;
- 252 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-isopropoxypropyl)-pyrrolidine-3-carboxylic acid;

- 253 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(5-methylhexyl)-pyrrolidine-3-carboxylic acid;
- 254 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(5-methyl-2-hexenyl)-pyrrolidine-3-carboxylic acid;
- 255 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(5-methyl-4-hexenyl)-pyrrolidine-3-carboxylic acid;
- 256 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3,5-dimethyl-2-hexenyl)-pyrrolidine-3-carboxylic acid;
- 257 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-methyl-N-isobutrylamino)ethyl)-pyrrolidine-3-carboxylic acid;
- 258 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-(2,2-dimethylpropyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 259 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-ethyl-N-butylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 260 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-methyl-N-benzylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 262 *trans,trans*-2-(4-Methoxyphenyl)-4-(5-indanyl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 262 *trans,trans*-2-(4-Methoxyphenyl)-4-(2,3-dihydrobenzofuran-5-yl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 263 *trans,trans*-2-(4-Methoxyphenyl)-4-(1-methylindol-5-yl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 264 *trans,trans*-2-(4-Methoxyphenyl)-4-(2-naphthyl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 265 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,2-dimethoxy-4-phenyl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 266 *trans,trans*-2-(4-Methoxyphenyl)-4-(1-methoxy-3-phenyl)-1-(N-methyl-N-propylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

Examples 267-288

Following the procedures described in Example 1 and Scheme II, the following compounds can be prepared.

- 267 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(propylaminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 268 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(aminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 269 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(4-fluorobenzyl)-piperidine-4-carboxylic acid;
- 270 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(2-ethoxyethyl)-piperidine-4-carboxylic acid;
- 271 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(2-propoxyethyl)-piperidine-4-carboxylic acid;
- 272 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-[2-(2-methoxyethoxy)ethyl]-piperidine-4-carboxylic acid;
- 273 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-[2-(2-pyridyl)ethyl]-piperidine-4-carboxylic acid;
- 274 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(morpholin-4-ylcarbonyl)-piperidine-4-carboxylic acid;
- 275 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(butylaminocarbonyl)-piperidine-4-carboxylic acid;
- 276 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(4-methoxyphenylaminocarbonyl)-3-piperidine-4-carboxylic acid;
- 277 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-acetyl-piperidine-3-carboxylic acid;
- 278 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(2-furoyl)-piperidine-3-carboxylic acid;
- 279 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(phenylaminocarbonyl)-piperidine-4-carboxylic acid;
- 280 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(allylaminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 281 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(n-butylaminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 282 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(N-n-butyl-N-methylaminocarbonylmethyl)-piperidine-4-carboxylic acid;

- 283 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(pyrrolidin-1-ylcarbonylmethyl)-piperidine-4-carboxylic acid;
- 284 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(isobutylaminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 285 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(cyclopentylaminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 286 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(morpholin-4-ylaminocarbonylmethyl)-piperidine-4-carboxylic acid;
- 287 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(2-phenoxyethyl)-piperidine-4-carboxylic acid;
- 288 *trans,trans*-3-(4-Methoxyphenyl)-5-(1,3-benzodioxol-5-yl)-1-(methoxyethylaminocarbonyl)-piperidine-4-carboxylic acid.

Example 289

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-dibutylaminophenyl)-pyrrolidine-3-carboxylic acid

- 5 4-Nitro-fluorobenzene, ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate (example 6A) and di-isopropyl ethylamine are heated in dioxane to give ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(4-nitrophenyl)-pyrrolidine-3-carboxylate. The nitro compound is hydrogenated to the corresponding aminophenyl compound. This is reacted with butyraldehyde and sodium cyanoborohydride
- 10 according to the method of Borch (J. Am Chem. Soc., 93, 2897, 1971) to give the corresponding N,N-dibutylaminophenyl compound, which is hydrolyzed with sodium hydroxide using the method of example 1D to give the title compound.

Example 290

- 15 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-dibutylamino-pyrimidine-4-yl)-pyrrolidine-3-carboxylic acid

- 2-(Dibutylamino) 4-chloropyrimidine is prepared from 2,4-dichloropyrimidine according to the method of Gershon (J. Heterocyclic Chem. 24, 205, 1987). This compound, ethyl *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate
- 20 (example 6A), and di-isopropyl ethylamine are heated in dioxane to give the intermediate ethyl ester, which is hydrolyzed with sodium hydroxide using the method of example 1D to give the title compound.

Example 291

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-butyl-N-phenylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared according to the general procedure of Example 1.

¹H NMR (CD₃OD) : δ 0.87 (t, 3H, J=8); 1.2-1.35 (m, 2H); 1.35-1.5 (m, 2H); 2.78 (m, 2H); 3.10

5 (t, 1H, J=9); 3.26 (d, 1H, J=15); 3.44 (dd, 1H, J=5, 10); 3.5-3.7 (m, 3H); 3.77 (m, 1H); 3.78 (s, 3H); 5.93 (s, 2H); 6.7-6.9 (m, 4H); 7.0-7.2 (m, 5H); 7.4 (m, 3H). MS (DCI/NH₃): m/e 531 (M+H)⁺.

Anal calcd for C₃₁H₃₄N₂O₆: C, 70.17; H, 6.46; N, 5.28. Found: C, 70.36; H, 6.52; N, 4.99.

Example 292

10 Sodium *trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylate*

Example 292A

Ethyl 3-(4-methoxyphenyl)-3-oxopropionate

15 Simultaneous reactions were run in both a 65-L reactor and a 35-L reactor that share the same reflux system. A nitrogen atmosphere was maintained in both. 4.0 kg (100 moles) of 60% sodium hydride in mineral oil and 32 L toluene were charged into the ambient temperature reactors. The mixture was agitated for 5 minutes and allowed to settle. 20 L of the toluene solution was aspirated. 28 L of toluene was added, agitated for 5 minutes, 20 allowed to settle and 28 L of the toluene solution was aspirated. 68 L of toluene and 8.4 L (69.7 moles) diethyl carbonate were added. The agitation was begun and the flow of Syltherm (Note 4) in reactor jackets was initiated. A solution of 5.0 kg (33.3 moles) 4-methoxyacetophenone in 12 L toluene was added over 20 minutes. When additions were complete, the jacket temperature was reduced to 10° C and stirring continued for 16 hours. A 25 solution of 6.7 L (117 moles) glacial acetic acid in 23 L deionized water was fed at the same rate that was previously used for the acetophenone solution. When addition was complete, agitation was stopped and the layers separated. The aqueous layer was washed once with 13 L toluene. The combined organic layers were washed twice with 6.7 L portions of 7% (w:w) aqueous sodium bicarbonate. The toluene solution was washed once with 6.7 L of 23% 30 (w:w) aqueous sodium chloride. The organic solution was dried over 10 kg sodium sulfate, filtered, and the solvent removed on the rotary evaporator to provide the desired product.

Example 292B

3,4-Methylenedioxy-1-(2-nitroethenyl)-benzene

35 In a 45-L cryogenic reactor with a contoured, anchor stirrer was dissolved 5.537 kg (36.9 moles) piperonal in 9 L methanol and 2.252 kg (36.9 moles) nitromethane at 15°-20° C. The jacket temperature was set to -5° C and the reaction solution cooled to a temperature of

+3.5° C. A 21° C solution of 3.10 kg (38.8 moles) 50% (w:w) aqueous sodium hydroxide diluted with 3.7 L water was pumped in. The reaction temperature was maintained between 10°-15° C. When addition was complete, the jacket temperature was reset to 1° C and stirring continued for 30 minutes. A mixture of 7 kg ice in 19 L water was added to dissolve most of the solid. The reaction mixture was filtered through canvas and then a 27R10SV Honeycomb filter. The filtered solution was metered into a 21° C mixture of 7.4 L concentrated hydrochloric acid in 11.1 L deionized water. The final reaction temperature was 26° C. The resulting product was centrifuged and washed until the wash pH rose to at least 6 (by pH indicating paper). The crude product was dissolved in 92 L dichloromethane and the layers separated. The aqueous layer was washed once with 8 L dichloromethane. The combined organics were dried over 1.32 kg magnesium sulfate and filtered through Whatman #1 paper. The volume was reduced to 20% and the solution cooled to 4° C. Filtration through Whatman #1 paper, followed by ambient temperature drying *in vacuo* with an air leak afforded 1.584 kg (22%) of a first crop. Concentration of the MLS to 25% followed by similar cooling, filtration, and drying afforded 0.262 kg (4%) of a second crop. The yellow product darkened on standing in light and air.

Example 292C

Ethyl 2-(4-methoxybenzoyl)-3-(1,3-benzodioxol-5-yl)-4-nitro-butanate

Into a 45-L stirred reactor at ambient temperature were charged 5.819 kg (30.1 moles) 3,4-methylenedioxy-1-(2-nitroethenyl)-benzene and 24 L ethyl acetate. A solution of 5.355 kg (24.1 moles) ethyl 3-(4-methoxyphenyl)-3-oxopropionate in 16 L ethyl acetate was added. 280 g (275 ml, 1.84 moles) of 1,8-diaza-bicyclo[5.4.0]undec-7-ene in four equal portions was added over a 2.5 hour period. The reaction mixture was filtered through dicalite and the resulting filtered solution was used in the next step without any further purification.

Example 292D

Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-4,5-dihydro-3H-pyrrol-3-carboxylate

The product of Example 292C (1316 ml solution consisting of 300 g Ethyl 2-(4-methoxybenzoyl)-3-(3,4-methylenedioxyphenyl)-4 nitrobutanoate in ethyl acetate) was added to a glass reactor containing RaNi # 28 (300 g). The reaction mixture was shaken under a hydrogen environment of 4 atm at room temperature for 18 hours and filtered through a nylon 0.20 micron 47 mm millipore.

The filtrate was concentrated to 1.4 kg of dark solution and purified by normal phase silica gel chromatography eluting with 85:15, hexanes: ethyl acetate. The pure fractions were combined and concentrated (as above) until crystals formed. The solution was cooled to 0° C and filtered. The solid was washed with 2 L of 85:15, hexane: ethyl acetate (0° C). The solids

were dried *in vacuo* at 50° C to a constant weight of 193.4 g (21% yield, melting point 80-81° C) of the title compound. A further 200 g (23% yield) of product was obtained from the mother liquors.

5

Example 292E

Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine 3-carboxylate

Into a 12-L flask equipped with magnetic stirring, addition funnel, temperature probe, and nitrogen inlet was charged 0.460 kg ethyl 2-(4-methoxyphenyl)-4-(3,4-methylenedioxyphenyl)-4,5-dihydro-3H-pyrrole-3-carboxylate (1.25 mol). The reaction vessel was degassed with nitrogen. Absolute 3.7 L ethanol and 1.12 L of THF were added. 31 mg bromocresol green and 94.26g sodium cyanoborohydride (1.5 mol) were added. A solution containing 400 mL absolute ethanol and 200 mL of 12 M HCl was then added. The reaction mixture was stirred for 30 minutes after addition was complete. After the starting material was consumed, 0.5 L of 7% aq. NaHCO₃ was added. The reaction mixture was concentrated and diluted with 5 L ethyl acetate. The organic layer was washed twice with 2 L of 7% aq. NaHCO₃ and once with 2.5 L of 23% aq. NaCl, the dried over 190g MgSO₄, filtered, and concentrated to give 447 g of the title compound as a thick yellow oil.

20

Example 292F

Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl) pyrrolidine 3-carboxylate

Into a 22-L flask equipped with overhead stirring, nitrogen inlet, and condenser was charged ethyl 2-(4-methoxyphenyl)-4-(3,4-methylenedioxyphenyl)-pyrrolidine-3-carboxylate (2.223 kg, 6.02 mol). The reaction vessel was degassed with nitrogen. 13.2 L of acetonitrile, 3.66 L diisopropylethylamine (2.71 kg, 20.9 mol), and 1.567 kg dibutylamidomethyl bromide (6.26 mol) were added. The mixture was refluxed at 78° C for 17 hrs. After the disappearance of starting material, the mixture was concentrated until crystals formed. The solid was filtered and washed with 4 L ethyl acetate (0° C). Concentrating of the filtrate was continued as above until all volatiles were removed. The residue was diluted with 40 L ethyl acetate and washed with 20 L deionized water. The organic layer was washed with 8 L of 23% aq. NaCl and dried over 0.399 kg MgSO₄ and filtered. Concentration as above provided 3.112 kg (96 % yield) of the title compound as a dark oil.

35

Example 292G

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine 3-carboxylate and preparation of trans,trans-2-(4-methoxyphenyl)-4-(3,4-dioxyphenyl)-pyrrolidine-3-carboxylic acid ethyl ester

Into a 35-L reactor equipped with overhead stirring, nitrogen inlet, and condenser was charged 3.112 kg ethyl 2-(4-methoxyphenyl)-4-(3,4-methylenedioxyphenyl)-pyrrolidine 3-carboxylate (5.78 mol). 16.4 L of absolute ethanol was added and the reaction vessel was degassed with nitrogen. 0.115 kg of sodium ethoxide (1.69 mol) was added and the mixture was refluxed at 79° C for 1 hr. The mixture was cooled to 15° C and 5 L of 7.6 M NaOH solution (38.1 mol) was added. The mixture was stirred at 15° C for 18 hrs. The solvent was evaporated and the residue dissolved in 15.8 L of deionized water and extracted with 28 L of ether. The ether solution was washed with 9.5 L deionized water. The aqueous wash was extracted with 3 L ether. 0.340 L of 12 M HCl was added to the aqueous layer. The aqueous layer was extracted with 24 L of ethyl acetate. The organic layer was washed with 9 L of 23% aq. NaCl, dried with 0.298 kg MgSO₄, filtered, and concentrated to give 2.132 kg of a dark oil. The oil was triturated with 18 L ether. The undesired solids were filtered and saved for later use. The mother liquors were concentrated to obtain 1.102 kg of light foam. The foam was dissolved in 5.5 L ethyl acetate with heating to 65° C. 14 L hexane was added slowly enough to keep the solution refluxing. The reaction mixture was cooled to 10° C and filtered. The crystals were washed with 2 L ether (0° C) and dried to constant weight *in vacuo* at 50° C to give 0.846 kg (43% yield, melting point 119-120) of crude product, which was further purified by normal phase silica gel chromatography.

Example 292H

Sodium *trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl) pyrrolidine 3-carboxylate

Into a 20-L flask was charged *trans,trans* 2-(4-methoxyphenyl)-4-(3,4-methylenedioxyphenyl)-1-(N,N-dibutylamino- carbonyl methyl) pyrrolidine 3-carboxylic acid (0.927 kg, 1.819 mol). A solution of 0.0720 kg NaOH (1.80 mol) dissolved in 4.65 L methanol was added. The reaction mixture was concentrated to an oil. Pentane (4 L) was added and the solution concentrated again. Pentane (4 L) was added again and concentration of this solution gave a light tan foam. The foam was dried *in vacuo* at 50° C to a constant weight of 0.937 kg (97% yield) of the title compound.

Example 293

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[decahydroisoquinolin-2-carbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) shows a mixture of isomers. MS (DCI/NH₃) *m/z* 521. Anal calcd for C₃₀H₃₆N₂O₆ . 1.3 TFA: C, 58.54; H, 6.62; N, 4.19 . Found: C, 58.34; H, 5.58; N, 4.00 .

Example 294

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[3,3-dimethylpiperidinyl-carboxylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) indicates presence of rotamers. δ 0.84 (s, 3H), 0.86 (s, 3H), 1.35-1.6 (m, 4H), 3.83 (s, 3H), 5.96 (s, 2H), 6.81 (d, 1H, J=8), 6.90 (dd, 1H, J=1,8), 7.01 (d, 2H, J=9), 7.03 (s, 1H), 7.47 (d, 2H, J=9). MS (DCI/NH₃) m/z 495. Anal calcd for C₂₈H₃₄N₂O₆ · 1.4 TFA: C, 56.55; H, 5.45; N, 4.28. Found: C, 56.52; H, 5.83; N, 4.26.

Example 295

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-isobutoxycarbonylamino)ethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods detailed in Example 61, but substituting propylamine for methylamine in Example 61B and isobutyl chloroformate for isobutyl chloride in Example 61C. The crude product was purified by trituration with 1:1 diethyl ether/ hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (t, 3H, J=7), 0.92 (m, 3H), 1.43 (h, 2H, J=7Hz), 1.7-1.9 (m, 1H), 2.72 (m, 1H), 2.90 (m, 2H), 3.10 (m, 2H), 3.25 (m, 2H), 3.40 (m, 1H), 3.55 (m, 1H), 3.62 (m, 1H), 3.7-3.9 (m, 2H), 3.78 (s, 3H), 5.95 (s, 2H), 6.72 (d, 1H, J= 8Hz), 6.82 (m, 3H), 7.00 (s, 1H), 7.30 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 527 (M+H)⁺. Anal calcd for C₂₉H₃₈N₂O₆ · 0.5 H₂O: C, 65.03; H, 7.34; N, 5.23. Found: C, 65.13; H, 6.96; N, 4.95.

Example 296

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[1,2,3,4-tetrahydroisoquinolin-2-carboxylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) indicates presence of rotamers. δ 2.97 (m, 2H), 4.68 (s, 3H), 5.97 (s, 2H), 6.83 (d, 1H, J=8), 6.9-7.0 (m, 3H), 7.03 (d, 1H, J=2), 7.1-7.3 (m, 4H), 7.4-7.5 (m, 2H). MS (DCI/NH₃) m/z 515.

Example 297

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-dimethylaminocarbonylamino)ethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared by the methods detailed in Example 61, but substituting propylamine for methylamine in Example 61B and dimethylcarbonyl chloride for isobutyl chloride in Example 61C. The crude product was purified by preparative HPLC

(Vydac μ C18) eluting with a 10-70% gradient of CH_3CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ^1H NMR (CDCl_3 , 300 MHz) δ 0.70 (t, 3H, $J=7$), 1.28 (m, 2H), 2.75 (s, 3H), 2.82 (m, 2H), 3.1-3.45 (m, 4H), 3.70 (m, 1H), 3.80 (s, 3H), 3.90 (m, 3H), 4.72 (m, 1H), 5.95 (s, 2H), 6.75 (d, 1H, $J=8\text{Hz}$), 6.87 (m, 3H), 7.05 (s, 1H), 7.40 (d, 2H, $J=8\text{Hz}$). MS (DCI/NH_3) m/e 498 ($\text{M}+\text{H}$) $^+$. Anal calcd for $\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_6 \cdot 1.25\text{ TFA}$: C, 55.35; H, 5.71; N, 6.56. Found: C, 55.41; H, 5.71; N, 6.41.

Example 298

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(4-nitrobenzenesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a yellow solid. m.p. 85-87°C. ^1H NMR (CDCl_3 , 300 MHz) δ 0.77 (t, $J=7.5\text{Hz}$, 3H), 1.38 (sextet, $J=7.5\text{Hz}$, 2H), 2.20-2.29 (m, 1H), 2.57-2.66 (m, 1H), 2.82-3.15 (m, 4H), 3.22 (t, $J=7.5\text{Hz}$, 2H), 3.38 (dd, $J=3\text{Hz}$, $J=9\text{Hz}$, 1H), 3.49-3.57 (m, 1H), 3.59 (d, $J=9\text{Hz}$, 1H), 3.83 (s, 3H), 5.96 (s, 2H), 6.73 (d, $J=8\text{Hz}$, 1H), 6.82 (dd, $J=1\text{Hz}$, $J=8\text{Hz}$, 1H), 6.87 (d, $J=9\text{Hz}$, 2H), 6.98 (d, $J=1\text{Hz}$, 1H), 7.27 (d, $J=9\text{Hz}$, 2H), 7.82 (d, $J=9\text{Hz}$, 2H), 8.23 (d, $J=9\text{Hz}$, 2H). MS (DCI/NH_3) m/e 612 ($\text{M}+\text{H}$) $^+$.

Example 299

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 59-61°C. ^1H NMR (CDCl_3 , 300MHz) δ 0.79 (t, $J=7.5\text{Hz}$, 3H), 0.90 (t, $J=6\text{Hz}$, 3H), 1.26-1.32 (m, 4H), 1.43 (sextet, $J=7.5\text{Hz}$, 2H), 1.67-1.76 (m, 2H), 2.23-2.32 (m, 1H), 2.70-3.08 (m, 7H), 3.15-3.32 (m, 2H), 3.42 (dd, $J=3\text{Hz}$, $J=9\text{Hz}$, 1H), 3.52-3.57 (m, 1H), 3.63 (d, $J=9\text{Hz}$, 1H), 3.80 (s, 3H), 5.95 (s, 2H), 6.73 (d, $J=7.5\text{Hz}$, 1H), 6.83 (dd, $J=1\text{Hz}$, $J=7.5\text{Hz}$, 1H), 6.87 (d, $J=8\text{Hz}$, 2H), 7.00 (d, $J=1\text{Hz}$, 1H), 7.32 (d, $J=8\text{Hz}$, 2H). MS (DCI/NH_3) m/e 561 ($\text{M}+\text{H}$) $^+$.

Example 300

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(4-trifluoromethoxybenzenesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 122-124°C. ^1H NMR (CD_3OD , 300MHz) δ 0.75 (t, $J=7.5\text{Hz}$, 3H), 1.26-1.45 (m, 2H), 2.96-3.08 (m, 2H), 3.23 (bs, 2H), 3.35-3.45 (m, 2H), 3.52 (t, $J=10\text{Hz}$, 1H), 3.81 (d, $J=9\text{Hz}$, 2H), 3.86 (s, 3H), 3.92 (t, $J=9\text{Hz}$, 1H), 4.63 (d, $J=10\text{Hz}$, 1H), 5.97 (s, 2H), 6.82 (d,

J=9Hz, 1H), 6.93 (dd, J=3Hz, J=9Hz, 1H), 7.06-7.08 (m, 3H), 7.46 (d, J=9Hz, 2H), 7.56 (d, J=9Hz, 2H), 7.89 (d, J=9Hz, 2H). MS (DCI/NH₃), m/e 651 (M+H)⁺.

Example 301

5 trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-methyl-2-propenesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 69-71°C.. ¹H NMR (CDCl₃, 300MHz) δ 0.79 (t, J=7.5Hz, 3H), 1.93 (sextet, J+7.5Hz, 2H), 1.92 (s, 3H), 2.25-2.35 (m, 1H), 2.68-2.77 (m, 1H), 2.85-3.28 (m, 7H), 3.40 (d, J=9Hz, 1H), 3.52-3.68 (m, 2H), 3.66 (d, J=9Hz, 1H), 3.80 (s, 3H), 4.92 (s, 1H), 5.07 (s, 1H), 5.97 (s, 2H), 6.74 (d, J=7Hz, 1H), 6.82-6.89 (m, 3H), 7.01 (s, 1H), 7.33 (d, J=9Hz, 2H). MS (DCI/NH₃), m/e 545 (M+H)⁺.

Example 302

15 trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-ethylpiperidinyl-carbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) shows a mixture of isomers. δ 0.75 (t, 3H, J=7), 1.4-1.7 (m, 8H), 3.84 (s, 3H), 5.96 (s, 2H), 6.83 (d, 1H, J=8), 6.91 (d, 1H, J=8), 7.0-7.1 (m, 3H), 7.52 (d, 2H, J=9). MS (DCI/NH₃) m/z 495. Anal calcd for C₂₈H₃₄N₂O₆. 1.6 TFA: C, 55.35; H, 5.30; N, 4.14. Found: C, 55.26; H, 5.37; N, 4.01.

Example 303

25 trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-methylpropanesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 72-73°C. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=7.5Hz, 3H), 1.04 (d, J=6Hz, 6H), 1.44(q, J=7.5Hz, 2H), 2.15-2.33 (m, 2H), 2.57-2.75 (m, 2H), 2.84-3.08 (m, 3H), 3.12-3.21 (m, 1H), 3.23-3.45 (m, 1H), 3.43 (d, J=11Hz, 1H), 3.55-3.62 (m, 1H), 3.66 (d, J=9Hz, 1H), 3.80 (s, 3H), 5.95 (s, 2H), 6.75 (d, J=9Hz, 1H), 6.83 (dd, J=1Hz, J=9Hz, 1H), 6.87(d, J=9Hz, 2H), 7.02 (d, J=1Hz, 1H), 7.33 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 547 (M+H)⁺.

Example 304

35 trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-heptanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 58-59°C. ¹H NMR (CDCl₃, 300MHz) δ 0.80(t, J=7.5Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.23-1.36 (m, 8H), 1.94 (q, J=7.5Hz, 2H), 1.71 (quintet, J=7Hz, 2H), 2.23-2.32 (m, 1H), 2.70-3.09 (m, 7H), 3.13-3.32 (m, 2H), 3.43 (dd, J=3Hz, J=9Hz, 1H), 3.52-3.58 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.80 (s, 3H), 5.96 (s, 2H), 6.73 (d, J=7Hz, 1H), 6.83 (dd, J=1Hz, J=7Hz, 1H), 6.87 (d, J=9Hz, 2H), 7.01 (d, J=1Hz, 1H), 7.32 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 589 (M+H)⁺.

Example 305

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-ethyl-N-ethoxycarbonylamino)ethyl]-pyrrolidine-3-carboxylic acid

Prepared by the methods detailed in Example 61, but substituting ethylamine for methylamine in Example 61B and ethyl chloroformate for isobutyryl chloride in Example 61C. The crude product was purified by preparative HPLC (Vydac μC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The desired fractions were lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.90 (t, 3H, J=7), 1.22 (m, 3H), 3.0-3.2 (m, 4H), 3.42 (m, 2H), 3.78 (s, 3H), 3.82 (m, 4H), 4.10 (q, 2H, J=7Hz), 3.5 (br s, 1H), 5.97 (dd, 2H, J=1,7Hz), 6.72 (d, 1H, J=8Hz), 6.84 (m, 3H), 7.00 (s, 1H), 7.42 (d, 2H, J=8Hz). MS (DCI/NH₃) m/e 485 (M+H)⁺. Anal calcd for C₂₆H₃₂N₂O₇ · 1.2 TFA: C, 54.90; H, 5.39; N, 4.51. Found: C, 55.01; H, 5.36; N, 4.56.

Example 306

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-hexanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 59-60°C. ¹H NMR (CDCl₃, 300MHz) δ 0.80(t, J=7.5Hz, 3H), 0.89(t, J=7Hz, 3H), 1.25-1.36(m, 6H), 1.53(sextet, J=7.5Hz, 2H), 1.72(quintet, J=7Hz, 2H), 2.23-2.32(m, 1H), 2.72-3.08(m, 7H), 3.15-3.32(m, 2H), 3.43(d, J=9Hz, 1H), 3.55-3.62(m, 1H), 3.65 (d, J=10Hz, 1H), 3.80(s, 3H), 5.96(s, 2H), 6.74(d, J=7.5Hz, 1H), 6.82(d, J=7.5Hz, 1H), 6.87(d, J=9Hz, 2H), 7.01(s, 1H), 7.32(d, J=9Hz, 2H). MS (DCI/NH₃), m/e 575 (M+H)⁺.

Example 307

trans-trans-2-(4-Ethylphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in examples 1 and 49, substituting ethyl 4-ethylbenzoate for ethyl 4-ethylbenzoate (prepared by the method of Krapcho et al., Org.

Syn. 47, 20 (1967) starting with 4'-ethylacetophenone) in procedure 49B. NMR (CDCl₃, 300 MHz) δ 7.31 (2H, d, J=8Hz), 7.16 (2H, d, J=8Hz), 7.03 (1H, d, J=3Hz), 6.86 (1H, dd, J=8&3Hz), 6.73 (1H, d, J=9Hz), 5.94 (1H, d, J=4Hz), 5.92 (1H, d, J=4Hz), 3.77 (1H, d, J=9Hz), 3.60 (1H, m), 3.53-3.23 (5H, m), 3.13-2.90 (4H, m), 2.73 (1H, d, J=14Hz), 2.62 (2H, q, J=9Hz), 1.45 (2H, m), 1.40-1.10 (6H, m), 1.02 (2H, m), 0.87 (3H, t, J=7Hz), 0.78 (3H, t, J=7Hz). m/e (DCl, NH₃) 509 (MH⁺) Anal.calc. for C₃₀H₄₀N₂O₅ C 70.84, H 7.93, N 5.51. Found C 70.80, H 7.85, N 5.25 .

Example 308

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-(2-chloroethoxy)carbonylamino)ethyl]-pyrrolidine-3-carboxylic acid

Prepared by the methods detailed in Example 61, but substituting propylamine for methylamine in Example 61B and 2-chloroethyl chloroformate for isobutyl chloride in Example 61C. The crude product was purified by titration with 1:1 diethyl ether/ hexane. The resulting solid was dissolved in CH₃CN and water and lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (t, 3H, J=7), 1.22 (m, 3H), 2.15 (m, 1H), 2.75 (m, 1H), 2.85 (m, 1H), 3.1 (m, 2H), 3.25 (m, 2H), 3.5 (m, 3H), 3.65 (m, 2H), 3.80 (s, 3H), 4.18 (m, 1H), 4.30 (m, 1H), 5.98 (s, 2H), 6.72 (m, 1H), 6.82 (m, 3H), 7.00 (m, 1H), 7.30(m, 2H). MS (DCl/NH₃) m/e 533 (M+H)⁺. Anal calcd for C₂₇H₃₃N₂O₇Cl: C, 60.84; H, 6.24; N, 5.26. Found: C, 60.48; H, 6.04; N, 5.10.

Example 309

trans-trans-2-(2-Methoxyethyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in example 1, substituting ethyl 5-methoxy-3-oxopentanoate for ethyl 4-methoxybenzoylacetate in Example 1A. The title compound is a yellow foam. ¹H NMR (CDCl₃, 300 MHz) δ 0.91 (t, J=7Hz) and 0.95 (t, J=7Hz, 6H total), 1.28-1.41 (br m, 4H), 1.45-1.63 (br m, 4H), 2.00-2.20 (br m, 2H), 3.06 (br t, J=9Hz, 1H), 3.30 (s) and 3.20-3.68 (br m, 11H total), 3.72-4.10 (br m, 4H), 5.92 (s, 2H), 6.72 (d, J=8.5Hz, 1H), 6.82 (dd, J=1.5, 8.5Hz, 1H), 6.91 (d, J=1.5Hz, 1H); MS (FAB) m/e 463 (M+H)⁺. Anal calcd for C₂₅H₃₈N₂O₅H₂O: C, 62.48; H, 8.39; N, 5.83. Found: C, 62.13; H, 8.15; N, 5.69.

Example 310

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-ethyl-N-n-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 57-58°C. ^1H NMR (CDCl_3 , 300 MHz) δ 0.89(t, J=7Hz, 3H), 1.06(t, J=7.5Hz, 3H), 1.26-1.37(m, 4H), 1.72(quintet, J=7.5Hz, 2H), 2.22-2.32(m, 1H), 2.71-2.96(m, 5H), 3.08-3.30(m, 4H), 3.95(d, J=9Hz, 1H), 3.53-3.60(m, 1H), 3.67(d, J=9Hz, 1H), 3.80(s, 1H), 5.97(s, 2H), 6.73(d, J=9Hz, 1H), 6.82(d, J=9Hz, 1H), 6.88(d, J=9Hz, 2H), 7.02(s, 1H), 7.33(d, J=9Hz, 2H). MS (CDI/NH_3) m/e 547 ($\text{M}+\text{H}^+$).

Example 311

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-dicyclohexylamino carbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD_3OD , 300 MHz) δ 1.0-2.0 (m, 20H), 3.0-3.1 (m, 2H), 3.80 (s, 3H), 5.95 (s, 2H), 6.75 (d, 1H, J=8), 6.86 (dd, 1H, J=2,8), 6.95 (d, 2H, J=9), 7.04 (d, 1H, J=2), 7.38 (d, 2H, J=9). MS (DCI/NH_3) m/z 563. Anal calcd for $\text{C}_{33}\text{H}_{42}\text{N}_2\text{O}_6 \cdot 0.5 \text{H}_2\text{O}$: C, 69.33; H, 7.58; N, 4.90. Found: C, 69.42; H, 7.29; N, 4.78.

Example 312

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-tert-butoxycarbonylamino)ethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 61, substituting propylamine for aqueous methylamine in Example 61B and di-tert-butylidicarbonate for isobutyl chloride in Example 61C. NMR (CD_3OD , 300 MHz) suggests presence of rotamers δ 0.81 (t, 3H, J=7), 1.2-1.5 (m, 11H), 3.78 (s, 3H), 5.92 (dd, 2H, J=1,2), 6.74 (d, 1H, J=8), 6.84 (dd, 1H, J=2,8), 6.92 (d, 2H, J=9), 6.99 (bd s, 1H), 7.35 (d, 2H, J=9). MS (DCI/NH_3) m/z 527. Anal calcd for $\text{C}_{29}\text{H}_{38}\text{N}_2\text{O}_7$: C, 66.14; H, 7.27; N, 5.32. Found: C, 66.05; H, 7.36; N, 5.15.

Example 313

trans-trans-2-(4-Methoxy-3-fluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the methods described in examples 1 and 43, using 4-methoxy-3-fluoro acetophenone in place of 4-methoxy acetophenone. m.p. 142-143 °C. NMR (CDCl_3 , 300 MHz) δ 0.82 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.03-1.50 (m, 8H), 2.82 (d, J=13Hz, 1H), 2.90-3.13 (m, 4H), 3.20-3.50 (m, 3H), 3.39 (d, J=13Hz, 1H), 3.55-3.65 (m, 1H), 3.82 (d, J=10Hz, 1H), 3.87 (s, 3H), 5.91 (dd, J=2Hz, 4Hz, 2H), 6.72 (d, J=8Hz, 1H), 6.83-6.91 (m, 2H), 6.99 (d, J=2Hz, 1H), 7.06 (m, 2H). Anal calcd for $\text{C}_{29}\text{H}_{37}\text{N}_2\text{O}_6\text{F}$: C, 65.89; H, 7.06; N, 5.30. Found: C, 65.82; H, 7.13; N, 5.29.

Example 314

trans,trans-2-(Propyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Example 314APropyl pentanesulfonamide

Pentane sulfonyl chloride (687 mg, 4.03 mmol) was dissolved in 5 mL CH₂Cl₂ and added to an ice-cooled solution of n-propylamine (0.40 mL, 4.82 mmol) and ethyldiisopropylamine (0.85 mL, 4.88 mmol) in 5 mL CH₂Cl₂ under a nitrogen atmosphere. The reaction was stirred at 0 °C for 30 min, then at 25 °C for 4 h. The solution was partitioned between 20 mL of 1.0 M aqueous NaHSO₄ and 25 mL CH₂Cl₂. The organic phase was washed sequentially with 25 mL H₂O and 25 mL brine, then dried (Na₂SO₄), filtered, and concentrated in vacuo to provide 739 mg (3.83 mmol, 95%) of the title compound as a white solid. TLC (25% EtOAc-hexane) R_f 0.23; ¹H NMR (CDCl₃, 300 MHz) δ 0.92 (t, J=7Hz, 3H), 0.97 (t, J=7Hz, 3H), 1.28-1.50 (br m, 4H), 1.52-1.68 (m, 2H), 1.75-1.90 (br m, 2H), 2.98-3.06 (m, 2H), 3.08 (q, J=6Hz, 2H), 4.10-4.23 (br m, 1H); MS (DCI/NH₃) m/e 211 (M+NH₄)⁺.

Example 3149B

Ethyl *trans,trans*-4-(1,3-benzodioxol-5-yl)-1-(2-bromoethyl)-2-propylpyrrolidine-3-carboxylate

The title compound was prepared according the procedure of Example 61A, substituting the compound of Example 94B for the pyrrolidine mixture.

Example 314C

Ethyl*trans,trans*-2-(Propyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylate

A solution of the compound of Example 314A (6.6 mg, 34 μmol) in 0.1 mL DMF was treated with sodium hydride (2 mg, 60% oil dispersion, 1.2 mg NaH, 50 μmol). The resulting mixture was stirred at room temperature for 15 min, then a solution of the compound of Example 189B (9.0 mg, 22 μmol) in 0.1 mL DMF was added, followed by 0.5 mg of tetra-n-butylammonium iodide. The reaction was sealed under argon and stirred at 60 °C overnight. The reaction was concentrated under high vacuum, and the residue was partitioned between 2 mL of saturated aqueous NaHCO₃, 1 mL water and 5 mL EtOAc. The organic phase was washed with 1 mL brine, dried by passing through a plug of Na₂SO₄, and the filtrate

concentrated in vacuo to an oil. The crude product was purified by preparative TLC (silica gel, 8 x 20 cm, 0.25 mm thickness, eluting with 20% EtOAc-hexane, providing 8.4 mg (73%) of the title compound as a wax.

Example 314D

trans,trans-4-(1,3-benzodioxol-5-yl)-2-(Propyl)-1-(2-(N-propyl-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

The title compound was prepared according to the procedure of Example 71C. ¹H NMR (CDCl₃, 300 MHz) δ 0.88-1.00 (m, 9H), 1.20-1.55 (br m, 6H), 1.55-1.68 (m, 3H), 1.70-1.85 (br m, 2H), 1.90-2.16 (br m, 2H), 2.84-3.26 (br m, 6H), 3.26-3.90 (br m, 6H), 5.95 (s, 2H), 6.76 (d, J=8Hz, 1H), 6.79 (m, 1H), 6.93 (br s, 1H); HRMS (FAB) calcd for C₂₅H₄₁N₂O₆S (M+H)⁺ 497.2685, found 497.2679.

Example 315

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-dimethylsulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared as a white solid. m.p. 59-61°C. ¹H NMR (CDCl₃, 300MHz) δ 0.79 (t, J=7.5Hz, 3H), 1.45 (sextet, J=7.5Hz, 2H), 2.22-2.31 (m, 1H), 2.65 (s, 6H), 2.70-2.79 (m, 1H), 2.85-3.04 (m, 4H), 3.09-3.32 (m, 2H), 3.40 (d, J=9Hz, 1H), 3.55 (t, J=9Hz, 1H), 3.65 (d, J=9Hz, 1H), 3.81 (s, 3H), 5.96 (s, 2H), 6.75 (d, J=9Hz, 1H), 6.83 (d, J=9Hz, 1H), 6.88 (d, J=9Hz, 2H), 7.02 (s, 1H), 7.34 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 534 (M+H)⁺.

Example 316

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-[4-methoxyphenyl]sulfonylamino)propyl]-pyrrolidine-3-carboxylic acid

Ethyl *trans-trans* and *cis-trans* 2-(4-Methoxyphenyl)-4-(1,3-benzodiox-5-yl) -1-(3-bromopropyl) pyrrolidine-3-carboxylate

A 2:1 mixture of *trans-trans* and *cis-trans* ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodiox-5-yl) -pyrrolidine-3-carboxylate (4.00 g; prepared according to example 1C), 32 ml dibromopropane, and 200 mg sodium iodide, were heated at 100° for 1.25 hrs. The excess dibromopropane was removed in vacuo and the residue was dissolved in toluene. After shaking with potassium bicarbonate, the solution was dried (Na₂SO₄) and the solution concentrated. The residue was chromatographed on silica gel eluting with 5:1 hexane:EtOAc. yielding 5.22 (98%) of the title compound.

Example 316BEthyl *trans-trans* and *cis-trans* 2-(4-Methoxyphenyl)-4-(1,3-benzodiox-5-yl)-1-(3-propylaminopropyl) pyrrolidine-3-carboxylate

The compound described in Example 316A (5.22 g) was heated at 80° for 2 hrs. with 35 ml. ethanol, 2.5 g. propylamine and 35 mg. sodium iodide. The solvents were removed in vacuo. The residue was dissolved in toluene, shaken with potassium bicarbonate solution and dried (Na₂SO₄). The solution was concentrated in vacuum to give 4.96 g of the title compound as an orange oil. This was used in the next step without purification.

Example 316C*trans-trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-[4-methoxyphenyl]sulfonylamino)propyl]-pyrrolidine-3-carboxylic acid

Using the method described in example 66, the compound prepared in Example 316B was reacted with 4-methoxybenzenesulfonyl chloride in acetonitrile containing diisopropylethylamine. The resulting product was chromatographed on silica gel (30% EtOAc in hexane), and hydrolyzed to the title compound by the method of example 1D. NMR (CDCl₃, 300 MHz) δ 0.83 (t, J=7Hz, 3H), 1.40-1.52 (m, 2H), 1.56-1.70 (m, 2H), 2.00-2.11 (m, 1H), 2.40-2.51 (m, 1H), 2.69-2.78 (m, 1H), 2.84-3.03 (m, 4H), 3.19-3.34 (m, 2H), 3.48-3.59 (m, 2H), 3.80 (s, 3H), 3.86 (s, 3H), 5.95 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.85 (d, J=8Hz, 3H), 6.93 (d, J=8Hz, 2H), 7.02 (d, J=2Hz, 1H), 7.29 (d, J=8Hz, 2H), 7.69 (d, J=8Hz, 2H). Anal calcd for C₃₂H₃₈N₂O₈S : C, 62.93; H, 6.27; N, 4.59. Found: C, 62.97; H, 6.39; N, 4.45.

Example 317*trans-trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-propylsulfonylamino)propyl]-pyrrolidine-3-carboxylic acid

Using the method described in example 66, the propylamino compound prepared in Example 316B was reacted with propanesulfonyl chloride in acetonitrile containing diisopropylethylamine. The resulting product was chromatographed on silica gel (30% EtOAc in hexane) and hydrolyzed to the title compound by the method of example 1D. NMR (CDCl₃, 300 MHz) δ 0.85 (t, J=7Hz, 3H), 1.02 (t, J=7Hz, 3H), 1.47-1.60 (m, 2H), 1.65-1.85 (m, 4H), 2.04-2.16 (m, 1H), 2.42-2.57 (m, 1H), 2.72-3.11 (m, 5H), 3.25-3.41 (m, 2H), 3.50-3.62 (m, 2H), 3.80 (s, 3H), 5.85 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.80-6.90 (m, 3H), 7.02 (d, J=2Hz, 1H), 7.30 (d, J=9Hz, 2H). Anal calcd for C₂₈H₃₈N₂O₇S : C, 61.52; H, 7.01; N, 5.12. Found: C, 61.32; H, 7.01; N, 5.01.

Example 318

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 313 and Example 66, the title compound was prepared as a white solid. m.p. 66-68°C. ¹H NMR (CDCl₃, 300MHz) δ 0.81(t, J=7.5Hz, 3H), 0.89(t, J=7Hz, 3H), 1.26-1.35(m, 4H), 1.45(sextet, J=7.5Hz, 2H), 1.68-1.76(m, 2H), 2.25-2.33(m, 1H), 2.72-2.92(m, 5H), 2.97-3.12(m, 2H), 3.16-3.33(m, 2H), 3.43(dd, J=3Hz, J=9Hz, 1H), 3.53-3.60(m, 1H), 3.66(d, J=10Hz, 1H), 3.88(s, 3H), 5.95(s, 2H), 6.74(d, J=8Hz, 1H), 6.82(dd, J=1Hz, J=8Hz, 1H), 6.92(t, J=8Hz, 1H), 6.97(d, J=1Hz, 1H), 7.12(d, J=8Hz, 1H), 7.18(dd, J=1Hz, J=12Hz, 1H). MS (DCI/NH₃) m/e 579 (M+H)⁺.

Example 319

trans-trans-2-(4-Pyridinyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(n-butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared using the methods described in examples 1 and 43, using methyl 3-oxo-3-(4-pyridyl)propanoate (J. Am. Chem. Soc. **1993**, 115, 11705) in place of ethyl (4-methoxybenzoyl)acetate. m.p. 131-132 °C. NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.05-1.50 (m, 8H), 2.90 (dd, J= 7Hz, 9Hz, 1H), 2.97 (d, J=13Hz, 1H), 3.00-3.25 (m, 4H), 3.32 (m, 1H), 3.39 (d, J=13Hz, 1H), 3.45-3.52 (m, 1H), 3.67-3.78 (m, 1H), 4.10 (d, J=9Hz, 1H), 5.92 (dd, J=2Hz, 4 Hz, 2H), 6.75 (d, J=9Hz, 1H), 6.90 (dd, J=9Hz, 2Hz, 1H), 7.02 (d, J=2Hz, 1H), 7.45 (d, J=8Hz, 2H), 8.50 (d, J=8Hz, 2H). Anal calcd for C₂₇H₃₅N₃O₅ : C, 67.34; H, 7.33; N, 8.73 . Found: C, 67.39; H, 7.45; N, 8.61.

Example 320

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(N-propyl-N-diethylaminocarbonylamino)ethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in example 61, substituting propylamine for aqueous methylamine in Example 61B and diethylcarbamyl chloride for isobutyryl chloride in Example 61C. NMR (CD₃OD, 300 MHz) δ 0.74 (t, 3H, J=7), 1.09 (t, 6H, J=7), 1.33 (m, 2H), 3.17 (q, 4H, J=7), 3.78 (s, 3H), 4.04 (m, 1H), 5.93 (s, 2H), 6.86 (d, 1H, J=8), 7.06 (dd, 1H, J=2,8), 6.94 (d, 2H, J=9), 7.04 (d, 1H, J=2), 7.40 (d, 2H, J=9). MS (DCI/NH₃) m/z 526. Anal calcd for C₂₉H₃₉N₃O₆ . 0.1 TFA: C, 65.31; H, 7.34; N, 7.82 . Found: C, 65.33; H, 7.43; N, 8.14.

Example 321

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[3,5-dimethylpiperidinyl-carbonylmethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) shows mixture of isomers. δ 0.88 (d, 3H, J=7), 0.93 (d, 3H, J=7), 3.82 (s, 3H), 5.95 (s, 2H), 6.82 (d, 1H, J=8), 6.89 (dd, 1H, J=1,8), 7.00 d, 2H, J=9), 7.03 (m, 1H), 7.47 (d, 2H, J=9). MS (DCI/NH₃) m/z 495.

5

Example 322

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-di(s-butyl)aminocarbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) suggests a mixture of isomers. δ 0.83 (t, 6H, J=8), 1.27 (d, 6H, J=7), 1.6 (m, 2H), 3.79 (s, 3H), 5.93 (s, 2H), 6.75 (d, 1H, J=8), 6.86 (d, 1H, J=8), 6.94 (d, 2H, J=9), 7.03 (d, 1H, J=2), 7.35 (d, 2H, J=9). MS (DCI/NH₃) m/z 511.

10
15
20
25
30
35

Example 323

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N-(2-Methylphenyl)-N-butylamino carbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. MS (DCI/NH₃) m/z 545. Anal calcd for C₃₂H₃₆N₂O₆ . 0.9 H₂O: C, 68.53; H, 6.79; N, 4.99 . Found: C, 68.56; H, 6.62; N, 4.71.

20

Example 324

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N-(3-Methylphenyl)-N-butylamino carbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared using the procedures described in example 1. NMR (CD₃OD, 300 MHz) δ 0.88 (t, 3H, J=7), 1.2-1.5 (m, 4H), 2.31 (s, 3H), 2.8 (m, 2H), 3.14 (t, 1H, J=10), 3.3 (m, 1H), 3.44 (dd, 1H, J=5,10), 3.53 (m, 1H), 3.60 (t, 2H, J=7), 3.79 (s, 3H), 3.82 (m, 1H), 5.93 (s, 2H), 6.74 (d, 1H, J=8), 6.8-6.9 (m, 5H), 7.06 (d, 1H, J=2), 7.09 (d, 2H, J=9), 7.18 (d, 1H, J=7), 7.27 (t, 1H, J=7). MS (DCI/NH₃) m/z 545. Anal calcd for C₃₂H₃₆N₂O₆ . 0.8 H₂O: C, 68.75; H, 6.78; N, 5.01 . Found: C, 68.70; H, 6.67; N, 4.85.

30

Example 325

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(benzyloxymethyl)-1-((N,N-dibutylaminocarbonylmethyl)pyrrolidine-3-carboxylic acid

35

Example 325A

Ethyl trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(benzyloxymethyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The procedures of Example 1A-1D were followed, substituting ethyl 4-benzyloxy-3-oxobutyrate for 4-methoxybenzoylacetate in Example 1A, to afford the title compound as a colorless oil. TLC (30% EtOAc-hexane) Rf 0.18; ¹H NMR (CDCl₃, 300 MHz) δ 0.88 (t, J=7Hz, 6H), 1.17 (t, J=7Hz, 3H), 1.20-1.34 (br m, 4H), 1.40-1.56 (br m, 3H), 2.85 (t, J=8Hz, 1H), 2.98-3.30 (m, 5H), 3.39-3.60 (m, 3H), 3.64-3.75 (m, 2H), 3.92 (d, J=14Hz, 1H), 4.10 (two overlapping q, J=6.5Hz, 2H), 4.53 (s, 2H), 5.91 (m, 2H), 6.69 (d, J=9Hz, 1H), 6.77 (dd, J=1.5, 9Hz, 1H), 6.91 (d, J=1.5Hz, 1H); MS (DCI/NH₃) m/e 553 (M+H)⁺.

Example 325B

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(benzyloxymethyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound was prepared according to the procedure of Example 71C, as a colorless glass. TLC (5% MeOH-CH₂Cl₂) Rf 0.13; ¹H NMR (CDCl₃, 300 MHz) δ 0.86 (t, J=7Hz), and 0.90 (t, J=7Hz, 6H total), 1.15-1.52 (br m, 8H), 2.96-3.35 (br m, 5H), 3.50-3.75 (br m, 2H), 3.80 (dd, J=3, 13Hz, 1H), 3.88-4.40 (br m, 6H), 4.45 (AB, 2H), 5.90 (s, 2H), 6.70 (d, J=8Hz, 1H), 6.84 (dd, J=1.8Hz, 1H), 6.93 (d, J=1Hz, 1H), 7.28-7.39 (m, 5H); MS (DCI/NH₃) m/e 524 (M+H)⁺.

Example 326

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(hydroxymethyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Example 326A

Ethyltrans,trans-4-(1,3-Benzodioxol-5-yl)-2-(hydroxymethyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The resultant product from Example 325A (128 mg, 0.232 mmol) and 25 mg of 20% Pd(OH)₂ on charcoal in 7 mL EtOH was stirred under 1 atm hydrogen for 48 h. The mixture was filtered through a plug of celite, and the catalyst was washed with 2 x 10 mL EtOH, then the combined filtrate and washes were concentrated under reduced pressure to afford the crude product. Purification by flash chromatography (40%EtOAc-hexane) provided the title compound.

Example 326B

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(hydroxymethyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound was prepared according to the procedure of Example 71C.

Example 327

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(N-methylpropenamid-3-yl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

5

Example 327A

Ethyl trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(formyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

10 The title compound is made by selective oxidation (e.g. using the Swern oxidation with DMSO, oxalyl chloride, ethyldiisopropylamine or using the Dess-Martin periodinane) of the compound of Example 326A.

Example 327B

Ethyl trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(O-tert-butylpropenoat-3-yl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

15

The title compound is produced by condensing the compound of Example 327A with tert-butyl triphenylphosphoranylidine acetate in CH_2Cl_2 solution.

Example 327C

Ethyl trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(propenoic acid-3-yl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

20

The title compound is produced by reacting the compound of Example 327B with trifluoacetic acid in CH_2Cl_2 (1:1).

25

Example 327D

Ethyl trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(N-methylpropenamid-3-yl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

30 The title compound is produced by condensing the compound of Example 327C with methylamine hydrochloride in the presence of a carbodiimide (e.g. N-ethyl-N-(3-dimethylamino)propylcarbodiimide, DCC).

Example 327E

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(N-methylpropenamid-3-yl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

35

The title compound is produced by reacting the compound of Example 327D with lithium hydroxide according to the procedure of Example 71C.

Example 328

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(1-hydroxy-2-propen-3-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

5

Example 328A

Ethyl *trans,trans*--4-(1,3-Benzodioxol-5-yl)-2-(1-hydroxy-2-propen-3-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The title compound is produced by reacting the compound of Example 327C with borane methyl sulfide complex.

10

Example 328B

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(1-hydrox-2-propen-3-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound is produced by condensing the compound of Example 328A with lithium hydroxide according to the procedure of Example 71C.

15

Example 329

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(N-benzylaminomethyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

20

Example 329A

Ethyl *trans,trans*--4-(1,3-Benzodioxol-5-yl)-2-(N-benzylaminomethyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The title compound is produced by condensing the compound of Example 327A with benzylamine in the presence of sodium cyanoborohydride in ethanol.

25

Example 329B

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(N-benzylaminomethyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

30

The title compound is produced by reacting the compound of Example 329A with lithium hydroxide according to the procedure of Example 71C.

Example 330

trans,trans--4-(1,3-Benzodioxol-5-yl)-2-(N-acetyl-N-benzylaminomethyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

35

Example 330A

Ethyl *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(N-acetyl-N-benzylaminomethyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The title compound is produced by reacting the compound of Example 3294A with acetic anhydride in the presence of pyridine or triethylamine.

Example 330B

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(N-acetyl-N-benzylaminomethyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound is produced by reacting the compound of Example 330A with lithium hydroxide according to the procedure of Example 71C.

Example 331

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(ethynyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Example 331A

Ethyl *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(ethynyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The title compound is made by employing the procedure of Corey and Fuchs (Tetrahedron Lett. **1972**, 3769-72), using the compound of Example 327A.

Example 331B

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(ethynyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound is produced by reacting the compound of Example 331A with lithium hydroxide according to the procedure of Example 71C.

Example 332

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(1-pentynyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Example 332A

Ethyl *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(pentynyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylate

The title compound is made by palladium-catalyzed coupling of the compound of Example 206A and propyl iodide, employing the procedure of Taylor, et. al. (J. Org. Chem. **1989**, 54(15), 3618-24).

Example 332B

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(1-pentynyl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

The title compound is produced by reacting the compound of Example 332A with lithium hydroxide according to the procedure of Example 71C.

Example 333

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[2-(2,6-dioxopiperidinyl) ethyl]-pyrrolidine-3-carboxylic acid

The compound of example 61A is added to a solution of the sodium salt of glutarimide in dimethylformamide. After stirring 24 hours, water is added and the mixture is extracted with ether. The resultant glutarimide is hydrolyzed to the title compound by the method of example 1D.

Example 334

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-diphenylaminocarbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared according to the procedures described in Example 1. ¹H NMR (300 MHz, CD₃OD) δ 2.83 (dd, 1, J = 8.1, 9.7), 2.99 (d, 1, J = 15.4), 3.19 (t, 1, J = 9.5), 3.49 (d, 1, J = 15.3), 3.51 (dd, 1, J = 4.6, 9.5), 3.57 (m, 1), 3.79 (s, 3), 3.85 (d, 1, J = 9.5), 5.90 (s, 2), 6.71 (d, 1, J = 8.0), 6.84 (m, 3), 7.04 (d, 1, J = 1.6), 7.14-7.16 (m, 6), 7.19-7.34 (m, 6); MS (DCI/NH₃) m/z 551; Anal Calcd for C₃₃H₃₀N₂O₆·0.65H₂O·0.35C₂H₅OCOCH₃: C, 69.77, H, 5.77, N, 4.76. Found: C, 69.75, H, 5.55, N, 4.64.

Example 335

trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-[N,N-diisopropylaminocarbonylmethyl]-pyrrolidine-3-carboxylic acid.

The title compound was prepared according to the procedures described in Example 1. ¹H NMR (300 MHz, CD₃OD) δ 0.95 (d, 3, J = 6.5), 1.24 (d, 3, J = 6.4), 1.30 (d, 6, J = 6.8), 2.85 (d, 1, J = 12.5), 3.04 (dd, 1, J = 8.1, 9.8), 3.14 (t, 1, J = 9.7), 3.32-3.55 (m, 3), 3.63 (m, 1), 5.92 (s, 2), 6.75 (d, 1, J = 8.1), 6.85 (dd, 1, J = 1.7, 8.1), 6.93 (m, 2), 7.02 (d, 1, J = 1.7), 7.35 (m, 2). MS (DCI/NH₃) m/z 483. Anal Calcd for C₂₇H₃₄N₂O₆·0.65 EtOAc: C, 65.86, H, 7.32, N, 5.19. Found: C, 5.74, H, 7.26, N, 5.52.

Example 336

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-N-propyl-N-butanefonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 313 and Example 66, the title compound was prepared as a white solid. m.p. 65-66°C. ^1H NMR (CDCl_3 , 300MHz) δ 0.82(t, J=7.5Hz, 3H), 0.92(t, J=7.5Hz, 3H), 1.34-1.52(m, 4H), 1.72(quintet, J=7.5Hz, 2H), 2.25-2.35(m, 1H), 2.72-2.94(m, 5H), 2.97-3.12(m, 2H), 3.19-3.46(m, 2H), 3.44(d, J=9Hz, 1H), 3.53-3.60(m, 1H), 3.67(d, J=9Hz, 1H), 3.89(s, 3H), 5.95(s, 2H), 6.74(d, J=8Hz, 1H), 6.82(d, J=8Hz, 1H), 6.92(t, J=9Hz, 1H), 6.97(s, 1H), 7.12(d, J=9Hz, 1H), 7.18(d, J=12Hz, 1H). MS (DCI/NH_3) m/e 565 ($\text{M}+\text{H}$) $^+$.

Example 337

Using methods described in the above examples, the compounds disclosed in Table 1 can be prepared.




Table 1




Table 1 cont.




Table 1 cont.







Table 1 cont.

R


40.

R


41.

R


42.




43.



44.



45.



46.



47.



48.



49.



50.



51.



52.



53.



54.




55.




56.




57.



58.



59.



60.

Table 1 cont.




Table 1 cont.







Table 1 cont.




97.




98.




99.




100.




101.




102.




103.




104.




105.




106.




107.




108.




109.




110.




111.




112.




113.




114.



115.



116.



117.

Table 1 cont.

R

118.



121.




124.




127.



130.



133.

R

119.



122.



125.




128.




131.




134.

R


120.




123.



126.



129.




132.



135.

Table 1 cont.

R


136.

R


137.

R


138.




139.




140.




141.




142.




143.




144.




145.




146.




147.




148.



149.



150.



151.



152.



153.


Table 1 cont.

R

154.

R

155.

R


156.



157.



158.



159.



160.



161.



162.




163.




164.




165.



166.




167.




168.


Table 1 cont.




169.




170.




171.




172.




173.




174.




175.




176.




177.




178.




179.




180.




181.




182.




183.




184.




185.




186.



187.




188.




189.


Table 1 cont.

R


190.

R


191.

R


192.




193.




194.




195.




196.




197.




198.




199.




200.




201.




202.




203.




204.



205.



206.



207.

Table 1 cont.


















Table 1 cont.




Table 1 cont.


















Table 1 cont.


















Table 1 cont.





















Table 1 cont.

R


283.

R


284.

R


285.

R


286.

R


287.

R


288.

R


289.

R


290.

R


291.

R


292.

R


293.

R


294.

R

295.


R

296.


R

297.


Table 1 cont.

R


298.

R


299.

R


300.




301.




302.




303.




304.




305.




306.




307.




308.




309.



310.




311.




312.


Table 1 cont.

R


313.




316.




319.




322.




325.

R


314.




317.




320.




323.




326.

R


315.




318.



321.




324.



327.

Table 1 cont.

R


328.




331.



334.




337.



340.

R


329.




332.



335.




338.



341.

R

330.




333.



336.



339.



342.

Table I cont.


















Table 1 cont.


















Table 1 cont.


















Table 1 cont.


















Table 1 cont.







Table 1 cont.

R


418.

R


419.

R

420.



421.



422.



423.



424.



425.



426.



427.




428.




429.



430.




431.




432.


Table 1 cont.

R


433.

R


434.

R


435.




436.




437.




438.




439.




440.




441.




442.




443.




444.



445.




446.




447.


Table 1 cont.

R


448.




451.




454.




457.




460.

R


449.




452.




455.




458.




461.

R


450.




453.



456.



459.



462.

Table 1 cont.




Table I cont.


















R**R****R**


Table 1 cont.

R


493.




496.




499.




502.




505.

R


494.




497.




500.




503.




506.

R


495.




498.



501.



504.



507.

Table 1 cont.


















Table 1 cont.


















R**R****R**


Table 1 cont.

R


538.




540.




544.




547.




550.

R


539.




542.




545.




548.




551.

R


540.




543.



546.



549.



552.

Table I cont.


















R**R****R**


Table 1 cont.




568.




569.




570.




571.




572.




573.




574.




575.




576.




577.




578.




579.




580.




581.




582.



583.




584.




585.


Table 1 cont.

R


586.




589.




592.




595.




598.




601.

R


587.




590.




593.




596.




599.




602.

R


588.




591.




594.



597.




600.




603.


Table 1 cont.

R


604.




607.




610.




613.




616.




619.

R


605.




608.




611.




614.




617.




620.

R


606.




609.




612.



615.




618.




621.


Table 1 cont.




622.




623.




624.




625.




626.




627.




628.




629.




630.




631.




632.




633.




634.




635.




636.



637.




638.




639.


Table 1 cont.

R


640.

R


641.

R


642.




643.




644.




645.




646.




647.




648.




649.




650.




651.



652.




653.




654.


Table 1 cont.

R


655.

R


656.

R


657.




658.




659.




660.




661.




662.




663.




664.




665.




666.



667.



668.



669.

Table 1 cont.




Table 1 cont.


































Table 1 cont.



Example 338

Using methods described in the above examples, compounds comprising a parent structure selected from those disclosed in Table 2A and an R substituent selected from those disclosed in Table 2B can be prepared.

5

Table 2A




Table 2A cont,















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.


















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.















Table 2A cont.


















Table 2A cont.




193.




194.




195.




196.




197.




198.




199.




200.




201.



202.




203.




204.


Table 2A cont.




205.




206.




207.




208.




209.




210.




211.




212.




213.



214.



215.



216.

Table 2A cont.















Table 2A cont.

















Table 2A cont.




Table 2A cont.




253.




254.




255.




256.




257.




258.




259.




260.




261.




262.




263.




264.



265.




266.




267.


Table 2A cont.




268.




269.




270.




271.




272.




273.




274.




275.




276.



277.



278.



279.

Table 2A cont.


















Table 2A cont.















Table 2A cont.


















Table 2A cont.





















Table 2A cont.




337.




338.




339.




340.




341.




342.




343.




344.




345.




346.




347.




348.



349.




350.




351.


Table 2A cont.




352.




353.




354.




355.




356.




357.




358.




359.




360.



361.



362.



363.

Table 2A cont.


















Table 2A cont.




376.




377.




378.




379.




380.




381.




382.




383.




384.



385.




386.




387.


Table 2A cont.




388.




389.




390.




391.




392.




393.




394.




395.




396.



397.



398.



399.

Table 2A cont.





















Table 2A cont.




430.




431.




432.




433.




434.




435.




436.




437.




438.




439.




440.




441.



442.




443.




444.


Table 2A cont.




445.




446.




447.




448.




449.




450.




451.




452.




453.



454.




455.




456.


Table 2A cont.




457.




458.




459.




460.




461.




462.



463.




464.




465.

Table 2B

R


1.




4.




7.




10.



13.




16.



19.

R


2.




5.




8.




11.



14.



17.




20.

R


3.




6.




9.




12.



15.



18.



21.

Table 2B cont.

























Table 2B cont.

R


43.

R


44.

R


45.



46.




47.




48.



49.



50.




51.



52.



53.




54.



55.



56.




57.



58.



59.



60.

Table 2B cont.



















Table 2B cont.




Table 2B cont.





















Table 2B cont.





















Table 2B cont.

R


133.

R


134.

R


135.




136.




137.




138.




139.




140.




141.




142.




143.




144.



145.



146.



147.

Table 2B cont.


















R**R****R**


Table 2B cont.

R


163.

R


164.

R


165.




166.




167.




168.




169.




170.




171.




172.




173.




174.



175.




176.




177.


Table 2B cont.

R


178.

R


179.

R


180.




181.




182.




183.




184.




185.




186.




187.




188.




189.



190.




191.




192.


Table 2B cont.

R


193.




196.




199.




202.




205.

R


194.




197.




200.




203.




206.

R


195.




198.



201.




204.




207.


Table 2B cont.

R


208.




211.




214.




217.




220.

R


209.




212.




215.




218.




221.

R


210.




213.



216.




219.




222.


Table 2B cont.




223.




224.




225.




226.




227.




228.




229.




230.




231.




232.




233.




234.



235.




236.




237.


Table 2B cont.

R


238.

R


239.

R


240.




241.



242.



243.



244.



245.




246.



247.




248.




249.



250.




252.



252.


Table 2B cont.

R


253.

R


254.

R


255.




256.




257.




258.




259.




260.




261.




262.




263.




264.



265.




266.




267.


Table 2B cont.

R


268.

R


269.

R


270.




271.




272.




273.




274.




275.




276.




277.




278.




279.



280.



281.



282.

Table 2B cont.



283.



284.



285.



Table 2B cont.





















Table 2B cont.

R


313.



316.




319.




322.




325.

R

314.



317.




320.




323.




326.

R

315.



318.



321.



324.



327.

Table 2B cont.



















Table 2B cont.

R


343.




346.



349.




352.




355.

R


344.




347.



350.




353.




356.

R


345.




348.



351.




354.




357.


Table 2B cont.

R


358.




361.




364.




367.




370.

R


359.




362.




365.




368.




371.

R


360.




363.



366.



369.



372.

Table 2B cont.
























Table 2B cont.

R


391.

R


392.

R


393.




394.




395.




396.




397.




398.




399.




400.




401.




402.




403.




404.




405.



406.



407.



408.

Table 2B cont.







Table 2B cont.




427.




428.




429.




430.




431.




432.




433.




434.




435.




436.




437.




438.




439.




440.




441.



442.




443.




444.


Table 2B cont.

R


445.




448.




451.




454.




457.

R


446.




449.




452.




455.




458.

R


447.




450.



453.



456.



459.

Table 2B cont.





















Table 2B cont.

R


475.




478.




481.




484.




487.

R


476.




479.




482.




485.




488.

R


477.




480.



483.




486.




489.


Table 2B cont.

R


490.




493.




496.




499.




502.




505.

R


491.




494.




497.




500.




503.




506.

R


492.




495.




498.



501.




504.




507.


Table 2B cont.

R


508.




511.




514.




517.




520.

R


509.




512.




515,




518.




521.

R


510.




513.



516.




519.




522.


Table 2B cont.

R


523.




526.




529.




532.




535.

R


524.




527.




530.




533.




536.

R


525.




528.



531.




534.




537.


Table 2B cont.

R


538.

R


539.

R


540.




541.




542.




543.




544.




545.




546.




547.




548.




549.



550.




551.




552.


Table 2B cont.

R


553.




556.




559.




562.




565.

R


554.




557.




560.




563.




566.

R


555.




558.



561.




564.




567.


Table 2B cont.

R


568.




571.




574.




577.




580.

R


569.




572.




575.




578.




581.

R


570.




573.



576.




579.




582.


Table 2B cont.

R


583.




586.




589.




592.




595.

R


584.




587.




590.




593.




596.

R


585.




588.



591.




594.




597.


Table 2B cont.

R


598.




601.




604.




607.




610.

R


599.




602.




605.




608.




611.

R


600.




603.



606.



609.



612.

Table 2B cont.





















Table 2B cont.

R


628.

R


629.

R

630.



631.



632.

5

10

Example 339

Using methods described in the above examples, compounds comprising a parent structure selected from those disclosed in Table 3A and an R substituent selected from those disclosed in Table 3B can be prepared.

15

Table 3A















Table 3A cont.












Table 3B







Table 3B cont.

R


22.




25.




28.




31.




34.




37.




40.

R


23.




26.




29.



32.




35.




38.




41.

R


24.




27.




30.



33.



36.




39.




42.


Table 3B cont.




43.




44.




45.



46.




47.




48.



49.



50.




51.



52.



53.




54.



55.



56.




57.



58.



59.



60.

Table 3B cont.



















Table 3B cont.




Table 3B cont.





















Table 3B cont.





















Table 3B cont.




133.




134.




135.




136.




137.




138.




139.




140.




141.




142.




143.




144.



145.




146.




147.


Table 3B cont.

R


148.




151.




154.




157.




160.

R


149.




152.




155.




158.




161.

R


150.




153.



156.



159.



162.

Table 3B cont.





















Table 3B cont.

R


178.

R


179.

R


180.




181.




182.




183.




184.




185.




186.




187.




188.




189.



190.




191.




192.


Table 3B cont.

R


193.




196.




199.




202.




205.

R


194.




197.




200.




203.




206.

R


195.




198.



201.




204.




207.


Table 3B cont.

R


208.

R


209.

R


210.




211.




212.




213.




214.




215.




216.




217.




218.




219.



220.




221.




222.


Table 3B cont.




223.




224.




225.




226.




227.




228.




229.




230.




231.




232.




233.




234.



235.



236.



237.

Table 3B cont.




















Table 3B cont.

R


253.

R


254.

R


255.




256.




257.




258.




259.




260.




261.




262.




263.




264.



265.




266.




267.


Table 3B cont.

R


268.




271.




274.




277.




280.

R


269.




272.




275.




278.




281.

R


270.




273.



276.




279.




282.


Table 3B cont.




283.




284.




285.




286.




287.




288.




289.




290.




291.




292.




293.




294.



295.



296.



297.

Table 3B cont.





Table 3B cont.

R


343.




346.



349.




352.




355.

R


344.




347.



350.




353.




356.

R


345.




348.



351.



354.



357.

Table 3B cont.


















R**R****R**


Table 3B cont.

R


373.

R


374.

R


375.




376.




377.




378.




379.




380.




381.




382.




383.




384.




385.




386.




387.



388.



389.



390.

Table 3B cont.



















R**R****R**

Table 3B cont.







Table 3B cont.




427.




428.




429.




430.




431.




432.




433.




434.




435.




436.




437.




438.




439.




440.




441.



442.




443.




444.


Table 3B cont.

R


445.




448.




451.




454.




457.

R


446.




449.




452.




455.




458.

R


447.




450.



453.



456.



459.

Table 3B cont.





















Table 3B cont.

R


475.

R


476.

R


477.




478.




479.




480.




481.




482.




483.




484.




485.




486.



487.




488.




489.


Table 3B cont.

R


490.




493.




496.




499.




502.




505.

R


491.




494.




497.




500.




503.




506.

R


492.




495.




498.



501.




504.




507.


Table 3B cont.

R


508.

R


509.

R


510.




511.




512.




513.




514.




515.




516.




517.




518.




519.



520.




521.




522.


Table 3B cont.

R


523.




526.




529.




532.




535.

R


524.




527.




530.




533.




536.

R


525.




528.



531.




534.




537.


Table 3B cont.

R


538.




541.




544.




547.




550.

R


539.




542.




545.




548.




551.

R


540.




543.



546.




549.




552.


Table 3B cont.

R


553.




556.




559.




562.




565.

R


554.




557.




560.




563.




566.

R


555.




558.



561.



564.



567.

Table 3B cont.





















Table 3B cont.

R


583.

R


584.

R


585.




586.




587.




588.




589.




590.




591.




592.




593.




594.



595.



596.



597.

Table 3B cont.




Table 3B cont.




















Table 3B cont.

R




628.

R




629.


R



630.



631.



632.

Example 340*trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-(3-methylbut-1-yl)-N-phenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.85 (d, J=6 Hz, 6H), 1.25 (q, J=7 Hz, 2H), 1.42-1.56 (m, 1H), 3.43-3.85 (m, 9H), 3.88s (3), 5.95 (s, 2H), 6.80 (d, J=7 Hz, 1H), 6.86 (dd, J=9 Hz, 1H), 6.89-7.00 (m, 2H), 6.97 (d, J=1 Hz, 1H), 7.04 (d, J=9 Hz, 2H), 7.37 (d, J=9 Hz, 2H), 7.40-7.47 (m, 3H). MS (C.I.) m/e C (53.12, 53.11), H (4.63, 4.80), N (3.33, 3.28).

Example 341*trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(4-methylphenylaminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.20-1.47 (m, 4H), 2.37 (s, 3H), 2.83 (q, J=7 Hz, 2H), 3.06-3.25 (m, 2H), 3.40-3.50 (m, 1H), 3.51-3.63 (m, 3H), 3.80 (s, 3H), 3.87 (d, J=9 Hz, 1H), 5.92 (s, 2H), 6.74 (d, J=8 Hz, 1H), 6.80-6.86 (m, 3H), 6.89 (d, J=8 Hz, 2H), 7.04 (d, J=2 Hz, 1H), 7.12 (d, J=8 Hz, 2H), 7.19

(d, J=8 Hz, 2H). MS (DCI) m/e 545 (M+H)⁺. Analysis calcd for C₃₂H₃₆N₂O₆: C, 70.57; H, 6.66; N, 5.14. Found: C, 70.20; H, 6.81; N, 5.03.

Example 342

5 trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-propoxyphenyl)-1-(N,N-di(n-butyl)amino)carbonylmethyl-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H (300MHz, CDCl₃) δ 7.30 (2H, d, J=9), 7.03 (1H, d, J=2), 6.83 (3H, m), 6.72 (1H, d, J=9), 5.95 (1H, d, J=2), 5.93 (1H, d, J=2), 3.88 (2H, t, J=7), 3.73 (1H, d, J=12), 3.58 (1H, m), 3.53-3.20 (4H, m), 3.10-2.90 (4H, m), 2.72 (1H, d, J=15), 1.79 (2H, q, J=8), 1.50-1.05 (8H, m), 1.02 (3H, t, J=7), 0.87 (3H, t, J=7), 0.80 (3H, t, J=7). MS (DCI/NH₃) m/e 539 (M+H)⁺. Anal calcd for C₃₁H₄₂N₂O₆ · 0.5H₂O: C, 67.98; H, 7.91; N, 5.11. Found: C, 68.24; H, 7.70; N, 5.03.

15 Example 343

20 trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-propylphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H (300MHz, CDCl₃) δ 7.31 (2H, d, J=9), 7.13 (2H, d, J=9), 7.03 (1H, d, J=2), 6.84 (1H, dd, J=6, 2), 6.73 (1H, d, J=9), 5.95 (1H, d, J=2), 5.93 (1H, d, J=2), 3.76 (1H, d, J=10), 3.60 (1H, m), 3.55-3.20 (4H, m), 3.13-2.88 (4H, m), 2.75 (1H, d, J=15), 2.55 (2H, t, J=8), 1.62 (2H, q, J=8), 1.50-1.00 (8H, m), 0.92 (3H, t, J=7), 0.85 (3H, t, J=7), 0.78 (3H, t, J=7). MS (DCI/NH₃) m/e 523 (MH)⁺. Anal calcd for C₃₁H₄₂N₂O₅·0.25 H₂O: C, 70.63; H, 8.13; N, 5.31. Found: C, 70.55; H, 8.08; N, 5.18.

Example 344

30 trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-(N-propyl-N-n-pentanesulfonylamino)propyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 316, the title compound was prepared. ¹H NMR (300MHz, CDCl₃) δ 0.85 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.3-1.4 (m, 4H), 1.5-1.6 (sextet, J=7, 2H), 1.65-1.8 (m, 4H), 2.05-2.15 (m, 1H), 2.43-2.56 (m, 1H), 2.72-3.1 (m, 7H), 3.27-3.4 (m, 2H), 3.5-3.6 (m, 2H), 3.80 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.8-6.9 (m, 1H), 6.85 (d, J=9Hz, 2H), 7.02 (d, J=2Hz, 1H), 7.80 (d, J=9Hz, 2H).

Example 345

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-ethylphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H (300MHz, CDCl₃) δ 7.40 (3H, m), 7.22 (2H, d, J=8), 7.13 (1H, dd, J=8, 3), 6.72 (1H, d, J=9), 5.28 (1H, d, J=12), 4.55 (2H, t, J=9), 4.15 (1H, d, J=18), 4.03 (2H, m), 3.75 (2H, m), 3.40 (2H, m), 3.20 (2H, t, J=9), 3.15 (1H, m), 3.10-2.90 (2H, m), 2.63 (2H, q, J=9), 1.47 (2H, m), 1.31 (4H, m), 1.12 (3H, t, J=8), 1.10 (2H, m), 0.92 (3H, t, J=9), 0.80 (3H, t, J=9). MS (DCI/NH₃) m/e 507 (M+H⁺). Anal calcd for C₃₁H₄₂N₂O₄ · 1.0 TFA: C, 63.86; H, 6.98; N, 4.51. Found: C, 63.95; H, 7.12; N, 4.43.

Example 346

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-(3-pentyl)-N-phenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.93 (t, J=7.3 Hz, 3H), 0.94 (t, J=7.3 Hz, 3H), 1.33 (m, 4H), 2.72 (d, J=15.2 Hz, 1H), 2.81 (m, 1H), 3.11-3.23 (m, 2H), 3.45-3.57 (m, 2H), 3.79 (s, 3H), 3.83 (d, J=9.8 Hz, 1H), 4.54 (m, 1H), 5.92 (s, 2H), 6.73 (d, J=7.8 Hz, 1H), 6.83 (m, 3H), 6.98 (bs, 2H), 7.04 (d, J=1.7 Hz, 1H), 7.07 (2), 7.37 (m, 3H). MS (DCI) m/e 545 (M+H⁺). Anal calcd for C₃₂H₃₃N₂O₆ · 0.35H₂O: C, 69.76; H, 6.71; N, 5.08. Found: C, 69.72; H, 6.66; N, 4.94.

Example 347

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl)-N-(3-trifluoromethylphenyl)amino)carbonyl)methylpyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=6.6 Hz, 3H), 1.17-1.45 (m, 4H), 2.65 (d, J=16.5 Hz, 1H), 2.72 (m, 1H), 3.10 (t, J=9.5 Hz, 1H), 3.21-3.27 (m, 1H), 3.40 (dd, J=4.1, 9.9 Hz, 1H), 3.54 (m, 1H), 3.61-3.74 (m, 3H), 3.77 (s, 3H), 5.93 (s, 2H), 6.73-6.85 (m, 4H), 7.02 (m, 3H), 7.33 (d, J=7.5 Hz, 1H), 7.40 (s, 1H), 7.58 (t, J=7.8 Hz, 1H), 7.69 (d, J=7.5 Hz, 1H). MS (DCI) m/e 599 (M+H⁺). Anal calcd for C₃₂H₃₃F₃N₂O₆: C, 64.21; H, 5.56; N, 4.68. Found: C, 64.09; H, 5.63; N, 4.57.

Example 348

trans, trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-propyl-N-(4-morpholinylcarbonyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.78 (t, J=7 Hz, 3H), 1.43 (q, J=7 Hz, 2H), 2.07-3.01 (m, 1H), 2.76 (dd, J=7, 9 Hz, 2H), 2.77-3.00 (m, 5H), 3.05 (3.70, J=m Hz, 11H), 3.76 (s, 3H), 5.88 (s, 2H), 6.67 (d, J=8 Hz, 1H), 6.80 (dd, J=7 Hz, 1H), 6.83-6.90 (m, 2H), 6.98 (d, J=2 Hz, 1H), 7.32-7.39 (m, 2H). MS m/e calc'd for (M+H) C₂₉H₃₉N₃O₇: (M+H) 540.2710. Found (M+H) 540.2713.

10

Example 349

trans, trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(cis-2,6-dimethylpiperidin-1-yl)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.94 (d, J=7 Hz, 3H), 1.15d (7, 3H), 1.10-1.70 (m, 6H), 1.70-1.90 (m, 1H), 2.9. (d, J=13 Hz, 1H), 3.00-3.20 (m, 2H), 3.50 (3.70, J=m Hz, 2H), 3.79 (s, 3H), 3.80-4.00 (m, 1H), 4.10-4.65 (m, 2H), 5.95 (s, 2H), 6.70 (7.10, J=m Hz, 5H), 7.35 (m, 2H). MS m/e calc'd for (M+H)⁺ C₂₈H₃₅N₂O₆: (M+H) 495.2495. Found (M+H) 495.2493.

20

Example 350

trans, trans-2-(4-Methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 57-59 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.78 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.28-1.36 (m, 4H), 1.93 (sextet, J=7Hz, 2H), 1.72 (t, J=7Hz, 2H), 2.20-2.32 (m, 1H), 2.72-3.10 (m, 7H), 3.18-3.41 (m, 2H), 3.43 (dd, J=3Hz, J=9Hz, 1H), 3.48 (s, 3H), 3.52-3.59 (m, 1H), 3.68 (d, J=9Hz, 1H), 5.15 (s, 2H), 5.94 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.82 (dd, J=1Hz, J=8Hz, 1H), 6.98-7.02 (m, 3H), 7.32 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 591 (M+H)⁺.

30

Example 351

trans, trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-(2-butyl)-N-phenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.79-0.89 (m, 6H), 1.14-1.21 (m, 1H), 1.25-1.40 (m, 1H), 2.64 (dd, J=4.6, 15.4 Hz, 1H), 2.76 (t, J=9.0 Hz, 1H), 3.05-3.13

35

(m, 2H), 3.37-3.49 (m, 2H), 3.70 (s, 3H), 3.80 (d, J=9.8 Hz, 1H), 4.53 (m, 1H), 5.83 (m, 2H), 6.65 (d, J=8.1 Hz, 1H), 6.72 (-6.76, J=m Hz, 3H), 6.87 (m, 2H), 6.95 (d, J=1.7 Hz, 1H), 7.03 (m, 2H), 7.29 (m, 3H). MS (DCI) m/e 531 (M+H⁺). Anal calcd for C₃₁H₃₄N₂O₆ · 0.4H₂O: C, 69.23; H, 6.52; N, 5.21. Found: C, 69.19; H, 6.52; N, 5.03.

Example 352

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-(2-propyl)-N-phenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.99 (d, J=6.8 Hz, 6H), 2.71 (d, J=15.6 Hz, 1H), 2.84 (m, 1H), 3.13-3.18 (m, 2H), 3.45-3.58 (m, 2H), 3.79 (s, 3H), 3.88 (d, J=9.8 Hz, 1H), 4.80 (m, 1H), 5.92 (s, 2H), 6.74 (d, J=8.1 Hz, 1H), 6.83 (m, 3H), 6.96 (br s, 2H), 7.04 (d, J=1.7 Hz, 1H), 7.13 (m, 2H), 7.38 (m, 3H). MS (DCI) m/e 517 (M+H⁺). Anal calcd for C₃₀H₃₂N₂O₆ · 0.4H₂O · 0.08CH₃CO₂C₂H₅: C, 68.65; H, 6.28; N, 5.28. Found: C, 68.64; H, 6.35; N, 5.14.

Example 353

trans,trans-4-(4-Propoxyphenyl)-2-(4-methoxyphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H (300MHz, CDCl₃) δ 7.42 (2H, d, J=10Hz), 7.38 (2H, d, J=10Hz), 6.92 (2H, d, J=10Hz), 6.88 (2H, d, J=10Hz), 5.13 (1H, bd, J=12Hz), 4.02 (2H, m), 3.90 (2H, t, J=8Hz), 3.80 (3H, s), 3.71 (3H, m), 3.40 (2H, m), 3.19 (1H, m), 3.10-2.90 (2H, m), 1.80 (2H, m), 1.48 (2H, m), 1.29 (4H, m), 1.13 (2H, m), 1.03 (3H, t, J=8Hz), 0.92 (3H, t, J=9Hz), 0.82 (3H, t, J=9Hz). MS (DCI/NH₃) m/e 525 (MH⁺). Anal calcd for C₃₁H₄₄N₂O₅ · 1 TFA : C, 62.06 H 7.10; N, 4.39 . Found: C, 62.43; H, 7.28; N, 4.39.

Example 354

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((1,2,3,4-tetrahydroquinolin-1-yl)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 1.88 (quintet, J=6.5 Hz, 2H), 2.67 (t, J=6.4 Hz, 2H), 2.87 (t, J=8.6 Hz, 1H), 3.14 (m, 2H), 3.42 (dd, J=4.6, 9.7 Hz, 1H), 3.53-3.70 (m, 3H), 3.72-3.78 (m, 1H), 3.77 (s, 3H), 3.86 (d, J=9.6 Hz, 1H), 5.91 (s,

2H), 6.73 (d, J=8.1 Hz, 1H), 6.83 (m, 3H), 6.98 (d, J=1.1 Hz, 1H), 7.02-7.23 (m, 6H). MS (DCI) m/e 515 (M+H⁺). Anal calcd for C₃₀H₃₀N₂O₆ · 0.3H₂O · 0.15 CH₃CO₂C₂H₅: C, 68.93; H, 6.01; N, 5.25. Found: C, 68.91; H, 5.86; N, 5.19.

5

Example 355

trans,trans-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 64-65 °C. ¹H NMR (CDCl₃,

10 300MHz) δ 0.79 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.07 (sextet, J=7Hz, 2H), 1.20-1.35 (m, 4H), 1.43 (sextet, J=7Hz, 2H), 2.83 (d, J=13.5Hz, 1H), 2.94-3.17 (m, 4H), 3.22-3.42 (m, 1H), 3.40-3.48 (m, 3H), 3.58-3.65 (m, 1H), 3.82 (s, 3H), 3.85 (s, 4H), 5.92 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.81 (d, J=8Hz, 1H), 6.86-6.96 (m, 3H), 7.07 (d, J=3Hz, 1H). MS (DCI/NH₃) m/e 541 (M+H⁺).

15

Example 356

trans,trans-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-*n*-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 75-86 °C. ¹H NMR (CD₃OD, 300

20 MHz) δ 0.75 (t, J=7Hz, 3H), 0.82 (t, J=7Hz, 3H), 1.32-1.43 (m, 6H), 1.65-1.77 (m, 2H), 3.0-3.09 (m, 4H), 3.23-3.27 (m, 2H), 3.44 (t, J=6Hz, 1H), 3.47-3.56 (m, 2H), 3.78 (d, J=9Hz, 1H), 3.83-3.93 (m, 2H), 3.87 (s, 3H), 3.92 (s, 3H), 4.63 (d, J=13Hz, 1H), 5.97 (s, 2H), 6.82 (d, J=7Hz, 1H), 6.93 (d, J=7Hz, 1H), 7.06 (d, J=7Hz, 1H),
25 7.08 (d, J=3Hz, 1H), 7.16 (dd, J=3Hz, J=7Hz, 1H), 7.27 (d, J=3Hz, 1H). MS (DCI/NH₃) m/e 591 (M+H⁺).

Example 357

trans,trans-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-*n*-hexanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 65-66 °C. ¹H NMR (CDCl₃, 300

30 MHz) δ 0.80 (t, J=7Hz, 3H), 0.89 (t, J=7Hz, 3H), 1.23-1.48 (m, 6H), 1.43 (sextet, J=7Hz, 2H), 1.72 (sextet, J=7Hz, 2H), 2.25-2.35 (m, 1H), 2.73-3.10 (m, 7H), 3.19-3.32 (m, 2H), 3.45 (dd, J=3Hz, J=9Hz, 1H), 3.53-3.59 (m, 1H), 3.68 (d, J=9Hz,

35

1H), 3.87 (s, 6H), 5.95 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.79-6.86 (m, 2H), 6.92-6.97 (m, 2H), 7.02 (s, 1H). MS (DCI/NH₃) m/e 605 (M+H)⁺.

Example 358

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(phthalimido)ethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 1C (250 mg), N-bromoethylphthalimide (206 mg), and diisopropylethylamine (175 mg) were dissolved in 1 mL of acetonitrile and heated for 2.5 hours at 95 °C. Toluene was added, and the mixture was washed with KHCO₃ solution. The solution was dried (Na₂SO₄) and concentrated. The crude product was purified by chromatography on silica gel eluting with 3:1 EtOAc-hexane to give 216 mg of an intermediate ethyl ester which was hydrolyzed by the method of Example 1D to give 130 mg of the title compound as a white powder. ¹H NMR (300 MHz, CDCl₃) δ 3.12-3.26 (m, 2H), 3.60-3.75 (m, 2H), 3.70 (s, 3H), 3.98-4.12 (m, 2H), 4.45-4.55 (m, 1H), 4.69 (d, J=9Hz, 1H), 4.76-4.88 (m, 1H), 5.96 (s, 2H), 6.55 (d, J=8Hz, 1H), 6.60-6.70 (m, 3H), 6.79 (d, J=8Hz, 1H), 7.05-7.45 (m, 5H), 7.75 (d, J=7Hz, 1H).

Example 359

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-(2-pentyl)-N-phenylamino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.86-0.98 (m, 6H), 1.17-1.22 (m, 1H), 1.23-1.41 (m, 3H), 2.70 (dd, J=11.2, 15.3 Hz, 1H), 2.83 (m, 1H), 3.10-3.21 (m, 2H), 3.45-3.60 (m, 2H), 3.79 (s, 3H), 3.86 (m, 1H), 4.74 (m, 1H), 5.91 (m, 2H), 6.73 (dd, J=1.1, 7.7 Hz, 3H), 6.82 (m, 2H), 7.04-7.14 (m, 3H), 7.36 (m, 3H). MS (DCI) m/e 545 (M+H⁺). Anal calcd for C₃₂H₃₆N₂O₆ · 0.25 CH₃CO₂C₂H₅: C, 69.95; H, 6.76; N, 4.94. Found: C, 70.03; H, 6.54; N, 4.78.

Example 360

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(2-naphthyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.83 (t, J=7 Hz, 3H), 1.23-1.39 (m, 4H), 1.40-1.55 (m, 3H), 2.60-2.72 (m, 2H), 3.00-3.80 (m, 5H), 3.66 (s, 3H), 5.87 (s, 2H), 6.39 (d, J=9 Hz, 2H), 6.74-6.85 (m, 3H), 7.17 (d, J=2 Hz, 1H), 7.40 (dd, J=8 Hz, 1H), 7.52-7.62 (m, 3H), 7.80-7.90 (m, 1H), 7.90-8.00 (m, 2H). MS (DCI) m/e 581 (M+H⁺). Analysis calcd for C₃₅H₃₆N₂O₆ · 0.3 H₂O: C, 71.73; H, 6.29; N, 4.78. Found: C, 71.74; H, 6.26; N, 4.72.

Example 361

trans,trans-2-(4-Propoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 53-54 °C. ¹H NMR (CDCl₃, 300MHz) δ 0.79 (t, J=7Hz, 3H), 0.89 (t, J=7Hz, 3H), 1.03 (t, J=7Hz, 3H), 1.24-1.34 (m, 4H), 1.43 (sextet, J=7Hz, 2H), 1.67-1.75 (m, 2H), 1.80 (sextet, 2H), 2.23-2.33 (m, 1H), 2.72-2.93 (m, 5H), 3.05 (septet, J=7Hz, 2H), 3.15-3.35 (m, 2H), 3.42 (d, J=9Hz, 1H), 3.54-3.62 (m, 1H), 3.67 (d, J=9Hz, 1H), 4.90 (t, J=7Hz, 2H), 5.95 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.85 (d, J=8Hz, 2H), 7.02 (s, 1H), 7.32 (d, J=8Hz, 2H). MS (DCI/NH₃) m/e 589 (M+H)⁺.

Example 362

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((2-methylindolin-1-yl)carbonyl)methylpyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ mixture of indole C₂ diastereomers, 0.95 (m, 1.5 (CH₃)), 1.05 (d, 6.3H, 1.5 (CH₃)), 2.62 (m, 1H), 3.01 (m, 2H), 3.14-3.25 (m, 1H), 3.37-3.52 (m, 1.5H), 3.56-3.80 (m, 2H), 3.65 (s, 1.5 (CH₃O)), 3.76 (s, 1.5 (CH₃O)), 3.93 (m, 0.5H), 4.05-4.13 (m, 0.5H), 4.42 (m, 0.5H), 4.65-4.74 (m, 1H), 5.91 (m, 2H), 6.72 (d, J=8.1 Hz, 0.5H), 6.75 (m, 0.5H), 6.85 (m, 2H), 6.92 (d, J=8.5 Hz, 1H), 7.00-7.06 (m, 2H), 7.14 (t, J=7.7 Hz, 1H), 7.21 (t, J=6.6 Hz, 1H), 7.38 (m, 2H), 7.99 (m, 1H). MS (DCI) m/e 515 (M+H)⁺. Anal calcd for C₃₀H₃₀N₂O₆ · 0.35H₂O · 0.3 CH₃CO₂C₂H₅: C, 68.47; H, 6.10; N, 5.12. Found: C, 68.46; H, 5.97; N, 5.07.

Example 363

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(2-hydroxy-3-propylthex-1-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 1.06 (m, 6H), 1.26-1.60 (m, 9H), 3.16 (dd, J=10.9, 12.6 Hz, 1H), 3.18 (d, J=11 Hz, 1H), 3.44 (d, J=2.0 Hz, 1H), 3.61 (t, J=11 Hz, 1H), 3.73 (t, J=11.0 Hz, 1H), 3.85 (m, 1H), 3.96-4.17 (m, 2H), 4.02 (s, 1.5 (CH₃O diastereomer)), 4.03 (s, 1.5 (CH₃O diastereomer)), 6.15 (s, 2H), 7.01 (d, J=8.1 Hz, 0.5H), 7.00 (d, J=8.1 Hz, 0.5H), 7.10 (m, 1H), 7.23 (m, 3H), 7.77 (m, 2H).

MS (DCI.) m/e 484 (M+H⁺). Anal calcd for C₂₈H₃₇NO₆ · 0.33 H₃PO₄: C, 65.34; H, 7.44; N, 2.72. Found: C, 65.30; H, 7.40; N, 2.60.

Example 364

5 *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-(4-heptyl)-N-(3,4-dimethoxybenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ: 1:1 mixture of rotamers, 0.61 (t, J=7.1 Hz, 1.5H), 0.72 (7.3, 1.5H), 0.76 (t, J=7.1, 1.5, 0.83, t, 7.3 Hz, 1.5H), 1.05-1.60 (m, 8H), 2.84-3.10 (m, J=2.5, 3.18, t, 9.7 Hz, 0.5H), 3.41-3.52 (m, 2H), 3.47-3.69 (m, 2H), 3.66 (s, 1.5H), 3.73 (s, 1.5H), 3.77 (s, 1.5H), 3.78 (s, 1.5H), 3.79 (s, 1.5H), 3.86 (d, J=9.8 Hz, 0.5H), 4.19 (d, J=17.7 Hz, 0.5H), 4.29 (d, J=15.2 Hz, 0.5H), 4.40-4.49 (m, 0.5H), 4.47 (d, J=15.3 Hz, 0.5H), 4.60 (d, J=17.6 Hz, 0.5H), 5.93 (m, 2H), 6.46 (dd, J=1.7, 8.2 Hz, 0.5H), 6.52 (d, J=2.0 Hz, 0.5H), 6.74 (m, 2.5H), 6.80 (s, 1H), 6.83-6.88 (m, 1H), 6.92 (m, 1.5H), 7.03 (dd, J=1.7, 6.8 Hz, 1H), 7.19 (m, 1H), 7.36 (m, 1H). MS (DCI) m/e 647 (M+H⁺). Anal calcd for C₃₇H₄₆N₂O₈: C, 68.71; H, 7.17; N, 4.33. Found: C, 68.41; H, 7.26; N, 4.11.

Example 365

20 *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((indolin-1-yl)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ: 2.97 (dd, J=8.1, 9.5 Hz, 1H), 3.10 (t, J=8.1 Hz, 2H), 3.16-3.22 (m, 2H), 3.51-3.68 (m, 3H), 3.73 (m, 3H), 3.83-4.05 (m, 3H), 5.90 (m, 2H), 6.73 (d, J=8.1 Hz, 1H), 6.86 (m, 3H), 6.99 (dt, J=1.1, 7.4 Hz, 1H), 7.08 (d, J=0.7 Hz, 1H), 7.11 (m, 1H), 7.18 (d, J=7.1 Hz, 1H), 7.38 (d, J=8.5 Hz, 2H), 8.02 (8.1, 1H). MS (C.I.) m/e 501 (M+H⁺). Anal calcd for C₂₉H₂₈N₂O₆ · 0.5 H₂O · 0.15 CH₃CO₂C₂H₅: C, 68.01; H, 5.82; N, 5.36. Found: C, 68.03; H, 5.65; N, 5.25.

Example 366

35 *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(2-chlorophenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ: 80.89 (dt, J=7 Hz, 3H), 1.23-1.51 (m, 4H), 2.52-4.00 (m, 8H), 3.78 (d, J=6 Hz, 3H), 5.92 (d, J=6 Hz, 2H), 6.70-6.87 (m, 4H),

7.02-7.21 (m, 4H), 7.27-7.52 (m, 3H). MS (DCI) m/e 565 (M+H)⁺. Analysis calcd for C₃₁H₃₂N₂O₆Cl · 0.6H₂O: C, 64.66; H, 5.99; N, 4.86. Found: C, 64.59; H, 6.00; N, 4.64.

5

Example 367

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3,4,5-trimethoxybenzyl)pyrrolidine-3-carboxylic acid

The compound resulting from Example 1C (0.25 g) was reacted with 0.169 g of 3,4,5-trimethoxybenzyl chloride and 0.175 g of diisopropylethylamine in 1 mL of acetonitrile for 2 hours at room temperature. The resulting ester was isolated and then hydrolyzed by the method of Example 1D to give 0.193 g of the title compound. m.p. 108-110 °C. ¹H NMR (300 MHz, CDCl₃) δ 2.75 (t, J=9Hz, 1H), 2.95-3.05 (m, 2H), 3.20 (d, J=11 Hz, 1H), 3.45-3.55 (m, 1H), 3.7-3.8 (m, 2H), 3.84 (s, 3H), 5.95 (dd, J=2Hz, 6Hz, 2H), 6.55 (s, 2H), 6.70 (d, J=8Hz, 1H), 6.30-6.35 (m, 1H), 6.90 (d, J=9Hz, 2H), 7.13 (d, J=2Hz, 1H), 7.43 (d, J=9Hz, 2H).

15

Example 368

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(3-chlorophenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.89 (t, J=7 Hz, 3H), 1.20-1.42 (m, 4H), 3.42-3.87 (m, 9H), 3.9 (s, 3H), 5.96 (s, 2H), 6.75 (7.10, J=m Hz, 7H), 7.33-7.50 (m, 4H). MS (C.I.) m/e 565 (M+H). Analysis calcd for C₃₁H₃₃N₂O₆Cl · 1.0CF₃COOH: C, 58.37; H, 5.05; N, 4.13. Found: C, 58.41; H, 4.99; N, 4.08.

20

25

Example 369

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(di-n-butylamino)pyrimidin-4-yl)pyrrolidine-3-carboxylic acid

The compound resulting from Example 1C (0.25 g) was reacted with 0.11 g of 2,4-dichloropyrimidine and 0.175 g of diisopropylethylamine in 1 mL of acetonitrile for 2 hours at room temperature to give 0.218 g of ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-chloro-4-pyrimidinyl)-pyrrolidine-3-carboxylate. This compound was reacted with 1 mL of dibutylamine in 2 mL of toluene at 125 °C for 17 hours. The resulting ethyl ester was hydrolyzed by

30

35

the method of Example 1D to give 0.142 g of the title compound as a white powder. ¹H NMR (300 MHz, CDCl₃) δ 0.75-0.90 (broad, 6H), 1.1-1.3 (br, 4H), 1.35-1.55 (br, 4H), 3.05 (m, 1H), 3.3-3.5 (br, 2H), 3.55-3.67 (m, 2H), 3.75 (s, 3H), 4.6 (br, 1H), 5.2 (br, 1H), 5.45 (br, 1H), 5.87 (s, 2H), 6.3 (br, 1H), 6.67 (d, J=8Hz, 1H), 6.7-6.85 (m, 4H), 7.10 (d, J=9Hz, 2H).

Example 370

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-(2-methylbut-2-yl)-N-phenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.90 (t, J=7.5 Hz, 3H), 1.12 (s, 3H), 1.14 (s, 3H), 2.06 (q, J=7.5 Hz, 2H), 2.73 (d, J=15.3 Hz, 1H), 2.91 (t, J=9.5 Hz, 1H), 3.11 (d, J=15.6 Hz, 1H), 3.21 (t, J=8.8 Hz, 1H), 3.50-3.61 (m, 2H), 3.80 (s, 3H), 4.00 (d, J=10.2 Hz, 1H), 5.91 (s, 2H), 6.74 (d, J=7.8 Hz, 1H), 6.85 (m, 3H), 6.93 (m, 1H), 6.98 (m, 1H), 7.03 (d, J=1.7 Hz, 1H), 7.17 (m, 2H), 7.36 (m, 3H). MS (DCI) m/e 545 (M+H⁺). Anal calcd for C₃₂H₃₆N₂O₆: C, 70.57; H, 6.66; N, 5.14. Found: C, 70.17; H, 6.53; N, 4.97.

Example 371

trans,trans-2-(4-Ethylphenyl)-4-(5-indanyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H (300MHz, CDCl₃) δ 7.25 (3H, m), 7.21 (1H, d, 3Hz), 7.17 (3H, m), 3.80 (1H, d, 10Hz), 3.65 (1H, ddd, 6, 5, 3Hz), 3.4 (4H, m), 3.10 (2H, m), 2.98 (2H, m), 2.88 (5H, m), 2.79 (1H, d, 16Hz), 2.62 (2H, q, 7Hz), 2.05 (2H, m), 1.42 (2H, m), 1.32 (1H, m), 1.21 (3H, t, 7Hz), 1.05 (2H, sext, 7Hz), 0.87 (3H, t, 7Hz), 0.79 (3H, t, 7Hz). MS (DCI, NH₃) m/e 505 (M+H⁺). Anal calcd for C₃₂H₄₄N₂O₃: C, 76.15; H, 8.79; N 5.55. Found: C, 75.96; H, 8.75; N, 5.36.

Example 372

trans,trans-2-(3,4-Difluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 62-63 °C. ¹H NMR (CDCl₃, 300 MHz), δ 0.83 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.13 (sextet, J=7Hz, 2H), 1.20-1.32 (m, 3H), 1.36-1.49 (m, 3H), 2.85-2.93 (m, 2H), 2.98-3.23 (m, 4H), 3.36-3.45 (m, 3H),

3.58-3.66 (m 1H), 3.94 (d, J=8Hz, 1H), 5.93 (s, 2H), 6.72 (d, J=7.5Hz, 1H), 6.84 (dd, J=1Hz, J=7.5Hz, 1H), 6.98 (d, J=7.5Hz, 1H), 7.08-7.15 (m, 2H), 7.22-7.28 (m, 1H). MS (CDI/NH₃) m/e 517 (M+H)⁺.

5

Example 373

trans,trans-2-(3,4-Difluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 71-72 °C. ¹H NMR (CDCl₃, 300

10 MHz) δ 0.82 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.25-1.38 (m, 4H), 1.46 (sextet, J=7Hz, 2H), 1.74 (quintet, J=7Hz, 2H), 2.26-2.36 (m, 1H), 2.72-2.95 (m, 5H), 2.98-3.12 (m, 2H), 3.15-3.34 (m, 2H), 3.45 (dd, J=3Hz, J=9Hz, 1H), 3.53-3.60 (m, 1H), 3.71 (d, J=9Hz, 1H), 5.96 (s, 2H), 6.75 (d, J=9Hz, 1H), 3.82 (cd., J=2Hz, J=9Hz, 1H), 5.96 (d, J=2Hz, 1H), 7.09-7.18 (m, 2H), 7.23-7.34 (m, 1H). MS (CDI/NH₃)
15 m/e 567 (M+H)⁺.

Example 374

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(ethoxymethyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

20 Using the procedures described in Example 1, the title compound was prepared. TLC (10% MeOH-CH₂Cl₂) R_f = 0.53. ¹H NMR (CDCl₃, 300 MHz, rotameric forms) δ 0.70 (t, J=7Hz), 0.80 (t, J=7Hz) and 0.96-1.04 (m, 6H total), 1.04-1.75 (m, 11H), 1.34-1.53 (br m, 4H), 2.65 (AB) and 2.80-3.08 (m, 2H total), 3.10-3.82 (br m, 12H), 4.03 (m) and 4.22-4.45 (br m, 2H total), 5.90 (s) and 5.91
25 (s, 2H total), 6.65-6.84 (m) and 6.93 (m) and 6.99 (m, 3H total). MS (FAB) m/e 463 (M+H)⁺. Anal calcd for C₂₅H₃₈N₂O₆ · 1.5 H₂O: C, 61.33; H, 8.44; N, 5.72. Found: C, 61.28; H, 7.78; N, 5.62.

Example 375

30

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(n-butyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a colorless wax. TLC (10% MeOH-CH₂Cl₂) R_f = 0.37.

¹H NMR (CDCl₃, 300 MHz, rotameric forms) δ 0.71 (t, J=7Hz) and 0.77-1.05 (m, 9H total), 1.05-1.20 (m, 2H), 1.20-1.72 (br m, 13H), 2.48-2.52 (m, 1H), 2.87-3.00 (m, 1H), 3.05-3.60 (m, 5H), 3.60-3.80 (br m, 2H), 3.88-4.05 (br m, 1H), 4.28 (br d,

J=15Hz, 1H total), 5.90 (s) and 5.92 (s, 2H total), 6.67-6.82 (m, 3H total). MS (FAB) m/e 461 (M+H)⁺. Anal calcd for C₂₆H₄₀N₂O₅ · 1.75 H₂O: C, 63.45; H, 8.90; N, 5.69. Found: C, 63.18; H, 8.22; N, 5.60.

Example 376

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(2-methylbutyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a colorless glass. TLC (10% MeOH-CH₂Cl₂) R_f = 0.49. ¹H NMR (CDCl₃, 300 MHz, rotameric forms and mixture of diastereomers) δ 0.69 (br t, J=7Hz) and 0.75-2.15 (several br m, approx. 26H total), 2.48-2.65 (br m, 1H), 2.87-3.01 (br m, 1H), 3.06-3.82 (br m, 7H), 3.90-4.40 (br m, 2H), 5.90 (s) and 5.92 (s, 2H total), 6.67-6.90 (m, 3H total). MS (FAB) m/e 475 (M+H)⁺.

Example 377.

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(3-methylbutyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. TLC (10% MeOH-CH₂Cl₂) R_f = 0.41. ¹H NMR (CDCl₃, 300 MHz, rotameric forms) δ 0.73 (t, J=7Hz) and 0.77-1.05 (m, 12H total), 1.07-1.75 (m, approx. 14H plus H₂O), 2.48-2.63 (m, 1H), 2.87-3.05 (m, 1H), 3.05-3.60 (several br m, 5H), 3.62-4.02 (br m, 2H), 4.29 (br d, J=15Hz, 1H), 5.89 (s) and 5.93 (s, 2H total), 6.65-6.90 (m, 3H total). MS (FAB) m/e 475 (M+H)⁺.

Example 378

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-((N-methyl-N-propylamino)sulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 58-59 °C. ¹H NMR (CDCl₃, 300MHz) δ 0.78 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.27 (sextet, J=7Hz, 2H), 1.48 (m, 4H), 2.22-2.30 (m, 1H), 2.62 (s, 3H), 2.68-2.78 (m, 1H), 2.84-3.03 (m, 5H), 3.08-3.31 (m, 3H), 3.39 (dd, J=3Hz, J=9Hz, 1H), 3.50-3.58 (m, 1H), 3.63 (d, J=9Hz, 1H), 3.79 (s, 3H), 5.95 (s, 2H), 3.73 (d, J=8Hz, 1H), 6.83 (dd, J=2Hz, J=8Hz, 1H), 3.87 (d, J=9Hz, 2H), 7.01 (d, J=2Hz, 1H), 7.33 (d, J=9Hz, 2H). MS (DCI/NH₃) m/e 576 (M+H)⁺.

Example 379trans,trans-2,4-Di(3,4-difluorophenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300MHz, CDCl₃) δ 7.35 (2H, m), 7.18 (4H, m), 4.87 (1H, d, J=12), 4.00-3.60 (5H, m), 3.60-3.10 (3H, m), 3.10-2.90 (2H, m), 1.45 (2H, m), 1.29 (4H, m), 1.15 (2H, m), 0.91 (3H, t, J=9), 0.83 (3H, t, J=9). MS (DCI/NH₃) m/e 509 (M+H⁺). Anal calcd for C₂₇H₃₂F₄N₂O₃· 0.75 TFA: C, 57.62; H, 5.56; N, 4.72. Found: C, 57.72; H, 5.67; N, 4.66.

Example 380trans,trans-4-(3,4-Dimethylphenyl)-2-(4-methoxyphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.43 (2H, d, J=9), 7.25 (1H, bs), 7.18 (1H, dd, J=8, 3), 7.11 (1H, d, J=9), 6.90 (2H, d, J=10), 5.48 (1H, d, J=12), 4.26 (1H, d, J=18), 4.16 (2H, m), 3.83 (2H, m), 3.81 (3H, s), 3.56 (1H, bd, J=18), 3.37 (1H, m), 3.20 (1H, m), 2.96 (2H, m), 2.24 (3H, s), 2.22 (3H, s), 1.47 (2H, m), 1.27 (4H, m), 1.10 (2H, m), 0.93 (3H, t, J=9), 0.81 (3H, t, J=9). MS (DCI/NH₃) m/e 495 (M+H⁺). Anal calcd for C₃₀H₄₂N₂O₄· 1.25 TFA: C, 61.26; H, 6.84; N, 4.40. Found: C, 61.16; H, 7.05; N, 4.38.

Example 381trans,trans-2,4-Di(3-fluoro-4-methoxyphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300MHz, CDCl₃) δ 7.20 (2H, m), 7.17 (2H, m), 6.93 (2H, m), 5.48 (1H, m), 4.26 (1H, m), 4.16 (2H, m), 3.83 (2H, m), 3.87 (6H, s), 3.56 (1H, m), 3.37 (1H, m), 3.20 (1H, m), 2.96 (2H, m), 1.47 (2H, m), 1.27 (4H, m), 1.10 (2H, m), 0.93 (3H, t, J=9), 0.81 (3H, t, J=9). MS (DCI/NH₃) m/e 533 (M+H⁺). Anal calcd for C₂₉H₃₈F₂N₂O₅· 0.75 H₂O: C, 63.78; H, 7.29; N, 5.13. Found: C, 63.77; H, 7.08; N, 4.99.

Example 382trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-(2-pentyl),N-(3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.90 (m, 3H), 0.95 (t, J=7.3 Hz, 3H), 1.13-1.37 (m, 4H), 2.30 (s, 3H), 2.34 (s (CH₃ rotamer)), 2.73-2.91 (m, 2H), 3.17-3.26 (m, 2H), 3.32-3.62 (m, 2H), 3.77-4.08 (m, 1H), 3.80 (s, 3H), 4.71 (m, 1H), 5.92 (m, 2H), 6.61-6.84 (m, 6H), 7.04-7.16 (m, 3H), 7.23-7.29 (m, 2H). MS (DCI) m/e 559 (M+H⁺). Anal calcd for C₃₃H₃₈N₂O₆ · 0.35 H₂O · 0.05 CH₃CO₂C₂H₅: C, 70.03; H, 6.92; N, 4.92. Found: C, 70.08; H, 6.82; N, 4.95.

Example 383

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(1-naphthyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.20-1.40 (m, 2H), 1.40-1.60 (m, 2H), 2.42-2.80 (m, 2H), 2.85-4.00 (m, 6H), 3.77 (d, J=1.5 Hz, 3H), 4.05-4.20 (m, 1H), 5.94 (d, J=2 Hz, 2H), 6.6 (dd, J=9, 10 Hz, 1H), 6.70-6.85 (m, 4H), 6.95-7.02 (m, 2H), 7.17 (dd, 8H, 1/2), 7.25 (dd, 8H, 1/2), 7.38-7.60 (m, 4H), 7.87-8.00 (m, 2H). MS (E.S.I.) m/e (M+H) 581. Analysis calcd for C₃₅H₃₆N₂O₆ · 1.4 H₂O: C, 69.38; H, 6.45; N, 4.62. Found: C, 69.36; H, 6.07; N, 4.41.

Example 384

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-phenyl-N-n-hexanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a tan solid. m.p. 67-68 °C. ¹H NMR (CD₃OD, 300 MHz) δ 0.88 (t, J=7Hz, 3H), 1.25-1.40 (m, 6H), 1.73 (quintet, J=7Hz, 2H), 2.13-2.23 (m, 1H), 2.64-2.88 (m, 3H), 3.02 (sextet, J=8Hz, 2H), 3.44-3.53 (m, 2H), 3.58 (d, J=9Hz, 1H), 3.56-3.75 (m, 1H), 3.78 (s, 3H), 3.88-3.98 (m, 1H), 5.93 (s, 2H), 6.72 (d, J=9Hz, 1H), 5.78-5.84 (m, 3H), 6.96 (d, J=2Hz, 1H), 7.20 (d, J=9Hz, 2H), 7.27-7.36 (m, 5H). MS (DCI/NH₃) m/e 609 (M+H)⁺.

Example 385

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(2-methyl-1,2,3,4-tetrahydroquinolin-1-yl)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 1.03 (m, 3H), 1.10-1.45 (m, 1H), 2.10-2.85 (m, 4H), 2.90-4.00 (m, 7H), 3.76 (s, 1.5H), 3.77 (s, 1.5H, isomer), 5.90 (m, 2H),

6.70-7.40 (m, 11H). MS (DCI) m/e 529 (M+H)⁺. Analysis calcd for C₃₁H₃₂N₂O₆ · 0.3 H₂O: C, 69.73; H, 6.15; N, 5.25. Found: C, 69.74; H, 6.10; N, 5.01.

5

Example 386

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(3-butyl-hept-2-en-1-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.86 (t, J=7.0 Hz, 3H), 0.90 (t, J=7.0 Hz, 3H), 1.20-1.41 (m, 8H), 1.95-2.06 (m, 4H), 3.24 (d, J=11.0 Hz, 1H), 3.51-3.59 (m, 3H), 3.60-3.71 (m, 1H), 3.77-3.84 (m, 1H), 3.81 (s, 3H), 4.45 (d, J=11.0 Hz, 1H), 5.52 (t, J=7.4 Hz, 1H), 5.93 (s, 2H), 6.77 (d, J=8.1 Hz, 1H), 6.87 (dd, J=1.8, 8.1 Hz, 1H), 6.99 (m, 3H), 7.46 (m, 2H). MS (DCI) m/e 494 (M+H)⁺. Anal calcd for C₃₀H₃₉NO₅: C, 72.99; H, 7.96; N, 2.84. Found: C, 72.73; H, 7.89; N, 2.64.

15

Example 387

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-hexanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 63-65 °C. ¹H NMR (CDCl₃, 300MHz) δ 0.82 (t, J=7Hz, 3H), 0.88 (t, J=6Hz, 3H), 1.23-1.47 (m, 6H), 1.44 (sextet, J=7Hz, 2H), 1.71 (quintet, J=6Hz, 2H), 2.24-2.34 (m, 1H), 2.70-2.93 (m, 5H), 2.96-3.12 (m, 2H), 3.15-3.35 (m, 2H), 3.43 (dd, J=3Hz, J=9Hz, 1H), 3.52-3.59 (m, 1H), 3.66 (d, J=9Hz, 1H), 3.87 (s, 3H), 5.95 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.82 (d, J=8Hz, 1H), 6.42 (t, J=8Hz, 1H), 6.96 (s, 1H), 7.12 (d, J=9Hz, 1H), 7.17 (d, J=12Hz, 1H). MS (DCI/NH₃) m/e 593 (M+H)⁺.

25

Example 388

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((3-pyridyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 2.87 (m, 2H), 3.04 (dd, J=3.2, 9.7 Hz, 1H), 3.21 (d, J=13.7 Hz, 1H), 3.51 (m, 1H), 3.76-3.85 (m, 2H), 3.79 (s, 3H), 5.90 (m, 2H), 6.71 (m, 1H), 6.79 (dd, J=1.7 Hz, 7.8Hz), 6.94 (m, 3H), 7.36-7.45 (m, 3H), 7.81 (m, 1H), 8.39 (m, 1H), 8.46 (dd, J=1.4 Hz, 1H). Anal calcd for C₂₅H₂₄N₂O₅ ·

35

0.70 H₂O · 0.05 CH₃CO₂C₂H₅: C, 67.34; H, 5.79; N, 6.23. Found: C, 67.31; H, 5.63; N, 5.90.

Example 389

5 *trans,trans*-2-(*n*-Hexyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) δ 0.82-1.00 (m, 9H), 1.20-1.40 (m, 12H), 1.45-1.60 (m, 4H), 1.70-1.90 (br m, 2H), 3.10-3.46 (m, 6H), 3.65 (t, J=10.8 Hz, 1H), 3.76 (t, J=11.0 Hz, 1H), 3.92-4.06 (m, 2H), 4.14-4.34 (m, 2H), 5.94 (s, 2H), 6.73 (d, J=8.1 Hz, 1H), 6.79 (dd, J=8.1, 1.8 Hz, 1H), 6.87 (d, J=1.8 Hz, 1H). MS(DCI/NH₃) m/e 489 (M+H)⁺. Anal calcd for C₂₈H₄₄N₂O₅ · 0.9 TFA: C, 60.53; H, 7.65; N, 4.74. Found: C, 60.62; H, 7.69; N, 4.61.

Example 390

15 *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((*N*-(2-pentyl)-*N*-(4-fluoro-3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.92 (m, 3H), 0.97 (t, J=7.1 Hz, 3H), 1.13-1.40 (m, 4H), 2.22 (m, 3H), 2.58-2.74 (m, 1H), 2.78-2.87 (m, 1H), 3.09-3.25 (m, 2H), 3.39-3.60 (m, 2H), 3.70-3.90 (m, 1H), 3.80 (s, 3H), 4.70 (m, 1H), 5.93 (m, 2H), 6.70-6.76 (m, 1H), 6.75 (dd, J=1.4, 8.1 Hz, 1H), 6.80-6.94 (m, 4H), 6.96-7.13 (m, 4H). MS (DCI) m/e 577 (M+H)⁺. Anal calcd for C₃₃H₃₇FN₂O₆ · 0.25 H₂O: C, 68.20; H, 6.50; N, 4.82. Found: C, 68.21; H, 6.46; N, 4.74.

Example 391

25 *trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((2-pyridyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 2.97 (dd, J=7.9, 9.7 Hz, 1H), 3.04 (t, J=9.6 Hz, 1H), 3.18 (dd, J=4.4 Hz, 9.9 Hz), 3.47 (d, J=14.0 Hz, 1H), 3.59 (m, 1H), 3.78 (s, 3H), 3.96 (d, J=9.9 Hz, 1H), 3.97 (d, J=13.6 Hz, 1H), 5.90 (m, 2H), 6.73 (d, J=8.1 Hz, 1H), 6.83 (dd, J=1.7, 7.9 Hz, 1H), 6.92 (m, 2H), 6.96 (d, J=1.8 Hz, 1H), 7.28 (m, 1H), 7.44 (m, 2H), 7.53 (d, J=8.1 Hz, 1H), 7.80 (dt, J=1.8, 7.7 Hz, 1H), 8.42 (m, 1H). MS (DCI) m/e 433 (M+H)⁺. Anal calcd for C₂₅H₂₄N₂O₅ · 0.35 H₂O: C, 68.43; H, 5.67; N, 6.38. Found: C, 68.44; H, 5.61; N, 6.24.

Example 392

trans, trans-2-(3-Phenylpropyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- 5 Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) δ 0.89-0.97 (m, 6H), 1.22-1.36 (m, 4H), 1.41-1.55 (m, 4H), 1.63-1.95 (m, 4H), 2.62 (dt, J=7.2, 2.1 Hz, 2H), 3.05-3.44 (m, 7H), 3.53-3.60 (m, 2H), 3.65-3.76 (m, 1H), 3.82-3.90 (m, 1H), 3.96-4.10 (m, 1H), 5.92 (s, 2H), 6.71 (d, J=8.1 Hz, 1H), 6.77 (dd, J=8.1, 1.5 Hz, 1H), 6.86 (d, J=1.2 Hz, 1H), 7.10-7.28 (m, 5H). MS(DCI/NH₃) m/e 523 (M+H)⁺. Anal calcd for C₃₁H₄₂N₂O₅ · 0.6 TFA: C, 65.43; H, 7.26; N, 4.74. Found: C, 65.28; H, 7.29; N, 4.50.

Example 393

- 15 trans-trans-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 115-117 °C. ¹H NMR (300 MHz, CDCl₃) δ 0.82 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.05-1.5 (m, 8H), 2.85 (d, J=13Hz, 1H), 2.90-3.17 (m, 5H), 3.20-3.35 (m, 1H), 3.35-3.50 (m, 3H), 3.55-3.65 (m, 1H), 3.84 (d, J=10Hz, 1H), 3.87 (s, 3H), 3.92 (s, 3H), 5.94 (dd, J=4Hz, 2Hz, 2H), 6.62 (s, 1H), 6.70 (s, 1H), 6.90 (t, J=8Hz, 1H), 7.05-7.20 (m, 2H).

Example 394

- 25 trans-trans-2-(1,4-Benzodioxan-6-yl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 107-110 °C. ¹H NMR (300 MHz, CDCl₃) δ 0.82 (t, J=7Hz, 3H), 0.88 (t, J=7Hz, 3H), 1.05-1.50 (m, 8H), 2.75 (d, J=13Hz, 1H), 2.90-3.12 (m, 4H), 3.32-3.60 (m, 5H), 3.69 (d, J= 8Hz, 1H), 3.90 (s, 3H), 4.23 (s, 4H), 5.95 (dd, J=4Hz, 2Hz, 2H), 6.62 (s, 1H), 6.70 (s, 1H), 6.78-6.93 (m, 3H).

Example 395

- 35 trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(3-butyl-2-fluoro-hept-2-en-1-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.84 (t, J=7.0 Hz, 3H), 0.88 (t, J=7.0 Hz, 3H), 1.16-1.37 (m, 8H), 1.83 (t, J=8.5 Hz, 2H), 2.03-2.08 (m, 2H), 2.76-2.92 (m, 2H), 3.02 (t, J=9.3 Hz, 1H), 3.32-3.42 (m, 2H), 3.50 (m, 1H), 3.71 (d, J=9.2 Hz, 1H), 3.78 (s, 3H), 5.91 (m, 2H), 6.72 (d, J=7.8 Hz, 1H), 6.83 (dd, J=1.7, 8.1 Hz, 1H), 6.90 (m, 2H), 7.02 (d, J=1.7 Hz, 1H), 7.34 (m, 2H). MS (DCI) m/e 512 (M+H)⁺. Anal calcd for C₃₀H₃₈FNO₅: C, 70.43; H, 7.49; N, 2.74. Found: C, 70.58; H, 7.54; N, 2.66.

Example 396

trans,trans-2-(3-Fluoro-4-ethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 65-66 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.26-1.36 (m, 4H), 1.41-1.52 (m, 5H), 1.73 (quintet, J=7Hz, 2H), 2.23-2.33 (m, 1H), 2.69-2.96 (m, 5H), 2.97-3.12 (m, 2H), 3.16-3.37 (m, 2H), 3.43 (d, J=9Hz, 1H), 3.52-3.59 (m, 1H), 3.66 (d, J=9Hz, 1H), 4.08 (q, J=7Hz, 2H), 5.95 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.82 (d, J=8Hz, 1H), 6.92 (t, J=8Hz, 1H), 6.97 (s, 1H), 7.07 (d, J=8Hz, 1H), 7.15 (d, J=12Hz, 1H). MS (DCI/NH₃) m/e 593 (M+H)⁺.

Example 397

trans,trans-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-butyl-N-propylamino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as a white solid. m.p. 118-120 °C. ¹H NMR (300 MHz, CDCl₃) δ 0.70-0.90 (4 triplets, J=7Hz), 1.05-1.55 (m, 8H), 2.80-3.50 (m, 9H), 3.55-3.65 (m, 1H), 3.82 (d, J= 10Hz, 1H), 3.85 (s, 3H), 3.92 (s, 3H), 5.96 (s, 2H), 6.62 (s, 1H), 6.70 (s, 1H), 6.90 (t, J=8Hz, 1H), 7.08-7.22 (m, 2H).

Example 398

trans,trans-4-(1,3-benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(4-chlorophenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.20-1.50 (m, 4H), 2.66-4.00 (m, 9H), 3.81 (s, 3H), 5.95 (s, 2H), 6.77 (d, J=7 Hz, 1H), 6.85 (d, J=8 Hz,

3H), 7.05 (m, 5H), 7.33-7.42 (m, 2H). MS (C.I.) m/e 565 (M+H). Analysis calcd for $C_{31}H_{33}N_2O_6Cl \cdot 0.25 H_3PO_4$: C, 63.16; H, 5.77; N, 4.75. Found: C, 63.14; H, 5.59; N, 4.53.

5

Example 399

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(4-methyl-1,2,3,4-tetrahydroquinolin-1-yl)carbonylmethylpyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. 1H NMR (300 MHz, CD_3OD) δ 1.27 (d, J=7 Hz, 1.5H), 1.28 (d, 7H, 1.5-diastereomer), 1.39-1.55 (m, 1H), 2.02-2.15 (m, 1H), 2.60-3.25 (m, 5H), 3.33-4.00 (m, 5H), 3.78 (s, 3H), 5.92 (d, J=3 Hz, 2H), 6.73 (dd, J=8 Hz, 1H), 6.75-6.90 (m, 3H), 6.91-7.35 (m, 7H). MS (DCI) m/e 529 (M+H)⁺. Analysis calcd for $C_{31}H_{32}N_2O_6$: C, 70.44; H, 6.10; N, 5.30. Found: C, 70.16; H, 6.04; N, 5.04.

15

Example 400

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-(piperidin-1-yl)ethanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 95-96 °C. 1H NMR ($CDCl_3$, 300MHz) δ 0.82 (t, J=7Hz, 3H), 1.43-1.55 (m, 4H), 1.63-1.72 (m, 4H), 2.29-2.38 (m, 1H), 2.64-2.78 (m, 5H), 2.87 (t, J=8Hz, 1H), 2.95-3.04 (m, 5H), 3.20-3.30 (m, 1H), 3.32-3.43 (m, 4H), 3.54-3.63 (m, 1H), 3.78 (d, J=8Hz, 1H), 3.87 (s, 3H), 5.92 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.78 (dd, J=2Hz, J=8Hz, 1H), 6.88 (t, J=8Hz, 1H), 6.94 (d, J=2Hz, 1H), 7.08-7.20 (m, 2H). MS (DCI/ NH_3) m/e 620 (M+H)⁺.

25

Example 401

trans,trans-2-(n-Heptyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. 1H NMR ($CDCl_3$, 300 MHz) δ 0.83-0.98 (s, 9H), 1.18-1.40 (m, 14H), 1.44-1.60 (m, 4H), 1.72-1.96 (br m, 2H), 3.12-3.45 (m, 6H), 3.65 (t, J = 10.5 Hz, 1H), 3.76 (t, J = 11.2 Hz, 1H), 3.90-4.06 (m, 2H), 4.13-4.33 (m, 2H), 5.93 (s, 2H), 6.73 (d, J = 7.8 Hz, 1H), 6.79 (dd, J = 7.8, 1.7 Hz, 1H), 6.87 (d, J = 1.7 Hz, 1H). MS (DCI/ NH_3) m/e 503 (M+H)⁺. Anal calcd for $C_{29}H_{46}N_2O_5 \cdot 0.75 TFA$: C, 62.28; H, 8.01; N, 4.76. Found: C, 62.20; H, 7.99; N, 4.50.

35

Example 402

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(3-methyl-1,2,3,4-tetrahydroquinolin-1-yl)carbonylmethylpyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.99 (d, 1.5H), 1.03 (d, J=6 Hz, 1.5H, second diastereomer), 2.60-4.00m (12), 3.78 (s, 1.5H), 3.79 (s, 1.5H, second diastereomer), 5.92 (s, 1H), 5.93 (s, 1H, diastereomer), 6.65-7.40 (m, 11H). MS (DCI) m/e 529 (M+H)⁺. Analysis calcd for C₃₁H₃₂N₂O₆ · 0.8 H₂O: C, 68.57; H, 6.24; N, 5.16. Found: C, 70.44; H, 6.10; N, 5.30.

Example 403

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(4-fluorophenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.2-1.47 (m, 4H), 2.7 (d, J=12 Hz, 1H), 2.80 (t, J=9 Hz, 1H), 3.09 (t, J=9 Hz, 1H), 3.25 (d, J=15 Hz, 1H), 3.40-3.47 (m, 1H), 3.49-3.65 (m, 3H), 3.75 (d, J=12 Hz, 1H), 3.80 (s, 3H), 5.94 (s, 2H), 6.72-6.86 (m, 4H), 7.00-7.15 (m, 7H). MS (DCI) m/e 549 (M+H)⁺. Analysis calcd for C₃₁H₃₃N₂O₆F · 0.4 H₂O: C, 66.99; H, 6.13; N, 5.04. Found: C, 66.99; H, 5.94; N, 4.99.

Example 404

trans,trans-1-(N-Butyl-N-(3-methylphenyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(5-benzofuranyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300MHz, CDCl₃) δ 7.66 (1H, bs), 7.60 (1H, d, J=3Hz), 7.45 (2H, s), 7.15 (4H, m), 6.75 (5H, m), 3.96 (1H, d, J=10Hz), 3.78 (3H, s), 3.74 (1H, m), 3.59 (3H, m), 3.21 (1H, t, J=9Hz), 3.19 (1H, d, J=16Hz), 2.92 (1H, t, J=9Hz), 2.70 (1H, d, J=16Hz), 2.29 (3H, s), 1.41 (2H, m), 1.24 (2H, m), 0.85 (3H, t, J=7Hz). MS (DCI, NH₃) m/e 541 (M+H)⁺. Anal. calcd for C₃₃H₃₄N₂O · 1 H₂O: C, 71.21; H, 6.52; N, 5.03. Found: C, 71.31; H, 6.30; N, 4.98.

Example 405

trans,trans-1-(N-Butyl-N-(3-methylphenyl)aminocarbonylmethyl)-2-(4-fluorophenyl)-4-(5-benzofuranyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 7.67 (1H, bs), 7.60 (1H, d, $J=3\text{Hz}$), 7.45 (2H, m), 7.18 (3H, m), 7.12 (1H, d, $J=7\text{Hz}$), 6.93 (2H, m), 6.76 (1H, d, $J=3\text{Hz}$), 6.70 (2H, bd), 4.02 (1H, m), 3.77 (1H, m), 3.59 (3H, m), 3.29 (1H, m), 3.19 (1H, m), 2.94 (1H, m), 2.71 (1H, m), 2.30 (3H, s), 1.45 (2H, m), 1.26 (2H, sext, $J=7\text{Hz}$), 0.84 (3H, t, $J=7\text{Hz}$). MS (DCI, NH_3) m/e 529 ($\text{M}+\text{H}^+$). Anal. calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{O}_5 \cdot 0.2 \text{ HOAc}$: C, 71.98; H, 6.30; N 5.18. Found : C, 71.68; H, 5.89; N, 5.25.

Example 406

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N,N-di-(3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CD_3OD) δ 2.27 (s, 6H), 2.81 (dd, $J=8.1, 9.5\text{Hz}$, 1H), 2.98 (d, $J=15.3\text{Hz}$, 1H), 3.20 (t, $J=16.6\text{Hz}$, 1H), 3.47-3.60 (m, 3H), 3.80 (s, 3H), 3.85 (d, $J=9.5\text{Hz}$, 1H), 5.91 (s, 2H), 6.73 (d, $J=7.8\text{Hz}$, 1H), 6.85 (m, 3H), 6.95 (m, 4H), 7.05 (d, $J=1.7\text{Hz}$, 1H), 7.06-7.24 (m, 6H). MS (DCI) m/e 579 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{35}\text{H}_{34}\text{N}_2\text{O}_6 \cdot 0.15 \text{ H}_2\text{O} \cdot 0.20 \text{ CH}_3\text{CO}_2\text{C}_2\text{H}_5$: C, 71.79; H, 6.04; N, 4.68. Found: C, 71.81; H, 5.79; N, 4.51.

Example 407

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H (300MHz, CDCl_3) δ 7.73 (2H, m), 7.40-7.10 (4H, m), 6.92 (2H, m), 6.72 (2H, d, $J=9$), 6.63 (1H, m), 5.40 (1H, m), 4.55 (2H, t, $J=9$), 4.30-4.10 (3H, m), 3.84 (3H, s), 3.82 (1H, m), 3.65 (1H, m), 3.39 (1H, m), 3.21 (2H, t, $J=9$), 3.10-2.90 (2H, m), 2.26 (3H, s), 1.55 (2H, m), 1.45 (2H, m), 0.92 (3H, t, $J=9$). MS (DCI/ NH_3) m/e 543 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{33}\text{H}_{38}\text{N}_2\text{O}_5 \cdot 0.65 \text{ H}_2\text{O}$: C, 71.50; H, 7.15; N, 5.05. Found: C, 71.47; H, 6.96; N, 4.83.

Example 408

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-(N,N-dimethylamino))ethanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 81-82 °C. ¹H NMR (CDCl₃, 300 MHz) δ 0.80 (t, J=7Hz, 3H), 1.43 (sextet, J=7Hz, 2H), 2.15-2.24 (m, 1H), 2.36 (s, 6H), 2.66-2.76 (m, 1H), 2.83-3.04 (m, 6H), 3.18-3.41 (m, 5H), 3.55-3.63 (m, 1H), 3.72 (d, J=8Hz, 1H), 3.85 (s, 3H), 5.90 (d, J=6Hz, 2H), 6.67 (d, J=8Hz, 1H), 6.78 (dd, J=2Hz, J=8Hz, 1H), 6.84 (t, J=8Hz, 1H), 7.94 (d, J=2Hz, 1H), 7.09 (d, J=8Hz, 1H), 7.20 (dd, J=2Hz, J=12Hz, 1H). MS (DCI/NH₃) m/e 580 (M+H)⁺.

Example 409

trans,trans-1-(N,N-Dibutylaminocarbonylmethyl)-2-(4-fluorophenyl)-4-(5-benzofuranyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.88 (1H, bs), 7.80 (2H, m), 7.61 (1H, d, J=3Hz), 7.55 (1H, bd, J=8Hz), 7.46 (1H, d, J=8Hz), 7.07 (2H, t, J=8Hz), 6.76 (1H, d, J=3Hz), 5.53 (1H, bd, J=11Hz), 4.18 (2H, m), 3.91 (3H, m), 3.55 (1H, d, J=16Hz), 3.30 (3H, m), 3.12 (1H, dd, J=10&9Hz), 2.95 (1H, m), 1.51 (2H, m), 1.31 (4H, m), 1.12 (2H, m), 0.92 (3H, m), 0.83 (3H, t, J=7Hz). MS m/e (DCI, NH₃) 595 (M+H)⁺.

Example 410

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-ethylphenyl)-1-(((N-butyl-N-(3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.35 (2H, m), 7.20-7.00 (7H, m), 6.70 (2H, d, J=9), 5.38 (1H, m), 4.55 (2H, t, J=9), 4.05 (1H, m), 3.64 (2H, m), 3.45 (1H, m), 3.21 (2H, t, J=9), 2.95 (1H, m), 2.75 (1H, m), 2.63 (2H, q, J=8), 2.38 (2H, m), 2.27 (3H, s), 1.43 (2H, m), 1.30 (2H, m), 1.22 (3H, t, J=9), 0.89 (3H, t, J=9). MS (DCI/NH₃) m/e 541 (M+H)⁺. Anal calcd for C₃₄H₄₀N₂O₄ · 1.6 AcOH: C, 70.17; H, 7.34; N, 4.40. Found: C, 70.11; H, 7.06; N, 4.80.

Example 411

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-fluorophenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.40 (2H, m), 7.28 (1H, bs), 7.18 (1H, dd, J=8, 3), 7.00 (2H, t, J=9), 6.72 (1H, d, J=9), 4.53 (2H, t, J=9), 3.92 (1H, m), 3.65 (1H, m), 3.42 (3H, m), 3.19 (2H, t, J=9), 3.15-2.90 (6H, m), 1.43 (3H, m), 1.25

(3H, m), 1.10 (2H, m), 0.90 (3H, t, J=8), 0.83 (3H, t, J=8). MS (DCI/NH₃) m/e 497 (M+H⁺). Anal calcd for C₂₉H₃₇FN₂O₄ · 0.25 H₂O: C, 69.51; H, 7.54; N, 5.59. Found: C, 69.45; H, 7.60; N, 5.44.

5 Example 412

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-fluorophenyl)-1-((N-butyl-N-(3-methylphenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.28 (1H, bs), 7.25-7.00 (5H, m), 6.91 (2H, m), 6.72 (3H, d, J=9), 4.54 (2H, t, J=9), 4.00 (1H, m), 3.60 (3H, m), 3.45 (1H, m), 3.19 (2H, t, J=9), 3.11 (2H, m), 2.84 (1H, m), 2.67 (1H, bd, J=18), 2.26 (3H, s), 1.42 (2H, m), 1.25 (2H, m), 0.88 (3H, t, J=8). MS (DCI/NH₃) m/e 531 (M+H⁺). Anal calcd for C₃₂H₃₅FN₂O₄ · 0.25 H₂O: C, 71.82; H, 6.69; N, 5.23. Found: C, 71.66; H, 6.55; N, 5.03.

15 Example 413

trans,trans-4-(Indan-5-yl)-2-(4-methoxyphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.32 (3H, m), 7.18 (2H, m), 6.85 (2H, d, J=9), 3.83 (1H, m), 3.79 (3H, s), 3.67 (1H, m), 3.50-3.20 (4H, m), 3.20-2.92 (4H, m), 2.87 (5H, m), 2.79 (1H, bd, J=15), 2.06 (2H, m), 1.43 (2H, m), 1.27 (4H, m), 1.08 (2H, m), 0.88 (3H, t, J=8), 0.82 (3H, t, J=8). MS (DCI/NH₃) m/e 507 (M+H⁺). Anal calcd for C₃₁H₄₂N₂O₄: C, 73.49; H, 8.36; N, 5.53. Found: C, 73.18; H, 8.29; N, 5.17.

25 Example 414

trans,trans-2-(4-Methoxyphenyl)-4-(3,4-difluorophenyl)-1-((N-butyl-N-(3-methylphenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300MHz, CDCl₃) δ 0.86 (t, J=7Hz, 3H), 1.10-1.35 (m, 2H), 1.35-1.52 (m, 2H), 2.29 (s, 3H), 2.63 (d, J=13Hz, 1H), 2.76 (t, J=7Hz, 1H), 3.06-3.20 (m, 2H), 3.42-3.53 (m, 1H), 3.50-3.64 (m, 3H), 3.80 (s, 3H), 3.86 (d, J=9Hz, 1H), 6.66-6.82 (m, 4H), 7.02-7.22 (m, 6H), 7.30-7.40 (m, 1H).

35 Example 415

trans,trans-1-(*N*-Butyl-*N*-(3-chlorophenyl)aminocarbonylmethyl)-2-(4-fluorophenyl)-4-(5-benzofuranyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.64 (1H, d, *J*=2Hz), 7.61 (1H, d, *J*=3Hz), 7.47 (1H, d, *J*=8Hz), 7.41 (1H, dd, *J*=8&3Hz), 7.30 (1H, dt, *J*=8&2Hz), 7.21 (1H, d, *J*=8Hz), 7.19 (2H, m), 7.00 (1H, bs), 6.94 (2H, t, *J*=8Hz), 6.83 (1H, bd, *J*=8Hz), 6.74 (1H, dd, *J*=2&1Hz), 3.96 (1H, d, *J*=10Hz), 3.75 (1H, ddd, 6, 5&3Hz), 3.59 (3H, m), 3.23 (1H, t, *J*=10Hz), 3.18 (1H, d, *J*=16Hz), 2.92 (1H, dd, *J*=10&9Hz), 2.69 (1H, d, *J*=16Hz), 1.41 (2H, m), 1.23 (2H, m), 0.87 (3H, t, *J*=7Hz). MS (DCI, NH₃) 549, 551 (M+H⁺). Anal. calcd for C₃₁H₃₀ClFN₂O₂: C, 67.82; H, 5.51; N, 5.10. Found: C, 67.43; H, 5.33; N, 4.78.

Example 416

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((*N*-propyl-*N*-(4-phenoxybenzyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ (rotamer) 7.40-7.20 (5H, m), 7.13 (2H, m), 6.98 (2H, m), 6.93-6.60 (7H, m), 5.93 (1H, d, *J*=2), 5.88 (5.85) (1H, d, *J*=2), 4.90 (4.50) (1H, d, *J*=15), 4.10 (4.25) (1H, d, *J*=15), 3.77 (3.73) (3H, s), 3.72 (1H, m), 3.60 (1H, m), 3.53-3.20 (3H, m), 3.15-2.75 (4H, m), 1.60-1.20 (2H, m), 0.83 (0.64) (3H, t, *J*=8). MS (DCI/NH₃) *m/e* 623 (M+H⁺). Anal calcd for C₃₇H₃₈N₂O₇ · 0.25 H₂O: C, 70.85; H, 6.19; N, 4.47. Found: C, 70.68; H, 6.10; N, 4.42.

Example 417

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-ethylphenyl)-1-(((*N*-(2-pentyl)-*N*-(4-fluoro-3-methylphenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.30 (1H, bs), 7.20-7.00 (5H, m), 6.87 (1H, m), 6.73 (2H, d, *J*=9), 6.57 (1H, m), 4.81 (1H, m), 4.55 (2H, t, *J*=9), 3.92 (1H, bd, *J*=11), 3.60 (1H, m), 3.43 (1H, m), 3.18 (2H, t, *J*=9), 3.17 (1H, m), 3.06 (1H, dd, *J*=15, 6), 2.88 (1H, dd, *J*=11, 9), 2.61 (2H, q, *J*=8), 2.59 (1H, m), 2.18 (3H, m), 1.40-1.10 (4H, m), 1.22 (3H, t, *J*=9), 1.00-0.80 (6H, m). MS (DCI/NH₃) *m/e* 573 (M+H⁺). Anal calcd for C₃₅H₄₁FN₂O₄ · 0.75 H₂O: C, 71.71; H, 7.31; N, 4.78. Found: C, 71.56; H, 7.33; N, 4.56.

Example 418

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-pyrimidinyl)amino)ethyl)pyrrolidine-3-carboxylic acid

- 5 Ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propylamino)propyl)pyrrolidine-3-carboxylate, prepared by the procedures of Example 61B (300 mg), 138 mg of 2-bromopyrimidine, and 150 mg of diisopropylethylamine were heated at 95 °C for 15 hours in 2 mL of acetonitrile. The resulting intermediate *trans-trans* ethyl ester was isolated by
- 10 chromatography on silica gel eluting with 5-10% ETOAc in CH₂Cl₂ and hydrolyzed with NaOH in ethanol/water to give 95 mg of the title compound. ¹H NMR (300 MHz, CDCl₃) δ 0.82 (t, J=7Hz, 3H), 1.50 (sextet, J=7Hz, 2H), 2.15-2.30 (m, 1H), 2.75-2.97 (m, 3H), 3.40-3.55 (m, 4H), 3.60-3.70 (m, 3H), 3.75 (s, 3H), 5.95 (s, 2H), 6.34 (t, J=4Hz, 1H), 6.65 (d, J=8Hz, 1H), 6.75-6.82 (m, 1H), 6.78 (d, J=9Hz, 2H), 6.96 (d, J=2Hz, 1H), 7.27 (d, J=9Hz, 2H), 8.20 (d, J=4Hz, 2H).
- 15

Example 419

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(3-butyl-2-chloro-
hept-2-en-1-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.84 (t, J=6.8 Hz, 3H), 0.88 (t, J=6.7 Hz, 3H), 1.19-1.39 (m, 8H), 2.05-2.09 (m, 2H), 2.17-2.23 (m, 2H), 2.78 (dd, J=6.6, 9.2 Hz, 1H), 2.95 (t, J=9.2 Hz, 1H), 3.32-3.37 (m, 2H), 3.49 (m, 1H), 3.70 (d, J=9.2 Hz, 1H), 3.77 (s, 3H), 5.91 (m, 2H), 6.72 (d, J=8.1 Hz, 1H), 6.85 (dd, J=1.9, 8.1 Hz, 1H), 6.89 (m, 2H), 7.08 (d, J=1.5 Hz, 1H), 7.36 (m, 2H). MS (DCI) m/e 528 (M+H⁺). Anal calcd for C₃₀H₃₈ClNO₅ · 0.25 H₂O: C, 67.66; H, 7.29; N, 2.63. Found: C, 67.62; H, 7.18; N, 2.40.

Example 420

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-methoxyphenyl)-1-(((N-(2-pentyl)-N-(4-fluoro-3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.28 (1H, bs), 7.15 (3H, m), 6.90 (1H, m), 6.77 (2H, dd, J=9, 3), 6.71 (2H, d, J=9), 6.56 (1H, m), 4.80 (1H, m), 4.53 (2H, t, J=9), 3.92 (1H, m), 3.79 (3H, s), 3.60 (1H, m), 3.45 (1H, m), 3.19 (2H, t, J=9), 3.18 (1H, m), 3.03 (1H, dd, J=15, 6), 2.85 (1H, m), 2.55 (1H, m), 2.18 (3H, m), 1.40-1.05 (4H, m), 1.00-0.80 (6H, m). MS (DCI/NH₃) m/e 575 (M+H⁺). Anal calcd for C₃₄H₃₉FN₂O₅ · 0.35 H₂O: C, 70.29; H, 6.89; N, 4.82. Found: C, 70.37; H, 6.92; N, 4.30.

Example 421

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(3-chlorophenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.29 (1H, d, J=3), 7.25-7.05 (5H, m), 6.98 (1H, bs), 6.80 (2H, m), 6.72 (2H, d, J=9), 4.53 (2H, t, J=9), 3.85 (1H, d, J=10), 3.79 (3H, s), 3.58 (3H, m), 3.42 (1H, dd, J=10, 6), 3.18 (4H, m), 2.87 (1H, m), 2.66 (1H, m), 1.40 (2H, m), 1.25 (2H, m), 0.86 (3H, t, J=9). MS (DCI/NH₃) m/e 563 (M+H⁺). Anal calcd for C₃₂H₃₅ClN₂O₅ · 0.25 H₂O: C, 67.72; H, 6.30; N, 4.94. Found: C, 67.72; H, 6.21; N, 4.55.

Example 422

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(5-ethylfuran-2-yl)-1-((N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.77 (1H, bs), 7.11 (1H, d, J=3), 7.02 (1H, dd, J=9, 3), 6.82 (1H, d, J=9), 6.52 (1H, d, J=4), 6.08 (1H, d, J=4), 5.98 (2H, s), 5.80 (1H, d, J=6), 4.70 (1H, bd, J=15), 4.37 (2H, m), 3.70 (2H, m), 3.39 (2H, m), 3.20 (1H, m), 3.10-2.82 (2H, m), 2.76 (2H, q, J=8), 1.45 (2H, m), 1.32 (3H, t, J=9), 1.30-1.10 (6H, m), 0.87 (3H, t, J=9), 0.85 (3H, t, J=9). MS (DCI/NH₃) m/e 499 (M+H⁺). Anal calcd for C₂₈H₃₈N₂O₆ · 1.75 HCl: C, 59.80; H, 7.12; N, 4.98. Found: C, 59.51; H, 6.96; N, 4.88.

Example 423

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-fluorophenyl)-1-(((N-(2-pentyl)-N-(4-fluoro-3-methylphenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.30-7.10 (4H, m), 6.92 (3H, m), 6.73 (2H, d, J=9), 6.59 (1H, m), 4.80 (1H, m), 4.53 (2H, t, J=9), 4.00 (1H, bd, J=10), 3.62 (1H, m), 3.45 (1H, m), 3.22 (1H, m), 3.21 (2H, t, J=9), 3.02 (1H, dd, J=15, 6), 3.85 (1H, t, J=10), 2.58 (1H, bd, J=18), 2.20 (3H, bs), 1.40-1.30 (3H, m), 1.15 (1H, m), 1.00-0.80 (6H, m). MS (DCI/NH₃) m/e 563 (M+H⁺). Anal calcd for C₃₃H₃₆F₂N₂O₄: C, 70.44; H, 6.45; N, 4.98. Found: C, 70.06; H, 6.47; N, 4.71.

Example 424

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-fluorophenyl)-1-(((N-butyl-N-(3-chlorophenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.30 (2H, m), 7.25-7.10 (4H, m), 6.95 (3H, m), 6.82 (1H, bd, J=9), 6.73 (1H, d, J=9), 4.55 (2H, t, J=9), 3.92 (1H, bd, J=11), 3.60 (3H, m), 3.43 (1H, dd, J=9, 6), 3.21 (2H, t, J=9), 3.16 (2H, m), 2.87 (1H, m), 2.69 (1H, m), 1.42 (2H, m), 1.26 (2H, m), 0.87 (3H, t, J=9). MS (DCI/NH₃) m/e 551 (M+H⁺). Anal calcd for C₃₁H₃₂ClFN₂O₄ · 0.25 H₂O: C, 67.02; H, 5.90; N, 5.04. Found: C, 66.98; H, 5.71; N, 4.76.

Example 425

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-ethylphenyl)-1-(((N-butyl-N-(3-chlorophenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.30 (1H, m), 7.21 (1H, d, J=9), 7.15 (2H, m), 7.09 (4H, bs), 6.96 (1H, bs), 6.80 (1H, bd, J=9), 6.73 (1H, d, J=9), 4.54 (2H, t, J=9), 3.89 (1H, bd, J=11), 3.60 (3H, m), 3.43 (1H, m), 3.22 (2H, t, J=9), 3.18 (2H, m), 2.92 (1H, m), 2.72 (1H, m), 2.62 (2H, q, J=8), 1.41 (2H, m), 1.26 (2H, m), 1.23 (3H, t, J=9), 0.87 (3H, t, J=9). MS (DCI/NH₃) m/e 561 (M+H⁺). Anal. calcd for C₃₃H₃₇ClN₂O₄ · 0.25 H₂O: C, 70.08; H, 6.68; N, 4.95. Found: C, 70.13; H, 6.59; N, 4.65.

Example 426

trans,trans-1-(N-Butyl-N-(3-chlorophenyl)carboxamidomethyl)-2-(4-methoxyphenyl)-4-(5-benzofuranyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 7.67 (1H, bs), 7.60 (1H, d, J=3Hz), 7.48 (1H, d, J=8Hz), 7.42 (1H, dd, J=8&3Hz), 7.29 (1H, dt, J=8&3Hz), 7.21 (1H, d, J=8Hz), 7.14 (2H, m), 6.99 (1H, bs), 6.76 (4H, m), 3.88 (1H, d, J=10Hz), 3.75 (1H, ddd, J=6, 5&3Hz), 3.59 (2H, m), 3.53 (1H, dd, J=10&3Hz), 3.22 (1H, t, J=9Hz), 3.19 (1H, m), 2.96 (1H, m), 2.70 (1H, d, J=16Hz), 1.42 (2H, m), 1.26 (2H, m), 0.87 (3H, t, J=7Hz). MS (DCI, NH₃) m/e 563, 561 (M+H⁺). Anal. calcd for C₃₂H₃₃ClN₂O₅ · 0.5 H₂O: C, 67.42; H, 6.01; N, 4.91. Found: C, 67.45; H, 5.82; N, 4.68.

Example 427

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-cyclohexyl-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) (rotamer) δ 0.78 (0.86) (t, 3H, J=7Hz), 0.90-1.90 (envelope, 14H), 2.69 (2.80) (d, 1H, J=12Hz), 2.9-3.8 (envelope, 10H), 3.78 (3.80) (s, 3H), 5.92 (s, 2H), 6.72 (d, 1H, J=9Hz) 6.86 (m, 3H) 7.03 (d, 1H, J=6Hz), 7.34 (m, 2H). MS (DCI/NH₃) m/e 537 (M+H)⁺. Anal. calc'd for C₃₁H₄₀N₂O₆ · 1 H₂O: C, 67.13; H, 7.63; N, 5.05. Found: C, 67.09; H, 7.34; N, 4.92.

Example 428

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-ethylphenyl)-1-(((N-(3-methylphenyl)-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 0.86 (t, 3H, J=7Hz), 1.22 (t, 3H, J=7Hz), 1.25 (m, 2H), 1.43 (m, 2H), 2.26 (s, 3H), 2.6 (q, 2H, J=7Hz), 2.68 (d, 1H, J=12Hz), 2.86 (t, 1H, J=8Hz), 3.19 (q, 2H, J=7Hz), 3.44 (dd, 1H, J= 3Hz, 10Hz), 3.59 (m, 3H), 3.94 (d, 1H, 9Hz), 5.92 (s, 2H), 6.75 (m, 3H), 6.86 (dd, 1H, J= 2Hz, 8Hz), 7.08 (m, 6H), 7.17 (t, 1H, J= 8Hz). MS (DCI/NH₃) m/e 543 (M+H)⁺. Anal. calc'd for C₃₃H₃₈N₂O₅ · 0.60 H₂O: C, 71.61; H, 7.14; N, 5.06. Found: C, 71.57; H, 6.80; N, 4.87.

Example 429

trans,trans-4-(Benzofuran-5-yl)-2-(4-ethylphenyl)-1-(((N-(3-methylphenyl)-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 0.90 (t, 3H, J=7Hz), 1.30 (t, 3H, J=7Hz), 1.31 (m, 2H), 1.43 (m, 2H), 2.27 (s, 3H), 2.73 (q, 2H, J=7Hz), 3.15 (d, 2H, J=17Hz), 3.61 (t, 2H, J= 8Hz), 3.82 (m, 2H), 4.00 (t, 1H, 12Hz), 4.26 (m, 2H), 5.53 (br d, 1H), 6.54 (br s, 2H), 6.76 (d, 1H, J= 2Hz), 7.14 (m, 3H), 7.28 (s, 1H), 7.40 (m, 3H), 7.48 (d, 1H, J= 8Hz), 7.63 (d, 1H, J=2Hz), 7.73 (s, 1H). HRMS. calc'd for C₃₄H₃₉N₂O₄ (M+H)⁺: 539.2910. Found: 539.2891

Example 430

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-ethylphenyl)-1-(((N-(3-methylphenyl)-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 0.87 (t, 3H, J=7Hz), 1.22 (t, 3H, J=7Hz), 1.24 (m, 2H), 1.42 (m, 2H), 2.30 (s, 3H), 2.61 (q, 2H, J=7Hz), 2.67 (d, 1H, J=14Hz), 2.86 (t, 1H, J= 8Hz), 3.18 (q, 2H, J=7Hz), 3.41 (dd, 1H, J=4,10Hz), 3.59 (m, 3H), 3.93 (d, 1H, J=10Hz), 4.25 (m, 4H), 6.74 (br s, 2H), 6.80 (d, 1H, J=8Hz), 6.93 (dd, 1H, J=2Hz,8Hz), 6.99 (d, 1H, J=2Hz), 7.07 (m, 5H), 7.17 (t, 1H, J=8Hz). MS (DCI/NH₃) m/e 557 (M+H)⁺. Anal. calc'd for C₃₄H₄₀N₂O₅ · 0.40 H₂O: C, 72.42; H, 7.29; N, 4.97. Found: C, 72.49; H, 7.16; N, 4.62.

Example 431

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-2-mesitylenesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 80-82 °C. ¹H NMR (CDCl₃, 300

- 5 MHz) δ 0.69 (t, J=7Hz, 3H), 1.37 (sextet, J=7Hz, 2H), 2.09-2.17 (m, 1H), 2.24 (s, 3H), 2.53 (s, 6H), 2.54-2.64 (m, 1H), 2.73-2.86 (m, 2H), 3.02 (sextet, J=7Hz, 2H), 3.13-3.28 (m, 3H), 3.44-3.53 (m, 1H), 3.57 (d, J=9Hz, 1H), 3.89 (s, 3H), 5.94 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.78 (dd, J=2Hz, J=8Hz, 1H), 6.85 (s, 2H), 6.92 (d, J=8Hz, 1H), 9.94 (d, J=2Hz, 1H), 7.06 (d, J=8Hz, 1H), 7.13 (dd, J=2Hz, J=12Hz, 1H). MS (DCI/NH₃) m/e 627 (M+H)⁺.
- 10

Example 432

trans,trans-2-(4-Methoxyphenyl)-4-(3,4-difluorophenyl)-1-((N-butyl-N-(3-chlorophenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 0.86 (t, J=7Hz, 3H), 1.18-1.32 (m, 2H), 1.35-1.48 (m, 2H), 2.64 (d, J=13Hz, 1H), 2.71 (t, J= 7Hz, 1H), 3.08-3.18 (m, 2H), 3.42-3.48 (m, 1H), 3.53-3.64 (m, 3H), 3.77 (s, 3H), 3.80 (d, J=9Hz, 1H), 6.73-6.85 (m, 3H), 6.94 (s, 1H), 7.04-7.40 (m, 7H).

20

Example 433

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(3-chloropropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.80 (t, 3H, J=7), 1.47 (bd hex, 2H, J=8), 2.15 (pen, 2H, J=7), 2.32 (m, 1H), 2.7-3.2 (m, 9H), 3.46 (dd, 1H, J=4, 10), 3.57 (m, 1H), 3.64 (t, 2H, J=6), 3.67 (d, 1H, J=9), 3.86 (s, 3H), 5.92 (s, 2H), 6.74 (d, 1H, J=8), 6.84 (dd, 1H, J=2, 8), 6.96 (d, 1H, J=2), 7.06 (t, 1H, J=9), 7.18 (m, 2H). MS (DCI/NH₃) m/e 585 (M+H; ³⁵Cl)⁺; 587 (M+H; ³⁷Cl)⁺. Anal calcd for C₂₇H₃₄N₂O₇ClFS: C, 55.43; H, 5.86; N, 4.79. Found: C, 55.65; H, 5.81; N, 4.70.

25

30

Example 434

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(3-chloropropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.79 (d, 3H, J=7), 0.84 (d, 3H,

35

J=7), 1.68 (hept, 1H, J=7), 2.18 (pen, 2H, J=7), 2.8-3.4 (m, 10H), 3.5-3.8 (m, 3H), 3.65 (t, 2H, J=6), 3.90 (s, 3H), 5.94 (s, 2H), 6.77 (d, 1H, J=8), 6.87 (dd, 1H, J=2, 8), 6.99 (d, 1H, J=2), 7.13 (t, 1H, J=9), 7.27 (m, 2H). MS (DCI/NH₃) m/e 599 (M+H)⁺. Anal calcd for C₂₈H₃₆N₂O₇ClFS · 0.3 TFA: C, 54.24; H, 5.78; N, 4.42.

5 Found: C, 54.19; H, 5.71; N, 4.01.

Example 435

trans,trans-2-Propoxymethyl-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- 10 Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) δ 0.87-0.98 (m, 9H), 1.21-1.39 (m, 4H), 1.43-1.57 (m, 4H), 1.58-1.70 (m, 2H), 3.13-3.29 (m, 4H), 3.34-3.43 (m, 3H), 3.45-3.55 (m, 3H), 3.69 (dd, J = 10.2, 4.5 Hz, 1H), 3.80-4.20 (m, 4H), 5.93 (s, 2H), 6.73 (d, J = 7.8 Hz, 1H), 6.84 (dd, J = 8.2, 1.7 Hz, 1H), 6.93 (d, J = 1.7 Hz, 1H).
- 15 MS(DCI/NH₃) m/e 477 (M+H)⁺. Anal calcd for C₂₆H₄₀N₂O₆·0.50 TFA: C, 60.77; H, 7.65; N, 5.25. Found: C, 60.73; H, 7.74; N, 5.22.

Example 436

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(4-methylbutanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

- 20 Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 65-67 °C. ¹H NMR (CDCl₃, 300MHz) δ 0.82 (t, J=7Hz, 3H), 0.88 (d, J=5Hz, 6H), 1.46 (sextet, J=7Hz, 2H), 1.56-1.64 (m, 3H), 2.24-2.33 (m, 1H), 2.68-2.93 (m, 5H), 2.98-3.12 (m, 2H), 3.15-3.35 (m, 2H), 3.43 (dd, J=3Hz, J=9Hz, 1H), 3.52-3.58 (t, 1H), 3.65 (d, J=12Hz, 1H), 3.87 (s, 3H), 5.95 (s, 2H), 6.73 (d, J=8Hz, 1H), 6.82 (dd, J=2Hz, J=8Hz, 1H), 6.92 (t, J=8Hz, 1H), 6.97 (d, J=2Hz, 1H), 7.10 (d, J=9Hz, 1H), 7.16 (dd, J=2Hz, J=12Hz, 1H). MS (DCI/NH₃) m/e 579 (M+H)⁺.
- 25

30 Example 437

trans,trans-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(n-pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

- Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (300MHz, CDCl₃) δ 0.81 (t, J=7Hz, 3H), 0.90 (t, J=9Hz, 3H), 1.25-1.35 (m, 4H), 1.44 (sextet, J=7Hz, 2H), 1.67-1.78 (m, 2H), 2.22-2.34 (m, 1H), 2.30-2.95 (m, 5H), 2.95-3.10 (m, 2H), 3.15-3.33 (m, 2H), 3.45 (dd, J=3Hz, 9Hz, 1H),
- 35

3.47-3.56 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.88 (s, 3H), 3.94 (s, 3H), 5.95 (s, 2H), 6.55 (s, 1H), 6.65 (s, 1H), 6.92 (t, J=7H, 1H), 7.11 (d, J=9Hz, 1H), 7.17 (d, J=12Hz, 1H).

Example 438

- 5 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2,2,3,3,3-pentafluoropropoxyethanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 63-64 °C. ¹H NMR (CDCl₃,

- 10 300MHz) δ 0.82 (t, J=7Hz, 3H), 1.45 (sextet, J=7Hz, 2H), 2.24-2.33 (m, 1H), 2.70-2.82 (m, 1H), 2.85-3.09 (m, 5H), 3.14-3.28 (m, 4H), 3.43 (dd, J=3Hz, J=9Hz, 1H), 3.52-3.58 (m, 1H), 3.65 (d, J=9Hz, 1H), 3.87 (s, 3H), 3.92-3.98 (m, 3H), 5.94 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.82 (dd, J=2Hz, J=8Hz, 1H), 6.92 (t, J=8Hz, 1H), 6.97 (d, J=2Hz, 1H), 7.10 (d, J=9Hz, 1H), 7.17 (dd, J=2Hz, J=12Hz, 1H). MS (DCI/NH₃)
15 m/e 685 (M+H)⁺.

Example 439

- trans,trans*-2-(1,4-Benzodioxan-6-yl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(*n*-pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid
- 20 Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CDCl₃) δ 0.81 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.23-1.36 (m, 4H), 1.45 (sextet, J=7Hz, 2H), 1.65-1.78 (m, 2H), 2.20-2.30 (m, 1H), 2.30-2.95 (m, 5H), 2.95-3.10 (m, 2H), 3.15-3.35 (m, 2H), 3.42 (dd, J=3Hz, 9Hz, 1H), 3.46-3.56 (m, 1H), 3.59 (d, J=9Hz, 1H), 3.91 (s, 3H), 4.24 (s, 4H), 5.95 (s, 2H), 6.57 (s, 1H),
25 6.68 (s, 1H), 6.82 (d, J=8Hz, 1H), 6.88 (dd, J=2Hz, 8Hz, 1H), 6.95 (d, J=2Hz, 1H).

Example 440

- trans,trans*-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(4-methoxybenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid
- 30 Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ (rotamer) 7.32 (1H, d, J=10), 7.22 (1H, m), 7.10 (1H, d, J=9), 7.03 (6.98) (1H, d, J=3), 6.90-6.80 (4H, m), 6.79 (2H, d, J=9), 6.77 (1H, t, J=8), 5.85 (2H, s), 4.92 (4.10) (1H, d, J=15), 4.42 (4.22) (1H, d, J=15), 3.81 (1H, m), 3.79 (3.78) (3H, s), 3.76 (3H, s), 3.62 (1H, m), 3.43 (2H, m),
35 3.30-2.70 (5H, m), 1.42 (1H, m), 1.23 (2H, m), 1.01 (1H, m), 0.83 (0.75) (3H, t,

J=8). MS (DCI/NH₃) m/e 575 (M+H⁺). Anal calcd for C₃₃H₃₈N₂O₇ · 0.5 H₂O: C, 67.91; H, 6.73; N, 4.80. Found: C, 67.78; H, 6.44; N, 4.55.

Example 441

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.76 (d, 3H, J=7), 0.84 (d, 3H, J=7), 0.92 (t, 3H, J=7), 1.36 (m, 4H), 1.70 (m, 3H), 2.90 (m, 2H), 3.02 (m, 2H), 3.1-3.8 (m, 7H), 3.84 (d, 2H, J=8), 3.91 (s, 3H), 5.96 (s, 2H), 6.80 (d, 1H, J=8), 6.88 (dd, 1H, J=2, 8), 7.00 (d, 1H, J=2), 7.19 (t, 1H, J=9), 7.35 (m, 2H). MS (DCI/NH₃) m/e 593 (M+H⁺). Anal calcd for C₃₀H₄₁N₂O₇F · 0.5 TFA: C, 57.31; H, 6.44; N, 4.31. Found: C, 57.08; H, 6.15; N, 3.95.

Example 442

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(3-fluorophenylamino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.10-1.30 (m, 4H), 2.70-2.90 (m, 2H), 3.13 (t, J=8 Hz, 1H), 3.40-3.90 (m, 6H), 3.79 (s, 3H), 5.93 (s, 2H), 6.75 (d, J=8 Hz, 1H), 6.80-7.20 (m, 9H), 7.40 (m, 1H). MS (DCI) m/e 549 (M+H⁺). Anal calcd for C₃₁H₃₃N₂O₆F · 0.8 H₂O: C, 66.13; H, 6.19; N, 4.98. Found: C, 66.21; H, 5.83; N, 4.84.

Example 443

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-fluorophenyl)-1-(N-butyl-N-(3-chlorophenylamino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.20-1.50 (m, 4H), 2.65-2.85 (m, 2H), 3.05-3.85 (m, 7H), 5.93 (s, 2H), 6.75 (d, J=8 Hz, 1H), 6.85 (dd, J=8 Hz, 1H), 6.90-7.10 (m, 4H), 7.10-7.25 (m, 3H), 7.33-7.45 (m, 2H). MS (DCI) m/e 553 (M+H⁺). Anal calcd for C₃₀H₃₀N₂O₅FCI: C, 65.16; H, 5.47; N, 5.07. Found: C, 65.37; H, 5.41; N, 4.98.

Example 444

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(3,4-dimethoxybenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ (rotamer) 7.33 (1H, d, J=10), 7.23 (1H, m), 7.03 (6.97) (1H, d, J=3), 6.90-6.60 (6H, m), 6.47 (1H, m), 5.93 (2H, m), 4.83 (4.09) (1H, d, J=15), 4.45 (4.22) (1H, d, J=15), 3.83 (3.86) (3H, s), 3.79 (1H, m), 3.77 (3.76) (3H, s), 3.75 (3.65) (3H, s), 3.60 (1H, m), 3.43 (2H, m), 3.28 (1H, m), 3.20-2.70 (4H, m), 1.43 (1H, m), 1.23 (2H, m), 1.02 (1H, m), 0.84 (0.77) (3H, t, J=8). MS (DCI/NH₃) m/e 605 (M+H⁺). Anal calcd for C₃₄H₄₀N₂O₈: C, 67.53; H, 6.67; N, 4.63. Found: C, 67.28; H, 6.63; N, 4.38.

Example 445

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(2-methoxybenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ (rotamer) 7.33 (1H, d, J=10), 7.11 (2H, m), 7.03 (1H, dd, J=8, 3), 6.90-6.60 (7H, m), 5.93 (2H, m), 4.83 (4.15) (1H, d, J=15), 4.47 (4.30) (1H, d, J=15), 3.81 (1H, m), 3.78 (3.73) (3H, s), 3.72 (3H, s), 3.59 (1H, m), 3.43 (2H, m), 3.30 (1H, m), 3.20-2.70 (4H, m), 1.42 (1H, m), 1.23 (2H, m), 1.01 (1H, m), 0.83 (0.77) (3H, t, J=8). MS (DCI/NH₃) m/e 575 (M+H⁺). Anal calcd for C₃₃H₃₈N₂O₇: C, 68.97; H, 6.66; N, 4.87. Found: C, 68.70; H, 6.56; N, 4.61.

Example 446

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(3-methoxybenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ (rotamer) 7.31 (1H, d, J=10), 7.13 (1H, d, J=9), 7.16 (1H, dt, J=8, 3), 7.03 (1H, dd, J=10, 2), 6.90-6.60 (6H, m), 6.50 (1H, m), 5.94 (2H, m), 4.82 (4.19) (1H, d, J=15), 4.50 (4.23) (1H, d, J=15), 3.78 (3.76) (3H, s), 3.77 (1H, m), 3.75 (3.67) (3H, s), 3.59 (1H, m), 3.57-3.35 (2H, m), 3.25 (1H, m), 3.20-2.70 (4H, m), 1.43 (1H, m), 1.23 (2H, m), 1.02 (1H, m), 0.84 (0.77) (3H, t, J=8). MS (DCI/NH₃) m/e 575 (M+H⁺). Anal calcd for C₃₃H₃₈N₂O₇: C, 68.97; H, 6.66; N, 4.87. Found: C, 68.72; H, 6.55; N, 4.60.

Example 447

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(3-chloropropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 2.15 (pen, 2H, J=7), 2.33 (m, 1H), 2.81 (m, 2H); 2.93 (t, 1H, J=9); 3.1-3.6 (m, 10H), 3.24 (s, 3H); 3.65 (t, 2H, J=6), 3.70 (d, 1H, J=9), 3.87 (s, 3H), 5.92 (s, 2H), 6.74 (d, 1H, J=8), 6.84 (dd, 1H, J=2, 8), 6.97 (d, 1H, J=2), 7.07 (t, 1H, J=9), 7.17 (m, 2H). MS (DCI/NH₃) m/e 601 (M+H)⁺. Anal calcd for C₂₇H₃₄N₂O₈ClFS: C, 53.95; H, 5.70; N, 4.66. Found: C, 53.65; H, 5.49; N, 4.26.

Example 448

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.93 (m, 3H), 1.34 (m, 4H), 1.69 (m, 2H), 2.33 (m, 1H), 2.75-3.1 (m, 7H), 3.23 (s, 3H), 3.3-3.6 (m, 6H), 3.70 (d, 1H, J=9), 3.86 (s, 3H), 5.92 (s, 2H), 6.74 (d, 1H, J=8), 6.84 (dd, 1H, J=2, 8), 6.97 (d, 1H, J=2), 7.07 (t, 1H, J=9), 7.18 (m, 2H). MS (DCI/NH₃) m/e 595 (M+H)⁺. Anal calcd for C₂₉H₃₉N₂O₈FS: C, 58.57; H, 6.61; N, 4.71. Found: C, 58.21; H, 6.29; N, 4.29.

Example 449

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-(4-heptyl)-N-(4-fluoro-3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.89 (m, 6H), 1.18-1.36 (m, 8H), 2.15 (bs, 1.5 (CH₃ rotamer)), 2.28 (bs, 1.5 (CH₃ rotamer)), 2.64 (t, J=14.9 Hz, 1H), 2.82 (m, 1H), 3.07-3.29 (m, 2H), 3.32-3.41 (m, 1H), 3.53-3.60 (m, 1H), 3.70-3.79 (m, 1H), 3.79 (s, 3H), 4.68 (m, 1H), 5.92 (m, 2H), 6.69-6.90 (m, 6H), 6.93-7.07 (m, 4H). MS (DCI) m/e 605 (M+H)⁺. Anal calcd for C₃₅H₄₁N₂O₆: C, 69.52; H, 6.83; N, 4.63. Found: C, 69.31; H, 6.78; N, 4.35.

Example 450

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-(5-nonyl)-N-(4-fluoro-3-methylphenyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CD_3OD) δ 0.81-0.90 (m, 6H), 1.30 (m, 12H), 2.14 (s, 1.5 (CH_3 rotamer)), 2.30 (s, 1.5 (CH_3 rotamer)), 2.60 (t, $J=14.8$ Hz, 1H), 2.80 (m, 1H), 3.09-3.24 (m, 2H), 3.33-3.42 (m, 1H), 3.50-3.55 (m, 1H), 3.65-3.77 (m, 1H), 3.79 (s, 3H), 4.64 (m, 1H), 5.93 (m, 2H), 6.70-6.84 (m, 5H), 6.91-7.13 (m, 5H). MS (DCI) m/e 633 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{37}\text{H}_{45}\text{FN}_2\text{O}_6$: C, 70.23; H, 7.17; N, 4.43. Found: C, 70.14; H, 7.13; N, 4.19.

Example 451

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-(5-nonylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CD_3OD) δ 0.80 (t, $J=7.0$ Hz, 3H), 0.84 (t, $J=7.1$ Hz, 3H), 1.15-1.55 (m, 12H), 2.88 (d, $J=15.9$ Hz, 1H), 3.07 (m, 2H), 3.26 (d, $J=16.3$ Hz, 1H), 3.36 (dd, $J=4.4, 9.8$ Hz, 1H), 3.64 (m, 1H), 3.76 (m, 1H), 3.79 (s, 3H), 3.98 (d, $J=9.5$ Hz, 1H), 5.93 (m, 2H), 6.77 (d, $J=7.8$ Hz, 1H), 6.85 (dd, $J=1.7, 8.1$ Hz, 1H), 6.93 (m, 2H), 6.99 (d, $J=1.7$ Hz, 1H), 7.39 (m, 2H). MS (DCI) m/e 525 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{30}\text{H}_{46}\text{N}_2\text{O}_6 \cdot 0.35 \text{H}_2\text{O}$: C, 67.86; H, 7.73; N, 5.28. Found: C, 67.87; H, 7.63; N, 5.11.

Example 452

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(2-fluorophenyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CD_3OD) δ 0.87 (dt, $J=7$ Hz, 3H), 1.15-1.32 (m, 4H), 3.77 (d, $J=2$ Hz, 3H), 2.65-5.92 (m, 9H), 5.93 (d, $J=4$ Hz, 2H), 6.70-6.90 (m, 4H), 7.00-7.45 (m, 7H). MS (DCI) m/e 549 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{31}\text{H}_{33}\text{N}_2\text{O}_6 \cdot 0.4 \text{H}_2\text{O}$: C, 66.99; H, 6.13; N, 5.04. Found: C, 67.01; H, 6.23; N, 4.68.

Example 453

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-benzothiazolyl)amino)ethyl)pyrrolidine-3-carboxylic acid

The title compound was prepared by the method of Example 418, substituting 2-chlorobenzothiazole for 2-bromopyrimidine. ^1H NMR (300 MHz, CDCl_3) δ 0.88 (t, $J=7$ Hz, 3H), 1.59 (sextet, $J=7$ Hz, 2H), 2.25-2.37 (m, 1H), 2.85-

2.97 (m, 3H), 3.28-3.36 (m, 2H), 3.50-3.58 (m, 3H), 3.60-3.65 (m, 1H), 3.67 (d, J=9Hz, 1H), 3.71 (s, 3H), 5.87 (d, J=2Hz, 1H), 5.91 (d, J=2Hz, 1H), 6.57 (d, J=8Hz, 1H), 6.73 (dd, J=2Hz, 9Hz, 1H), 6.76 (d, J=8 Hz, 2H), 6.91 (d, J=2Hz, 1H), 7.01 (t, J=8Hz, 1H), 7.22 (t, J=8Hz, 1H), 7.29 (d, J=8Hz, 2H), 7.40 (d, J=7Hz, 1H), 7.55 (d, J=7Hz, 1H).

Example 454

trans,trans-2-(2-Ethoxyethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CDCl₃, 300 MHz) δ 0.91 (t, J = 7.4 Hz, 3H), 0.94 (t, J = 7.4 Hz, 3H), 1.19 (t, J = 7.0 Hz, 3H), 1.24-1.38 (m, 5H), 1.46-1.60 (m, 4H), 2.03-2.12 (m, 2H), 3.07 (t, J = 8.0 Hz, 1H), 3.07-3.34 (m, 6H), 3.43-3.52 (m, 3H), 3.59-3.74 (m, 3H), 3.80-4.01 (m, 2H), 5.93 (s, 2H), 6.72 (d, J = 8.1 Hz, 1H), 6.79 (dd, J = 8.2 Hz, 1.7 Hz, 1H), 6.87 (d, J = 1.7 Hz, 1H). MS(DCl/NH₃) m/e 477 (M+H)⁺. Anal calcd for C₂₆H₄₀N₂O₆ · 0.4 TFA: C, 61.64; H, 7.80; N, 5.36. Found: C, 61.63; H, 7.84; N, 5.29.

Example 455

trans,trans-2-(4-Methoxy-3-fluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-(morpholin-4-ylethyl)sulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Ethyl 2-(4-methoxy-3-fluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-vinylsulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid, prepared by the procedures of Example 125, was reacted with excess morpholine for 4 hours at room temperature. Chromatography on silica gel eluting with EtOAc gave a 65% yield of an intermediate ethyl ester which was hydrolyzed to the title compound with NaOH in ethanol/water. ¹H NMR (300 MHz, CDCl₃) δ 0.81 (t, J=7Hz, 3H), 1.46 (sextet, J=7Hz, 2H), 2.43-2.52 (m, 4H), 2.70-2.92 (m, 5H), 2.97-3.33 (m, 6H), 3.60 (dd, J=3Hz, 9Hz, 1H), 3.51-3.59 (m, 1H), 3.62-3.70 (m, 5H), 3.88 (s, 3H), 5.95 (s, 2H), 6.72 (d, J=8Hz, 1H), 6.70 (dd, J=2Hz, 8Hz, 1H), 6.90 (t, J=9Hz, 1H), 6.96 (d, J=2Hz, 1H), 7.10 (d, J=8Hz, 1H), 7.18 (dd, J=2Hz, 12Hz, 1H).

Example 456

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-((2,2,2-trifluoroethoxyethane)sulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 95-96 °C. ¹H NMR (CD₃OD, 300MHz) δ 0.80 (t, J=7Hz, 3H), 1.35-1.48 (m, 2H), 3.07 (sextet, J=7Hz, 2H), 3.23-3.55 (m, 8H), 3.80-3.87 (m, 2H), 3.93 (s, 3H), 3.94-4.02 (m, 4H), 4.66 (d, J=12Hz, 1H), 5.96 (s, 2H), 6.83 (d, J=8Hz, 1H), 6.94 (d, J=8Hz, 1H), 7.06 (d, J=2Hz, 1H), 7.23 (t, J=9Hz, 1H), 7.43 (d, J=9Hz, 1H), 7.49 (dd, J=2Hz, J=12Hz, 1H). MS (DCI/NH₃) m/e 635 (M+H)⁺.

Example 457

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-fluorophenyl)-1-(N-butyl-N-(3-methylphenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.20-1.50 (m, 4H), 2.31 (s, 3H), 2.65-2.80 (m, 2H), 3.19 (t, J=7 Hz, 1H), 3.25 (d, J=10 Hz, 1H), 3.35-3.65 (m, 4H), 3.79 (d, J=10 Hz, 1H), 5.93 (s, 2H), 6.74 (d, J=7 Hz, 1H), 6.80-6.90 (m, 3H), 6.91-7.09 (m, 3H), 7.10-7.35 (m, 4H). MS (DCI) m/e 533 (M+H)⁺. Anal calcd for C₃₁H₃₃N₂O₅F: C, 69.91; H, 6.25; N, 5.26. Found: C, 69.56; H, 6.26; N, 5.23.

Example 458

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(butanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.94 (m, 3H), 1.23 (hex, 2H, J=8), 1.69 (m, 2H), 3.08 (m, 2H), 3.20 (s, 3H), 3.3-3.5 (m, 10H), 3.77 (m, 2H), 3.92 (s, 3H), 4.60 (m, 1H), 5.96 (s, 2H), 6.81 (d, 1H, J=8), 6.88 (dd, 1H, J=2, 8), 6.99 (d, 1H, J=2), 7.22 (t, 1H, J=9), 7.38 (m, 2H). MS (APCI) m/e 581 (M+H)⁺. Anal calcd for C₂₈H₃₇N₂O₈F₅ · 1.1 TFA: C, 51.37; H, 5.44; N, 3.97. Found: C, 51.27; H, 5.35; N, 4.11.

Example 459

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-methylpropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared and isolated as a white solid. m.p. 77-78 °C. ¹H NMR (CDCl₃,

300MHz) δ 0.83 (t, J=7Hz, 3H), 1.06 (d, J=6Hz, 6H), 1.45 (q, J=7Hz, 2H), 2.20 (septet, J=6Hz, 1H), 2.26-2.36 (m, 1H), 2.62-2.78 (m, 3H), 2.85-2.95 (m, 2H), 2.97-3.10 (m, 2H), 3.15-3.35 (m, 2H), 3.43 (dd, J=3Hz, J=9Hz, 1H), 3.53-3.62 (m, 1H), 3.66 (d, J=9Hz, 1H), 3.88 (s, 3H), 5.95 (s, 2H), 6.74 (d, J=8Hz, 1H), 6.82 (dd, J=2Hz, J=8Hz, 1H), 6.92 (t, J=8Hz, 1H), 6.97 (d, J=2Hz, 1H), 7.12 (d, J=9Hz, 1H), 7.18 (dd, J=2Hz, J=12Hz, 1H). MS (DCI/NH₃) m/e 565 (M+H)⁺.

Example 460

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(4-nitrobenzyl)amino)carbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ (rotamer) 8.11 (2H, m), 7.32 (3H, dd, J=9, 2), 7.16 (7.07) (1H, bd, J=10), 6.98 (6.94) (1H, d, J=2), 6.85 (2H, d, J=9), 6.83-6.70 (2H, m), 5.99 (5.97) (2H, d, J=2), 5.02 (4.18) (1H, d, J=15), 4.63 (4.38) (1H, d, J=15), 3.79 (3.77) (3H, s), 3.72 (1H, d, J=10), 3.61 (1H, m), 3.48 (1H, bd, J=15), 3.43-3.20 (2H, m), 3.06 (2H, m), 2.90 (1H, m), 3.79 (1H, bd, J=14), 1.43 (1H, m), 1.23 (2H, m), 1.02 (1H, m), 0.84 (0.78) (3H, t, J=8). MS (DCI/NH₃) m/e 590 (M+H)⁺. Anal calcd for C₃₂H₃₅N₃O₈: C, 65.18; H, 5.98; N, 7.13. Found: C, 65.89; H, 5.85; N, 6.85.

Example 461

trans,trans-2-(4-Ethylphenyl)-4-(3,4-difluorophenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.78 (t, 3H, J=7), 0.87 (t, 3H, J=7), 1.02 (hex, 2H, J=7), 1.22 (t, 3H, J=7), 1.27 (m, 2H), 1.45 (m, 2H, J=7), 2.63 (q, 2H, J=7), 2.77 (d, 1H, J=14), 2.94 (dd, 1H, J=7, 9), 3.05 (m, 3H), 3.3-3.5 m, 3H), 3.44 (d, 1H, J=14), 3.66 (m, 1H), 3.75 (d, 1H, J=10), 7.20 (td, 2H, J=1,8), 7.22 (m, 2H), 7.32 (td, 2H, J=1,8), 7.43 (ddd, 1H, J=2,8,12). MS (DCI/NH₃) m/e 501 (M+H)⁺. Anal calcd for C₂₉H₃₈N₂O₃F₂ · 0.6 H₂O: C, 68.11; H, 7.73; N, 5.48. Found: C, 68.03; H, 7.53; N, 5.37.

Example 462

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.20-1.50 (m, 4H), 2.21 (d, J=2 Hz, 3H), 2.64 (d, J=14 Hz, 1H), 2.75 (dd, J=10 Hz, 1H), 3.05 (t, J=7 Hz, 1H), 3.25 (d, J=15 Hz, 1H), 3.35-3.70 (m, 5H), 3.77 (s, 3H), 5.92 (s, 2H), 6.70-6.92 (m, 6H), 6.96-7.10 (m, 4H). MS (DCI) m/e 563 (M+H)⁺. Anal calcd for C₃₂H₃₅N₂O₆F · 0.5 H₂O: C, 67.24; H, 6.35; N, 4.90. Found: C, 67.16; H, 6.06; N, 4.81.

Example 463

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-((3-isopropyl)phenyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, 3H), 1.17 (d, J=7 Hz, 6H), 1.20-1.50 (m, 4H), 2.63 (d, J=15 Hz, 1H), 2.75 (t, J=7 Hz, 1H), 2.85 (m, 1H), 3.00 (t, J=7 Hz, 1H), 3.25 (d, J=15 Hz, 1H), 3.40-3.70 (m, 5H), 3.75 (s, 3H), 5.90 (s, 2H), 6.65-6.80 (m, 3H), 6.71 (dt, J=7 Hz, 3H), 7.07 (m, 3H), 7.20-7.35 (m, 2H). MS (DCI) m/e 573 (M+H)⁺. Anal calcd for C₃₄H₄₀N₂O₆ · 0.15 H₃PO₄: C, 69.52; H, 6.94; N, 4.77. Found: C, 63.31; H, 6.72; N, 4.43.

Example 464

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N-butyl-N-(3-ethylphenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (m, J=7 Hz, 3H), 1.16 (t, J=7 Hz, 3H), 1.20-1.47 (m, 4H), 2.50 (q, J=7 Hz, 2H), 2.70-2.85 (m, 2H), 3.13 (t, J=7 Hz, 1H), 3.20-4.5 (m, 6H), 3.78 (s, 3H), 3.83 (d, J=8 Hz, 1H), 5.92 (s, 2H), 6.72 (d, J=8 Hz, 1H), 6.80-6.90 (m, 5H), 7.02-7.13 (m, 3H), 7.15-7.25 (m, 2H). MS (DCI) m/e 559 (M+H)⁺. Anal calcd for C₃₃H₃₈N₂O₆ · 0.3 H₂O: C, 70.27; H, 6.90; N, 4.97. Found: C, 70.31; H, 6.63; N, 4.60.

Example 465

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-ethylphenyl)-1-(((N-(3-chlorophenyl)-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 0.87 (t, 3H, $J=7\text{Hz}$), 1.23 (t, 3H, $J=7\text{Hz}$), 1.28 (m, 2H), 1.41 (m, 2H), 2.63 (q, 2H, $J=7\text{Hz}$), 2.67 (m, 1H), 2.92 (m, 1H), 3.20 (m, 2H), 3.42 (m, 1H), 3.60 (q, 2H, $J=7\text{Hz}$), 3.93 (m, 1H), 5.92 (s, 2H), 6.75 (d, 1H, $J=8\text{Hz}$), 6.84 (m, 3H), 6.95 (br s, 1H), 7.02 (s, 1H), 7.10 (br s, 3H), 7.25 (m, 2H). MS (APCI) m/e 563 ($\text{M}+\text{H}$) $^+$. Anal. calc'd for $\text{C}_{32}\text{H}_{35}\text{N}_2\text{O}_5\text{Cl} \cdot 0.80 \text{H}_3\text{PO}_4$: C, 59.92; H, 5.88; N, 4.37. Found: C, 59.90; H, 5.83; N, 4.07.

Example 466

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-ethylphenyl)-1-((N-(3-chlorophenyl)-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 0.86 (t, 3H, $J=7\text{Hz}$), 1.23 (t, 3H, $J=7\text{Hz}$), 1.25 (m, 2H), 1.40 (m, 2H), 2.64 (q, 2H, $J=7\text{Hz}$), 2.70 (m, 1H), 2.95 (m, 1H), 3.20 (m, 2H), 3.40 (m, 1H), 3.57 (m, 3H), 3.90 (m, 1H), 4.25 (s, 4H), 6.80 (d, 1H, $J=8\text{Hz}$), 6.95 (d, 1H, $J=2\text{Hz}$), 6.95 (m, 2H), 7.07 (br s, 3H), 7.22 (m, 3H). MS (APCI) m/e 577. ($\text{M}+\text{H}$) $^+$. Anal. calc'd for $\text{C}_{33}\text{H}_{37}\text{N}_2\text{O}_5\text{Cl} \cdot 0.85 \text{H}_2\text{O}$: C, 66.90; H, 6.58; N, 4.73. Found: C, 66.92; H, 6.25; N, 4.36.

Example 467

trans,trans-4-(Benzofuran-5-yl)-2-(4-ethylphenyl)-1-((N-(3-chlorophenyl)-N-butylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 0.85 (t, 3H, $J=7\text{Hz}$), 1.26 (t, 3H, $J=7\text{Hz}$), 1.30 (m, 2H), 1.40 (m, 2H), 2.60 (q, 2H, $J=7\text{Hz}$), 2.72 (m, 1H), 2.93 (m, 1H), 3.22 (m, 2H), 3.50 (m, 1H), 3.55 (m, 2H), 3.75 (m, 1H), 3.90 (br d, 1H), 6.75 (d, 1H, $J=1\text{Hz}$), 6.80 (br d, 1H), 6.95 (br s, 1H), 7.08 (m, 4H), 7.20 (t, 1H, $J=8\text{Hz}$), 7.28 (t, 1H, $J=8\text{Hz}$), 7.42 (m, 2H), 7.58 (d, 1H, $J=1\text{Hz}$), 7.63 (s, 1H). MS (APCI) m/e 559 ($\text{M}+\text{H}$) $^+$. Anal. calc'd for $\text{C}_{33}\text{H}_{35}\text{N}_2\text{O}_4\text{Cl} \cdot 0.45 \text{H}_2\text{O}$: C, 69.88; H, 6.38; N, 4.94. Found: C, 69.83; H, 6.04; N, 4.87.

Example 468

trans,trans-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-butyl-N-phenylamino)ethyl)pyrrolidine-3-carboxylic acid

Ethyl 2-(4-methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(bromoethyl)-pyrrolidine-3-carboxylate, prepared using the procedures

of Example 61A (300 mg), was reacted with N-butyl aniline (190 mg) in 1 mL of dioxane containing 130 mg of diisopropylethylamine to give the ethyl ester. The ester was hydrolyzed with sodium hydroxide to give 148 mg of the title compound as a white powder. ¹H NMR (300 MHz, CDCl₃) δ 0.90 (t, J=9Hz, 3H), 1.28 (sextet, J=7Hz, 2H), 1.46 (quintet, J=7Hz, 2H), 2.20-2.32 (m, 1H), 2.68-2.77 (m, 1H), 2.82-2.95 (m, 2H), 3.12-3.22 (m, 2H), 3.30-3.44 (m, 3H), 3.45-3.55 (m, 1H), 3.62 (d, J=9Hz, 1H), 3.83 (s, 3H), 3.90 (s, 3H), 5.95 (s, 2H), 6.51 (d, J=7Hz, 2H), 6.55-6.62 (m, 2H), 6.69 (d, J=2Hz, 1H), 6.84 (t, J=8Hz, 1H), 7.02-7.15 (m, 3H), 7.19 (dd, J=2Hz, 12Hz, 1H).

Example 469

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-ethylphenyl)-1-((N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 0.78 (t, 3H, J=7Hz), 0.88 (t, 3H, J=7Hz), 1.05 (q, 2H, J=7Hz), 1.23 (t, 3H, J=7Hz), 1.28 (m, 2H), 1.45 (m, 2H), 2.64 (q, 2H, J=7Hz), 2.78 (m, 1H), 2.9-3.2 (envelope, 4H), 3.30 (m, 1H), 3.40 (m, 3H), 3.60 (m, 1H), 3.80 (m, 1H), 4.25 (s, 4H), 6.80 (d, 1H, J=8Hz), 6.90 (m, 1H), 6.98 (d, 1H, J=2Hz), 7.17 (d, 2H, J=8Hz), 7.30 (m, 2H). MS (APCI) m/e 523 (M+H)⁺. Anal. calc'd for C₃₁H₄₂N₂O₅ · 1.1 HOAc: C, 67.73; H, 7.94; N, 4.76. Found: C, 67.81; H, 7.55; N, 4.48.

Example 470

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(3-methylphenylamino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7.1 Hz, 3H), 1.30 (m, 2H), 1.44 (m, 2H), 2.30 (s, 3H), 2.80 (d, J=15.2 Hz, 1H), 2.85 (t, J=9.3 Hz, 1H), 3.19 (t, J=9.3 Hz, 1H), 3.33 (d, J=10.2 Hz, 1H), 3.42-3.61 (m, 3H), 3.79 (s, 3H), 3.91 (d, J=9.8 Hz, 1H), 4.22 (m, 4H), 6.75-6.86 (m, 6H), 6.95 (d, J=2.0 Hz, 1H), 7.09 (d, J=8.8 Hz, 2H), 7.22 (d, J=10.2 Hz, 1H), 7.26 (t, J=7.6 Hz, 1H). MS (DCI) m/e 559 (M+H)⁺. Anal. calc'd for C₃₃H₃₈N₂O₆ · 0.4 CH₃CO₂C₂H₅: C, 69.97; H, 6.99; N, 4.72. Found: C, 0.06; H, 6.66; N, 4.48.

Example 471

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(3-chlorophenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7.0 Hz, 3H), 1.25 (m, 2H), 1.40 (m, 2H), 2.78 (d, J=14.6 Hz, 1H), 2.86 (t, J=9.0 Hz, 1H), 3.16 (t, J=9.5 Hz, 1H), 3.34-3.43 (m, 2H), 3.48-3.62 (m, 3H), 3.79 (s, 3H), 3.85 (d, J=9.5 Hz, 1H), 4.22 (m, 4H), 6.78 (d, J=8.5 Hz, 1H), 6.81-6.86 (m, 3H), 6.93-7.09 (m, 5H), 7.33-7.38 (m, 2H). MS (DCI) m/e 579 (M+H⁺). Anal calcd for C₃₂H₃₅ClN₂O₆ · 1.1 CH₃CO₂C₂H₅ · 0.15 H₃PO₄: C, 63.30; H, 6.46; N, 4.06. Found: C, 63.54; H, 6.09; N, 3.98.

Example 472

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(4-pyridylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 2.84 (t, J=9.6 Hz, 1H), 2.88 (dd, J=9.6, 7.3 Hz, 1H), 3.09 (dd, J=3.3, 9.6 Hz, 1H), 3.21 (d, J=14.3 Hz, 1H), 3.53 (m, 1H), 3.78 (s, 3H), 3.81 (m, 2H), 5.92 (m, 2H), 6.73 (d, J=8.1 Hz, 1H), 6.82 (dd, J=1.8, 8.1 Hz, 1H), 6.93 (m, 2H), 6.95 (d, J=1.5 Hz, 1H), 7.43 (m, 4H), 8.44 (d, J=5.2 Hz, 2H). MS (DCI) m/e 433 (M+H⁺). Anal calcd for C₂₅H₂₄N₂O₅ · 0.3 CH₃CO₂C₂H₅: C, 68.57; H, 5.80; N, 6.10. Found: C, 68.68; H, 5.60; N, 5.81.

Example 473

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(3-*tert*-butylphenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.88 (t, J=7.2 Hz, 3H), 1.23 (s, 9H), 1.26-1.45 (m, 4H), 2.74 (dd, J=15.1 Hz, 1H), 2.84 (m, 1H), 3.13 (t, J=9.0 Hz, 1H), 3.29 (d, J=15.1 Hz, 1H), 3.50-3.66 (m, 4H), 3.77 (s, 3H), 3.84 (d, J=9.6 Hz, 1H), 5.92 (s, 2H), 6.74 (d, J=7.7 Hz, 1H), 6.79-6.85 (m, 4H), 6.86-6.90 (m, 1H), 6.99 (t, J=1.8 Hz, 1H), 7.06 (d, J=1.8 Hz, 1H), 7.13 (m, 2H), 7.33 (t, J=7.7 Hz, 1H), 7.42 (m, 1H). MS (DCI) m/e 587 (M+H⁺). Anal calcd for C₃₅H₄₂N₂O₆: C, 71.65; H, 7.22; N, 4.77. Found: C, 71.56; H, 7.33; N, 4.69.

Example 474

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(3-n-butylphenylamino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.88 (t, J=7.3 Hz, 3H), 0.92 (t, J=7.3 Hz, 3H), 1.23-1.59 (m, 8H), 2.58 (t, J=7.6 Hz, 2H), 2.75 (d, J=15.3 Hz, 1H), 2.80 (dd, J=8.5, 9.5 Hz, 1H), 3.12 (t, J=9.3 Hz, 1H), 3.29 (d, J=15.6 Hz, 1H), 3.46 (dd, J=4.9, 9.7 Hz, 1H), 3.52-3.64 (m, 3H), 3.78 (s, 3H), 3.83 (d, J=9.8 Hz, 1H), 5.92 (s, 2H), 6.74 (d, J=8.1 Hz, 1H), 6.79-6.87 (m, 4H), 7.05 (d, J=1.7 Hz, 1H), 7.10 (d, J=8.8 Hz, 2H), 7.20 (d, 7.8H), 7.29 (t, J=7.6 Hz, 1H). MS (DCI) m/e 587 (M+H)⁺. Anal calcd for C₃₅H₄₂N₂O₆: C, 71.65; H, 7.22; N, 4.77. Found: C, 71.33; H, 7.28; N, 4.74.

Example 475

trans,trans-4-(3,4-Difluorophenyl)-2-(4-ethylphenyl)-1-(N-(n-butyl)-N-(3-methylphenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.87 (t, 3H, J=7), 1.19 (t, 3H, J=7), 1.28 (m, 2H), 1.43 (m, 2H), 2.28 (s, 3H), 2.60 (q, 2H, J=7), 2.66 (m, 2H), 3.06 (m, 1H), 3.21 (d, 1H, J=15), 3.42 (dd, 1H, J=4,9), 3.58 (m, 3H), 3.71 (d, 1H, J=9), 6.80 (s, 2H), 7.06 (s, 4H), 7.18 (m, 4H), 7.45 (m, 1H). MS (APCI) m/e 535 (M+H)⁺. Anal calcd for C₃₂H₃₆N₂O₃F₂ · 1.3 HOAc: C, 67.83; H, 6.78; N, 4.57. Found: C, 67.83; H, 6.46; N, 4.70.

Example 476

trans,trans-2-(4-Ethylphenyl)-4-(3,4-difluorophenyl)-1-(N-(n-butyl)-N-(3-chlorophenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.82 (t, 3H, J=7), 1.16 (t, 3H, J=7), 1.23 (m, 2H), 1.35 (m, 2H), 2.55 (q, 2H, J=7), 2.66 (m, 2H), 3.01 (t, 1H, J=9), 3.16 (d, 1H, J=15), 3.32 (dd, 1H, J=4,9), 3.56 (m, 3H), 3.67 (d, 1H, J=9), 6.94 (d, 1H, J=7), 7.02 (m, 5H), 7.14 (m, 2H), 7.32 (m, 3H). MS (APCI) m/e 555 (M+H)⁺. Anal calcd for C₃₁H₃₃N₂O₃ClF₂ · 0.6 TFA: C, 61.88; H, 5.42; N, 4.48. Found: C, 61.90; H, 5.62; N, 3.98.

Example 477

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-fluorophenyl)-1-(N-butyl-N-(3-chlorophenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7 Hz, 3H), 1.10-1.30 (m, 4H), 2.60-2.75 (m, 2H), 3.03 (t, J=7 Hz, 1H), 3.15-3.75 (m, 6H), 4.02 (m, 4H), 6.75 (d, J=6 Hz, 1H), 6.85 (dd, J=7 Hz, 1H), 6.90 (7.19, J=m Hz, 6H), 7.32-7.43 (m, 3H). MS (DCI) m/e 567 (M+H)⁺. Anal calcd for C₃₁H₃₂N₂O₅Cl · 1.6 H₂O: C, 62.49; H, 5.95; N, 4.70. Found: C, 62.20; H, 5.54; N, 4.42.

Example 478

trans,trans-4-(Benzofuran-5-yl)-2-(4-ethylphenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CDCl₃) δ 0.78 (t, 3H, J=7Hz), 0.84 (t, 3H, J=7Hz), 1.05 (q, 2H, J=7Hz), 1.21 (t, 3H, J=7Hz), 1.25 (m, 2H), 1.45 (m, 2H), 2.62 (q, 2H, J=7Hz), 2.80 (d, 1H, J=13Hz), 3.0 (m, 2H), 3.15 (m, 2H), 3.35 (m, 1H), 3.43 (m, 2H), 3.52 (m, 1H), 4.40 (m, 2H), 6.73 (d, 1H, J=1Hz), 7.14 (d, 2H, J=8Hz), 7.26 (s, 1H), 7.31 (d, 2H, J=8Hz), 7.44 (s, 2H), 7.60 (d, 1H, J=1Hz), 7.65 (s, 1H). MS (APCI) m/e 505 (M+H)⁺. Anal. calc'd for C₃₁H₄₀N₂O₄: C, 73.78; H, 7.99; N, 5.55. Found: C, 73.69; H, 7.97; N, 5.21.

Example 479

trans,trans-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(pyrrolidine-1-carbonylmethyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Ethyl 2-(4-methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-propyl-aminoethyl)-pyrrolidine-3-carboxylate, prepared according to the procedures of Example 61B (300 mg), N-bromoacetyl pyrrolidine (132 mg) and diisopropylethylamine (154 mg) were heated for 1 hour at 50 °C in 1 mL of acetonitrile to give the intermediate ethyl ester. The ester was hydrolyzed to the title compound by the method of Example 1D. ¹H NMR (300 MHz, CDCl₃) δ 0.88 (t, J=7Hz, 3H), 1.30-1.45 (m, 2H), 1.75-1.92 (m, 4H), 2.30-2.40 (m, 1H), 2.47-2.58 (m, 2H), 2.70-3.00 (m, 5H), 3.24-3.45 (m, 6H), 3.50-3.70 (m, 2H), 3.83 (s, 3H), 3.86 (d, J=9Hz, 1H), 3.88 (s, 3H), 5.93 (s, 2H), 6.58 (d, J=2Hz, 1H), 6.70 (d, J=2Hz, 1H), 6.87 (t, J=8Hz, 1H), 7.10 (d, J=9Hz, 1H), 7.21 (dd, J=2Hz, 12Hz, 1H).

Example 480

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-((N-(perhydroazepinylcarbonyl)-(D)-leucyl)amino)ethyl)pyrrolidine-3-carboxylic acid

Example 480AD-Leucine O-benzyl ester Tosylate salt

To benzyl alcohol (8.2 g) dissolved in benzene (30 mL) was added D-leucine (5.0 g) and *p*-toluenesulfonic acid monohydrate (8.0 g). The reaction was warmed to reflux with removal of water overnight. Once TLC indicated consumption of starting material, the reaction was cooled, and the resulting solid was filtered and washed with EtOAc to give the title compound as a white powder (14.26 g, 99%).

Example 480BN-Perhydroazepinylcarbonyl-D-Leucine O-Benzyl ester

To the compound resulting from Example 480A (1.0 g) dissolved in chloroform (20 mL) was added triethylamine (0.4 mL). The solution was cooled to 0 °C, and carbonyldiimidazole was added. After 1.5 hours, TLC indicated complete consumption of starting material, so hexamethylene imine (0.327 mL) was added. After 1 hour, an additional amount of hexamethylene imine (0.330 mL) was added, and the reaction was stirred at ambient temperature overnight. The solution was washed with sodium bicarbonate (2 x 20 mL), 1 N H₃PO₄ (2 x 20 mL), and brine (20 mL), dried over Na₂SO₄, decanted and evaporated. The residue was purified by flash chromatography on silica gel eluting with 25 - 50% EtOAc in hexanes to give the title compound as a crystalline solid (0.835 g, 89%).

Example 480CN-Perhydroazepinylcarbonyl-D-Leucine

To the compound resulting from Example 480B (200 mg) dissolved in dry ethanol (1.0 mL) was added 10% palladium on carbon (10 mg). After flushing the flask with nitrogen, the reaction was stirred vigorously under an atmosphere of hydrogen for 1 hour. The reaction was filtered through infusorial earth and evaporated to give the title compound (140 mg).

Example 480D*trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(cyanomethyl)-pyrrolidine-3-carboxylic acid ethyl ester*

To the compound resulting from Example 1C (510 mg of a 50 % wt. solution in toluene) dissolved in acetonitrile (2.0 mL) was added diisopropylethylamine (0.24 mL), followed by bromoacetonitrile (0.072 mL). After 2 hours, TLC indicated complete consumption of starting material. The solvent was evaporated, and the residue was purified by flash chromatography on silica gel eluting with 20 - 40% EtOAc in hexanes to give the title compound as a colorless oil (0.28 g, 99%).

Example 480E*trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-aminoethyl)-pyrrolidine-3-carboxylic acid ethyl ester*

To the compound resulting from Example 480D (275 mg) dissolved in 10 mL each of triethylamine and ethanol was added Raney nickel catalyst (0.2 g), and the reaction was placed under a hydrogen atmosphere (4 atmospheres) for 3 days. The reaction was filtered and evaporated. The residue was dissolved in methylene chloride (10 mL) and extracted with 1 M HCl (5 x 1 mL). The combined aqueous extracts were basified and then extracted with methylene chloride (5 x 2 mL). The combined organic extracts were dried with MgSO₄, filtered and evaporated to give the title compound as an unstable oil (0.14 g).

Example 480F*trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-((N-(perhydroazepinylcarbonyl)leucyl)amino)ethyl)-pyrrolidine-3-carboxylic acid, ethyl ester*

The compound resulting from Example 480E (0.10 g) was dissolved in methylene chloride (3.0 mL), and the compound resulting from Example 480C (0.07 g) was added. The solution was cooled to 0 °C, and EDCI (0.052 g) was added. After 4 hours, the reaction was evaporated and partitioned between water (1 mL), and EtOAc (10 mL). The organic solution was washed with water (1 mL) and brine (1 mL), dried over MgSO₄, filtered and evaporated. The residue was purified by flash chromatography on silica gel eluting with 50 - 60% EtOAc in hexanes to give the title compound as a colorless oil (0.075 g, 48%).

Example 480G

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-((N-(perhydroazepinylcarbonyl)leucyl)amino)ethyl)pyrrolidine-3-carboxylic acid

The compound resulting from Example 480F (0.75 g) was dissolved in ethanol (1.0 mL) and 5 M NaOH (0.050 mL) was added. After 2 hours, additional 5 M NaOH (0.090 mL) was added. After an additional 3.5 hours, the reaction was evaporated. The residue was dissolved in water (5 mL) and washed with diethyl ether (2 x 2 mL). The aqueous solution was acidified with 1 N H₃PO₄ to pH • 3. The solid which precipitated dissolved when the mixture was extracted with chloroform (3 x 3 mL). The chloroform extracts were washed with brine (2 mL), dried with MgSO₄, filtered and evaporated to give the title compound as a tan solid (0.053 g). Purification by HPLC (Vydac mC18) eluting with a 10 - 70% gradient of CH₃CN in 0.1%TFA provided suitable material (0.049 g) after lyophilization of the desired fractions. ¹H NMR (CDCl₃, 300 MHz) δ 0.82 (dd, 6.4, 4.4 Hz, 6H), 0.87 (dd, J = 5.7, 5.7 Hz, 6H), 1.04-1.28 (m, 3H), 1.34-1.65 (m, 19H), 2.95 (br m, 2H), 3.15-3.40 (m, 14H), 3.40-3.55 (m, 4H), 3.58-3.68 (m, 2H), 3.70-3.76 (br m, 2H), 3.80 (s, 3H), 3.81 (s, 3H), 4.15 (br m, 2H), 5.10 (br m, 2H), 5.93 (s, 3H), 5.95 (s, 3H), 6.70-6.97 (m, 13H), 7.43-7.56 (br m, 3H), 8.2 (br s, 1H), 8.5 (br s, 1H). MS(DCI/NH₃) m/e 623 (M+H)⁺. Anal calcd for C₃₄H₄₆N₄O₇ • 2.00 TFA: C, 53.65; H, 5.69; N, 6.58. Found: C, 53.66; H, 5.66; N, 6.54.

Example 481

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(N,N-di(n-hexyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.80-0.95 (m, 6H), 1.0 (m, 2H), 1.07 (1.55, J=m Hz, 14H), 2.70 (d, J=13 Hz, 1H), 2.85-3.15 (m, 4H), 3.20-3.60 (m, 9H), 3.64 (d, J=10 Hz, 1H), 3.79 (s, 3H), 5.90 (m, 2H), 6.70 (d, 8H), 1, 6.80-6.93 (m, 3H), 7.05 (2, 1H), 7.35 (d, J=10 Hz, 2H). Anal calcd for C₃₃H₄₆N₂O₆ • 1.7 H₂O: C, 66.35; H, 8.34; N, 4.69. Found: C, 66.32; H, 8.04; N, 4.52.

Example 482

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-fluorophenyl)-1-(N-butyl-N-(3-methylphenyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CD_3OD) δ 0.87 (t, $J=7$ Hz, 3H), 1.20-1.35 (m, 2H), 1.35-1.40 (m, 2H), 2.32 (s, 3H), 2.55-2.70 (m, 2H), 2.97 (t, $J=7$ Hz, 1H), 3.22 (d, $J=14$ Hz, 1H), 3.25-3.70 (m, 5H), 4.20 (m, 4H), 6.97 (d, $J=2$ Hz, 1H), 7.09 (m, 2H), 7.15-7.35 (m, 2H). MS (DCI) m/e 547 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{32}\text{H}_{35}\text{N}_2\text{O}_5\text{F} \cdot 1.2 \text{H}_2\text{O}$: C, 67.64; H, 6.63; N, 4.93. Found: C, 67.73; H, 6.37; N, 4.70.

Example 483

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(3-nitrobenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ (rotamer) 8.14 (2H, m), 8.05 (7.83) (1H, m), 7.60-7.30 (3H, m), 7.13 (1H, m), 7.10-6.70 (5H, m), 5.94 (2H, m), 5.43 (5.33) (1H, d, $J=12$), 4.75 (1H, bd, $J=15$), 4.60-4.20 (2H, m), 4.10 (2H, m), 3.80 (3.76) (3H, s), 3.75-3.40 (3H, m), 3.20-2.80 (2H, m), 1.50 (1H, m), 1.30 (1H, m), 1.20-1.00 (2H, m), 0.91 (0.78) (3H, t, $J=8$). MS (DCI/ NH_3) m/e 590 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{32}\text{H}_{35}\text{N}_3\text{O}_8 \cdot 2.1 \text{ TFA}$: C, 52.44; H, 4.51; N, 5.07. Found: C, 52.25; H, 4.83; N, 5.71.

Example 484

trans,trans-4-(1,2-Dihydrobenzofuran-5-yl)-2-(4-ethylphenyl)-1-(((N-butyl-N-(3,4-dimethoxybenzyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H (300MHz, CDCl_3) δ (rotamer) 7.40 (2H, m), 7.30-7.10 (4H, m), 6.90-6.70 (3H, m), 6.48 (1H, m), 5.45 (1H, m), 4.65 (1H, d, $J=15$), 4.57 (2H, dt, $J=9, 3$), 4.40-4.00 (5H, m), 3.87 (3.85) (3H, s), 3.84 (1H, m), 3.83 (3.79) (3H, s), 3.56 (2H, m), 3.20 (2H, t, $J=10$), 2.90 (1H, m), 2.64 (2H, q, $J=8$), 1.52 (1H, m), 1.31 (2H, m), 1.22 (3H, dt, $J=9, 2$), 1.07 (1H, m), 0.92 (0.78) (3H, t, $J=8$). MS (DCI/ NH_3) m/e 601 ($\text{M}+\text{H}^+$). Anal calcd for $\text{C}_{36}\text{H}_{44}\text{N}_2\text{O}_6 \cdot 1.35 \text{ TFA}$: C, 61.59; H, 6.06; N, 3.71. Found: C, 61.69; H, 6.04; N, 3.63.

Example 485

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-(((N-butyl-N-(4-heptyl)amino)carbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CD_3OD) δ 0.71-1.04 (m, 11H), 1.07-1.35 (m, 6H),

1.73-1.53 (m, 4H), 2.79-3.25 (m, 5H), 3.35-3.44 (m, 1H), 3.51-3.68 (m, 3H), 3.78-3.89 (m, 1H), 3.79 (s, 3H), 5.92 (m, 2H), 6.74 (dd, J=1.7, 8.1 Hz, 1H), 6.85 (td, J=1.7, 8.1 Hz, 1H), 6.93 (m, 2H), 7.02 (dd, J=1.7, 9.5 Hz, 1H), 7.36 (m, 2H). MS (C.I.) m/e 553 (M+H⁺). Anal calcd for C₃₂H₄₄N₂O₆: C, 69.54; H, 8.02; N, 5.07.

5 Found: C, 69.31; H, 7.89; N, 5.06.

Example 486

trans,trans-2-(4-Methylcyclohexyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

10 Using the procedures described in Example 1, the title compound was prepared and isolated as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.88 (3H, d, J = 7Hz), 0.92 (3H, t, J = 7Hz), 0.96 (3H, t, J = 7Hz), 1.05 (1H, m), 1.22-1.40 (7H, m), 1.45-1.65 (6H, m), 1.67-1.84 (4H, m), 3.17-3.45 (6H, m), 3.70 (1H, brn), 3.82 (1H, dd, J = 9Hz, 15Hz), 3.86 (1H, d, J = 15Hz), 5.93 (2H, s), 6.73 (1H, d, J = 8Hz), 6.78 (1H, dd, J = 2Hz, 8Hz), 6.88 (1H, d, J = 2Hz). MS (DCI/NH₃) m/e 501 (M+H⁺). Anal calcd for C₂₉H₄₄N₂O₅ · 0.25 CF₃CO₂H: C, 66.9%; H, 8.43; N, 5.29. Found: C, 66.79; H, 8.60; N, 4.87.

Example 487

trans,trans-2-(2-Propylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

20 Using the procedures described in Example 1, the title compound was prepared and isolated as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (6H, m), 0.92 (3H, t, J = 7Hz), 0.97 (3H, t, J = 7Hz), 1.12-1.40 (13H, m), 1.42-1.68 (6H, m), 2.90 (1H, m), 3.14-3.30 (2H, m), 3.33 (4H, m), 3.72 (1H, brn), 3.90 (1H, brn), 5.93 (2H, dd, J = 2Hz, 4Hz), 6.73 (1H, d, J = 8Hz), 6.78 (1H, dd, J = 2Hz, 8Hz), 6.88 (1H, d, J = 2Hz). MS (DCI/NH₃) m/e 517 (M+H⁺). Anal calcd for C₃₀H₄₈N₂O₅ · 0.35 CF₃CO₂H: C, 66.24; H, 8.76; N, 5.03. Found: C, 66.26; H, 8.82; N, 4.98.

30

Example 488

trans,trans-4-(1,4-Benzodioxan-6-yl)-2-(4-fluorophenyl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

35 Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.83 (t, J=7 Hz, 3H), 0.89 (t, J=7 Hz, 3H), 0.90-1.17 (m, 4H), 1.20-1.65 (m, 5H), 2.77d (13, 1H), 2.87 (dd, J=8, 2 Hz, 1H),

2.95-3.60 (m, 7H), 3.71 (d, J=9 Hz, 1H), 4.21 (s, 4H), 6.72 (d, 1H), 6.91 (dd, J=8 Hz, 1H), 6.97 (d, J=2 Hz, 1H), 7.05 (t, J=7 Hz, 2H), 7.40-7.50 (m, 2H). MS (DCI) m/e 513 (M+H)⁺. Anal calcd for C₂₉H₃₇N₂O₅F · 1.2C F₃COOH: C, 58.07; H, 5.93; N, 4.31. Found: C, 57.94; H, 5.81; N, 4.56.

5

Example 489

trans,trans-2-(3-Methylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.83 (3H, t, J = 7Hz), 0.85 (3H, d, J = 7Hz), 0.91 (3H, t, J = 7Hz), 0.97 (3H, t, J = 7Hz), 1.05-1.22 (2H, m), 1.22-1.41 (7H, m), 1.43-1.68 (5H, m), 1.89 (1H, m), 2.94 (1H, t, J = 6Hz), 3.15-3.27 (3H, m), 3.29-3.60 (5H, m), 3.72 (1H, brd, J = 6Hz), 3.92 (1H, brd, J = 13.5Hz), 5.93 (2H, dd, J = 2Hz, 4Hz), 6.73 (1H, d, J = 8Hz), 6.78 (1H, dd, J = 2Hz, 8Hz), 6.88 (1H, d, J = 2Hz). MS (DCI/NH₃) m/e 489 (M+H)⁺. Anal calcd for C₂₈H₄₄N₂O₅ · 0.30 CF₃CO₂H: C, 65.70; H, 8.54; N, 5.36. Found: C, 65.93; H, 8.81; N, 4.84.

15

Example 490

trans,trans-2-(2-Ethylbutyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared and isolated as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (6H, m), 0.92 (3H, t, J = 7Hz), 0.97 (3H, t, J = 7Hz), 1.13-1.41 (13H, m), 1.43-1.72 (6H, m), 2.96 (1H, brm), 3.12-3.52 (6H, m), 3.55-3.70 (1H, m), 3.70-3.86 (2H, m), 3.99 (1H, brm), 5.93 (2H, dd, J = 2Hz, 4Hz), 6.73 (1H, d, J = 8Hz), 6.78 (1H, dd, J = 2Hz, 8Hz), 6.88 (1H, d, J = 2Hz). MS (DCI/NH₃) m/e 489 (M+H)⁺. Anal calcd for C₂₈H₄₄N₂O₅ · 0.45 CF₃CO₂H: C, 64.28; H, 8.30; N, 5.19. Found: C, 64.16; H, 8.38; N, 5.08.

30

Example 491

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(butanesulfonylamino))ethyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 66, the title compound was prepared. ¹H NMR (CD₃OD, 300 MHz) δ 0.74 (d, 3H, J=7), 0.83 (d, 3H, J=7), 0.94 (t, 3H, J=7), 1.44 (hex, 2H), 1.67 (m, 4H), 2.91 (d, 2H, J=8), 3.04 (dd, 2H,

35

J=8, 10), 3.1-3.6 (m, 5H), 3.78 (m, 2H), 3.92 (s, 3H), 4.60 (m, 1H), 5.97 (s, 2H), 6.82 (d, 1H, J=8), 6.89 (dd, 1H, J=2, 8), 7.01 (d, 1H, J=2), 7.22 (t, 1H, J=9), 7.39 (m, 2H). MS (ESI) m/e 579 (M+H)+.

5

Example 492

trans,trans-2-(4-Methoxy-3-fluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(4-ethylpyrimidin-2-yl)amino)ethyl)pyrrolidine-3-carboxylic acid

1-Dimethylamino-1-pentene-3-one, prepared by the method described in Syn. Comm. 12 (1), 35 (1982), was converted to 2-amino-4-ethylpyrimidine with guanidine by the method of Chem. Ber. 97, 3397 (1964). This material was converted to 2-bromo-4-ethyl-pyrimidine with NaNO₂ and HBr, using the method of Helv. Chim. Acta 75, 1629 (1992). This

10

bromopyrimidine was reacted with ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propylamino)propyl)-pyrrolidine-3-carboxylate, prepared using the procedures of Example 61B, using the procedure for Example 418, to give the title compound as a white powder. ¹H NMR (300 MHz, CDCl₃) δ 0.83 (t, J=7Hz, 3H), 1.11 (t, J=7Hz, 3H), 1.45 (sextet, J=7Hz, 2H), 2.18-2.27 (m, 1H), 2.45 (q, J=7Hz, 2H), 2.80-2.97 (m, 3H), 3.40-3.75 (m, 7H), 3.83 (s, 3H), 5.95 (s, 2H), 6.25 (d, J=4Hz, 1H), 6.68 (d, J=8Hz, 1H), 6.79 (dd, J=2Hz, 8Hz, 1H), 6.82 (t, J=9Hz, 1H), 6.92 (d, J=2Hz, 1H), 7.05 (d, J=9Hz, 1H), 7.15 (dd, J=2Hz, 12Hz, 1H), 8.10 (d, J=4Hz, 1H).

15

20

Example 493

trans,trans-4-(1,3-Benzodioxol-5-yl)-2-(4-methoxyphenyl)-1-((N-butyl-N-(3,4-dimethylphenyl)aminocarbonyl)methyl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ¹H NMR (300 MHz, CD₃OD) δ 0.87 (t, J=7.3 Hz, 3H), 1.23-1.36 (m, 2H), 1.38-1.43 (m, 2H), 2.22 (s, 3H), 2.29 (s, 3H), 2.79 (d, J=14.9 Hz, 1H), 2.84 (dd, J=8.6, 9.7 Hz, 1H), 3.16 (t, J=9.5 Hz, 1H), 3.32 (d, J=15.3 Hz, 1H), 3.43-3.61 (m, 4H), 3.79 (s, 3H), 3.88 (d, J=9.8 Hz, 1H), 5.93 (s, 2H), 6.74 (m, 3H), 6.83 (m, 3H), 7.04 (d, J=1.7 Hz, 1H), 7.11 (m, 3H). MS (C.I.) m/e 559(MH⁺). Anal calcd for C₃₃H₃₈N₂O₆•0.3H₂O: C, 70.27; H, 6.90; N, 4.97. Found: C, 70.24; H, 6.62; N, 4.58.

25

30

35

Example 494

trans,trans-2-(3-Methylpent-3-en-1-yl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedure described in Example 1, the title compound was prepared and isolated as an amorphous solid. ^1H NMR (CDCl_3 , 300 MHz) δ 0.92 (3H, t, $J = 7\text{Hz}$), 0.97 (3H, t, $J = 7\text{Hz}$), 1.22-1.40 (5H, m), 1.44-1.61 (8H, m), 1.82 (1H, brn), 2.02 (2H, m), 3.05-3.30 (4H, m), 3.3.8 (1H, m), 3.55 (1H, brn), 3.85 (2H, m), 4.12 (1H, brd, $J = 15\text{Hz}$), 5.11 (1H, dd, $J = 6\text{Hz}$, 12Hz), 5.93 (2H, s), 6.73 (1H, d, $J = 8\text{Hz}$), 6.78 (1H, dd, $J = 2\text{Hz}$, 8Hz), 6.88 (1H, d, $J = 2\text{Hz}$). MS (DCI/NH_3) m/e 487 ($\text{M}+\text{H}$) $^+$. Anal calcd for $\text{C}_{28}\text{H}_{42}\text{N}_2\text{O}_5 \cdot 0.7 \text{CF}_3\text{CO}_2\text{H}$: C, 62.34; H, 7.60; N, 4.95. Found: C, 62.49; H, 7.43; N, 4.73.

Example 495

1-(N-Phenylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Example 495A

N-Phenylbromoacetamide

To a stirred solution of aniline (7.40 mmol) in methylene chloride (25 mL) at -50°C was added successively N,N-diisopropylethylamine (1.58 mL, 8.14 mmol, 1.1 eq) and bromoacetyl bromide (0.72 mL, 7.40 mmol, 1 eq) such that the temperature did not exceed -40°C . On completion of the addition, the cooling bath was removed, and the reaction mixture was allowed to warm to room temperature. After stirring for a further 30 minutes, the mixture was diluted with ether (70 mL) and poured into 1 N sodium bisulfate solution. The phases were separated, and the upper layer was washed successively with water and brine. The organic phase was dried (Na_2SO_4) and the solvent evaporated to half volume, at which point the product crystallized. The crystals were removed by vacuum filtration to afford the title compound.

Example 495B

trans,trans-1-(N-Phenylaminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 and the compound resulting from Example 495A, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 8.8 (bs, 1H) 7.49 (2H, d, $J=8\text{Hz}$), 7.38 (4H, m), 7.11 (1H, tt, $J=8\&2\text{Hz}$), 6.99 (1H, d, $J=2\text{Hz}$), 6.91 (2H, d, $J=8\text{Hz}$), 6.86 (1H, d, $J=2\text{Hz}$), 6.81 (1H, d, $J=8\text{Hz}$), 5.99 (1H, d, $J=2\text{Hz}$), 5.98 (1H, d, $J=2\text{Hz}$), 3.94 (1H, d, $J=10\text{Hz}$), 3.78 (3H, s), 3.70 (1H, ddd, $J=6$, $5\&3\text{Hz}$), 3.42 (1H, dd, $J=10\&3\text{Hz}$), 3.41 (1H, d,

$J=16\text{Hz}$), 3.18 (1H, dd, $J=11\&9\text{Hz}$), 3.01 (1H, t, $J=10\text{Hz}$), 2.93 (1H, d, $J=16\text{Hz}$). MS (DCI, NH_3) m/e 475 ($\text{M}+\text{H}^+$). Anal. Calc for $\text{C}_{27}\text{H}_{26}\text{N}_2\text{O}_6 \cdot 1\text{H}_2\text{O}$: C, 65.85, H, 5.73, N 5.69. Found: C, 65.95, H, 5.52, N, 5.38.

5

Example 496*trans,trans*-1-(N-(2,3-Dimethylphenyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 8.68 (1H, bs), 7.64 (d, $J=8\text{Hz}$), 7.38, (2H, d, $J=8\text{Hz}$), 7.09 (1H, t, $J=8\text{Hz}$), 6.97, (1H, d, $J=8\text{Hz}$), 6.90 (1H, d, $J=2\text{Hz}$), 6.88 (2H, d, $J=8\text{Hz}$), 6.82 (1H, dd, $J=8\&3\text{Hz}$), 6.76 (1H, d, $J=8\text{Hz}$), 5.97 (1H, d, $J=2\text{Hz}$), 5.96 (1H, d, $J=2\text{Hz}$), 3.95 (1H, d, $J=10\text{Hz}$), 3.80 (3H, s), 3.70 (1H, ddd, $J=6, 5\&3\text{Hz}$), 3.48 (1H, dd, $J=10\&3\text{Hz}$), 3.44 (1H, d, $J=16\text{Hz}$), 3.18 (1H, dd, $J=11\&9\text{Hz}$), 3.06 (1H, t, $J=10\text{Hz}$), 2.96 (1H, d, $J=16\text{Hz}$), 2.31 (3H, s), 2.16 (3H, s). MS (DCI, NH_3) m/e 503 ($\text{M}+\text{H}^+$). Anal. Calc for $\text{C}_{29}\text{H}_{30}\text{N}_2\text{O}_6 \cdot 0.5\text{H}_2\text{O}$: C, 68.09, H, 6.11, N, 5.48. Found: C, 68.13, H, 5.91, N, 5.29.

15

Example 497*trans,trans*-1-(N-(2,4-Dimethylphenyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 8.60 (1H, bs), 7.78 (d, $J=8\text{Hz}$), 7.38, (2H, d, $J=8\text{Hz}$), 6.99 (1H, m), 6.95, (1H, d, $J=8\text{Hz}$), 6.94 (1H, d, $J=2\text{Hz}$), 6.88 (2H, d, $J=8\text{Hz}$), 6.82 (1H, dd, $J=8\&3\text{Hz}$), 6.77 (1H, d, $J=8\text{Hz}$), 5.97 (1H, d, $J=2\text{Hz}$), 5.96 (1H, d, $J=2\text{Hz}$), 3.92 (1H, d, $J=10\text{Hz}$), 3.79 (3H, s), 3.68 (1H, ddd, $J=6, 5\&3\text{Hz}$), 3.43 (1H, dd, $J=10\&3\text{Hz}$), 3.42 (1H, d, $J=16\text{Hz}$), 3.18 (1H, dd, $J=11\&9\text{Hz}$), 3.04 (1H, t, $J=10\text{Hz}$), 2.95 (1H, d, $J=16\text{Hz}$), 2.29 (3H, s), 2.24 (3H, s). MS (DCI, NH_3) m/e 503 ($\text{M}+\text{H}^+$). Anal. Calc for $\text{C}_{29}\text{H}_{30}\text{N}_2\text{O}_6 \cdot 0.75\text{H}_2\text{O}$: C, 67.50, H, 6.15, N 5.43. Found: C, 67.42; H, 5.95; N, 5.13.

25

30

Example 498*trans,trans*-1-(N-(2,5-Dimethylphenyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 8.62 (1H, bs), 7.79 (1H, bs), 7.38, (2H, d, $J=8\text{Hz}$), 7.03 (1H, d, $J=8\text{Hz}$), 6.95, (1H, d, $J=8\text{Hz}$), 6.94 (1H, d, $J=2\text{Hz}$), 6.88

35

(2H, d, $J=8\text{Hz}$), 6.82 (1H, dd, $J=8\&3\text{Hz}$), 6.77 (1H, d, $J=8\text{Hz}$), 5.97 (2H, s), 3.92 (1H, d, $J=10\text{Hz}$), 3.78 (3H, s), 3.70 (1H, ddd, $J=6, 5\&3\text{Hz}$), 3.48 (1H, dd, $J=10\&3\text{Hz}$), 3.42 (1H, d, $J=16\text{Hz}$), 3.18 (1H, dd, $J=11\&9\text{Hz}$), 3.04 (1H, t, $J=10\text{Hz}$), 2.95 (1H, d, $J=16\text{Hz}$), 2.29 (3H, s), 2.24 (3H, s). MS (DCI, NH_3) m/e 503 ($\text{M}+\text{H}^+$).

5 Anal. Calc for $\text{C}_{29}\text{H}_{30}\text{N}_2\text{O}_6 \cdot 0.5 \text{H}_2\text{O}$: C, 68.09; H, 6.11; N, 5.48. Found: C, 67.72; H, 5.89; N, 5.25.

Example 499

trans,trans-1-(N-(3,4-Dimethylphenyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 8.73 (1H, bs), 7.38 (2H, bd, $J=8\text{Hz}$), 7.30, (1H, d, $J=3\text{Hz}$), 7.20 (1H, bs), 7.08, (1H, d, $J=8\text{Hz}$), 7.01 (1H, bs), 6.90 (2H, d, $J=8\text{Hz}$), 6.85 (1H, bs), 6.80 (1H, d, $J=8\text{Hz}$), 5.99 (1H, d, $J=3\text{Hz}$), 5.98 (1H, d, $J=3\text{Hz}$), 3.92 (1H, d, $J=10\text{Hz}$), 3.78 (3H, s), 3.70 (1H, ddd, $J=6, 5\&3\text{Hz}$), 3.48 (1H, dd, $J=10\&3\text{Hz}$), 3.42 (1H, d, $J=16\text{Hz}$), 3.18 (1H, dd, $J=11\&9\text{Hz}$), 3.04 (1H, t, $J=10\text{Hz}$), 2.95 (1H, d, $J=16\text{Hz}$), 2.25 (3H, s), 2.21 (3H, s). MS (DCI, NH_3) m/e 503 ($\text{M}+\text{H}^+$). Anal. Calc for $\text{C}_{29}\text{H}_{30}\text{N}_2\text{O}_6 \cdot 0.75 \text{H}_2\text{O}$: C, 67.50; H, 6.15; N 5.43. Found: C, 67.24; H, 5.94; N, 5.20.

Example 500

trans,trans-1-(N-(3,5-Dimethylphenyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1, the title compound was prepared. ^1H NMR (300 MHz, CDCl_3) δ 8.75 (1H, bs), 7.35, (2H, d, $J=8\text{Hz}$), 7.10 (2H, s), 7.02 (1H, d, $J=3\text{Hz}$), 6.90 (2H, d, $J=8\text{Hz}$), 6.84 (1H, d, $J=2\text{Hz}$), 6.80, (1H, d, $J=8\text{Hz}$), 6.76 (1H, bs), 5.99 (1H, d, $J=3\text{Hz}$), 5.98 (1H, d, $J=3\text{Hz}$), 3.92 (1H, d, $J=10\text{Hz}$), 3.79 (3H, s), 3.68 (1H, ddd, $J=6, 5\&3\text{Hz}$), 3.40 (2H, m), 3.18 (1H, dd, $J=11\&9\text{Hz}$), 2.98 (1H, t, $J=10\text{Hz}$), 2.88 (1H, d, $J=16\text{Hz}$), 2.3 (6H, s). MS (DCI, NH_3) m/e 503 ($\text{M}+\text{H}^+$). Anal. Calc for $\text{C}_{29}\text{H}_{30}\text{N}_2\text{O}_6 \cdot 0.5 \text{H}_2\text{O}$: C, 68.09; H, 6.11; N 5.48. Found: C, 67.93; H, 6.01; N, 5.19.

Example 501

Alternate Preparation of

(+)-*trans,trans*-1-(N-Di(*n*-butyl)aminocarbonylmethyl)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylic acid Hydrochloride Salt

Example 501AN, N-Dibutyl bromoacetamide

To a solution of bromoacetyl bromide (72.3 mL, 830 mmol) in toluene (500 mL) cooled to 0 °C was added a solution of dibutylamine (280.0 mL, 1.66 mol) in toluene (220 mL) via an addition funnel maintaining the reaction temperature below 10 °C. Upon completion of the addition, the reaction mixture was stirred at 0 °C for 15 minutes. A solution of 2.5% aqueous H₃PO₄ (500 mL) was slowly introduced, and the reaction mixture was allowed to warm to room temperature with vigorous stirring. The solution is 2.5% phosphoric acid by weight. The layers were separated and the organic phase washed with water (500 mL) and concentrated to provide the bromoacetamide as a solution in toluene.

Example 501B5-(2-Nitrovinyl)-1,3-benzodioxole

To piperonal (15.55 kg, 103.5 mol) under mechanical stirring and under nitrogen was added ammonium acetate (13.4 kg, 173.8 mol), acetic acid (45.2 kg), and nitromethane (18.4 kg, 301.4 mol) sequentially. The mixture was warmed to 70 °C. After about 30 minutes, the yellow product began to crystallize. The reaction temperature was raised to 80 °C and stirred for about 10 hours until minimal piperonal remains. The somewhat thick reaction mixture was cooled to 10 °C and filtered. The precipitate was washed with acetic acid (2 x 8 kg) and then water (2 x 90 kg). The product was dried under a nitrogen purge and then in a vacuum oven at 50 °C for 2 days to afford 15.94 kg (80%) of the title compound as a bright yellow solid.

Example 501C4-Methoxybenzoyl acetate

To potassium t-amylate (25 wt %, 50.8 kg, 99.26 mol) in toluene (15.2 kg) cooled to 5 °C under mechanical stirring and under nitrogen was added a mixture of 4-methoxyacetophenone (6.755 kg, 44.98 mol) and diethyl carbonate (6.40 kg, 54.18 mol) in toluene over 1 hour maintaining the temperature below 10 °C. The reaction mixture was heated to 60 °C for 8 hours until no 4-methoxyacetophenone was detected by HPLC. The mixture was cooled to 20 °C and quenched by adding to a mixture of acetic acid (8 kg) and water (90 kg) over 30 minutes while maintaining the temperature at

<20 °C. The layers were separated, and the organic layer was washed with 5% sodium bicarbonate solution (41 kg) and concentrated to 14.65 kg. The temperature is maintained below 50 °C during the distillation. The yellow product concentrate was assayed by HPLC against an external standard and the yield was found to be 9.40 kg (94%).

Example 501D

Ethyl 2-(4-methoxybenzoyl)-4-nitromethyl-3-(1,3-benzodioxol-5-yl) butyrate

To the compound resulting from Example 501B (7.5 kg, 37.9 mol) suspended in THF (56 kg) with mechanical stirring under nitrogen was added the compound resulting from Example C (8.4 kg, 37.9 mol). The mixture was cooled to 17 °C, sodium ethoxide (6.4 g, 0.095 mol) was added, and the reaction was stirred for 30 minutes. After about 15 minutes, the nitrostyrene was completely dissolved. Sodium ethoxide (6.4 g, 0.095 mol) was added, and the mixture was stirred at 25 °C until HPLC shows less than 1 area % ketoester remaining. The reaction was concentrated to 32.2 kg which was determined by HPLC assay to be ~14.9 kg (95%).

Example 501E

Ethyl *cis, cis*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl) pyrrolidine-3-carboxylate

Raney nickel (20.0 g), from which the water had been decanted, was charged to a stirred hydrogenator equipped with a thermocouple. THF (20 mL), the crude compound resulting from Example 501D (40.82 g, 0.0482 mol), and acetic acid (2.75 mL, 0.0482 mol) were added sequentially. The mixture was put under a hydrogen atmosphere at 60 psi until the hydrogen uptake slowed dramatically. TFA was added, and the mixture was hydrogenated at 200 psi until HPLC shows no residual imine and <2 area % nitron. The catalyst was filtered away and washed with 100 mL of methanol. The filtrate was assayed by HPLC and found to contain 13.3 g (75% yield) of the *cis, cis*-pyrrolidine compound. The filtrate was concentrated and chased with additional THF (200 mL) to give a final volume of 100 mL. The mixture was neutralized with 2 N NaOH solution (50 mL), diluted with water (200 mL), and extracted with ethyl acetate (2 x 100 mL). The combined nearly colorless ethyl acetate layers were assayed against an external standard by HPLC to be 13.0 g (73%) of the title compound.

Example 501F

Ethyl trans, trans-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl) pyrrolidine-3-carboxylate

- The solution of the compound resulting from Example 501E (38.1 g, 0.103 mol) was chased with ethanol (200 mL) to a final volume of 100 mL and sodium ethoxide (3.40 g, 0.050 mol) was added. The mixture was heated to 75 °C. When HPLC shows <3% of the cis,cis isomer remaining, the mixture was cooled to room temperature. The product was assayed by HPLC against an external standard and found to contain 34.4 g (90% yield) of the title compound. The crude compound solution was concentrated and the residue taken up in isopropyl acetate (400 mL). The organic layer was washed with water (2 x 150 mL) and then extracted with 0.25 M phosphoric acid solution (2 x 400 mL). The combined phosphate layers were stirred with ethyl acetate (200 mL) and neutralized to pH 7 with solid sodium bicarbonate (21 g). The organic layer was separated and found to contain 32.9 g (87%) of the title compound.

Example 501G

Ethyl (2R,3R, 4S)-(+)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl) pyrrolidine-3-carboxylate, (S)-(+)- mandelate salt

- The solution resulting from Example 501F was chased with acetonitrile (100 mL) to give a final volume of 50 mL. (S)-(+)-Mandelic acid (2.06 g, 0.0136 mmol) was added and allowed to dissolve. The mixture was seeded with the product and allowed to stir at room temperature for 16 hours. The reaction mixture was cooled to 0 °C and stirred for 5 hours. The product was filtered and dried in a vacuum oven with a nitrogen purge for 1 day at 50 °C to give 5.65 g (40%) of the title compound. The purity of the product can be determined by chiral HPLC using Chiralpak AS, isocratic elution with 95:5:0.05 hexane-ethanol-diethylamine; flow - 1 mL/min.; UV detection at 227 nm. Retention times: (+)-enantiomer: 15.5 min.; (-)-enantiomer: 21.0 min.

Example 501H

(2R,3R,4S)-(+)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)- pyrrolidine-3-carboxylic acid

- The compound resulting from Example 501G (20.0 g, 0.0383 mol) was suspended in ethyl acetate (150 mL) and 5% sodium bicarbonate solution (150 mL). The mixture was stirred at room temperature until the salt dissolved

and carbon dioxide evolution had ceased. The organic layer was separated and concentrated. The residue was chased with acetonitrile (200 mL) to a final volume of 100 mL and cooled to 10 °C. Diisopropylethylamine (11.8 mL, 0.0574 mol) and the compound resulting from Example A (10.5 g, 0.0421 mol) were added, and the mixture was stirred for 12 hours at room temperature. The reaction mixture was concentrated and chased with ethanol (200 mL) to a final volume of 100 mL. Sodium hydroxide solution (40%, 20 mL, 0.200 mol) was added, and the mixture was heated at 60 °C for 4 hours until HPLC showed no starting material remaining. The reaction mixture was poured into water (400 mL) and washed with hexanes (2 x 50 mL). The aqueous layer was washed with hexane (2 x 20 mL). A stirred mixture of the aqueous layer and ethyl acetate (400 mL) was neutralized to pH 5 with concentrated HCl (12 mL). The organic layer was separated and found to contain 18.3 g (94% yield) of the title compound.

Example 501I

(2R,3R,4S)-(+)-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid hydrochloride salt

To a solution of the compound of Example 501H in ethyl acetate at room temperature in a mechanically stirred vessel equipped with a thermocouple, was added 39.4 mL of 1 N HCl in ethanol (0.0394 mol). The resultant solution was filtered to remove foreign matter, concentrated in vacuo, and chased with ethyl acetate (400 mL). The solution was seeded repeatedly, as the solvent was removed, until crystallization was initiated. The mixture was concentrated to a volume of 100 mL, and the product was filtered and washed with ethyl acetate (25 mL). The resultant white solid was dried in a vacuum oven under a nitrogen purge at 50 °C to afford 17.6 g (90%) of the title compound.

Example 502

trans, trans-2-(2-Methylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 502A

(±)-Ethyl 3-methylhexanoate

To a slurry of 60% sodium hydride (2.26g, 57 mmol) in 10mL of hexanes and 100mL of diethyl ether was added triethylphosphonoacetate (10.3mL, 52mmol). Once gas evolution ceased, 2-pentanone (6.0mL, 64mmol) was added. After 3 hours at room temperature, the reaction was quenched with water, and partitioned into ether. The organic layer was washed with water and brine, dried with anhydrous sodium sulfate, filtered, and the solvent was removed under reduced pressure. The residue was dissolved in 50mL of ethanol and 10% palladium on carbon (6.0g) was added. The vessel was pressurized to 4 atmosphere of hydrogen, and was shaken at room temperature for 3 hours. The reaction was filtered and the solvent was removed under reduced pressure to give 3.0g of the title compound.

Example 502B

(±)-Ethyl 5-methyl-3-oxooctanoate

To a solution of ethyl 3-methylhexanoate in 150mL of ethanol was added sodium hydroxide (2.3g, 57.6mmol). After 48 hours at room temperature, solvent was removed under reduced pressure, and the residue was dissolved in 150mL of water. The solution was washed with ether, then acidified with concentrated hydrochloric acid and washed with methylene chloride. The organic layer was dried with anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure to give 2.7g of the corresponding acid from which 3.9g of the title compound was prepared by the method of Bram and Vilkas, *Bul. Chem. Soc. Fr.*, 945 (1964).

Example 502C

trans, trans-2-(2-Methylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 1 and substituting ethyl 5-methyl-3-oxooctanoate for ethyl (4-methoxybenzoyl)acetate afforded the title compound, which was isolated by lyophilization from dilute aqueous TFA/CH₃CN. Note that the multiplicity of the signals in the aryl region of the NMR spectrum reflects a 1:1 mixture of diastereomers on the alkyl chain. ¹H NMR (CDCl₃, 300 MHz) δ 0.8-1.0 (m, 12H), 1.2-1.4 (m, 7H), 1.45-1.6 (m, 6H), 1.6-1.74 (m, 1H), 1.8-2.0 (m, 1H), 3.1-3.4 (m, 5H), 3.67-3.78 (m, 1H), 3.8-3.91 (m, 1H), 4.0-4.2 (m, 2H), 4.3-4.5 (m, 2H), 5.93 (d, J=1.5 Hz, 2H), 6.73 (dd, J=8.1, 1.2 Hz,

1H), 6.79 (ddd, J=7.8, 1.8, 1.8 Hz, 1H), 6.86 (dd, J=3.9, 1.5 Hz, 1H). MS (DCI/NH₃) m/e 489 (M+H)⁺. Anal calcd for C₂₈H₄₄N₂O₅•1.0 TFA•0.5 H₂O: C, 58.91; H, 7.58; N, 4.58. Found: C, 58.91; H, 7.58; N, 4.45.

5

Example 503

trans,trans-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

10

Ethyl 3,3-dimethylhexanoate was prepared using the general procedure of Cahiez *et al.*, Tetrahedron Lett., 31, 7425 (1990). Using the procedures described in Example 502 and substituting ethyl 3,3-dimethylhexanoate for ethyl 3-methylhexanoate afforded the title compound, which was isolated by lyophilization from dilute aqueous TFA/CH₃CN. ¹H NMR (CDCl₃, 300 MHz) δ 0.80-0.99 (m, 15H), 1.10-1.37 (m, 8H), 1.43-1.58 (m, 4H), 1.77-1.97 (m, 2H), 3.48-3.12 (m, 5H), 3.60-3.69 (m, 1H), 3.75-3.86 (m, 1H), 3.95-4.16 (m, 2H), 4.28-4.4 (m, 2H), 5.94 (s, 2H), 6.74 (d, J=7.8 Hz, 1H), 6.8 (dd, J=8.1, 1.5 Hz, 1H), 6.87 (d, J=1.8 Hz, 1H). MS (DCI/NH₃) m/e 503 (M+H)⁺. Anal calcd for C₂₉H₄₆N₂O₅•1.05 TFA: C, 60.01; H, 7.62; N, 4.50. Found: C, 60.21; H, 7.37; N, 4.33.

20

Example 504

trans,trans-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

25

Example 504A

Ethyl 5-(1,3-dioxolyl)-3-oxopentanoate

30

The title compound was synthesized from ethyl acetoacetate and 2-bromomethyl-1,3-dioxane, according to the procedure of Huckin and Weiler, Tetrahedron Lett. 3927, (1971).

35

Sodium hydride 4.97 g (0.124 mol), as a 60% mineral oil dispersion, was weighed into a 250 mL flask, into which 80 ml of tetrahydrofuran was directly added. The flask was capped with septum cap, flushed with nitrogen, and cooled in an ice bath. To above stirred slurry was added dropwise 15.0 mL (0.118 mol) ethyl acetoacetate. After the addition was complete, the resulting mixture was stirred at 0 °C for additional 10 min. To above mixture

was then added 48.4 mL (0.121 mol) *n*-butyl lithium, a 2.50 M solution in hexane, in a dropwise manner. The resulting orange color solution was stirred for 10 min before 13.5 mL (0.130 mol) bromomethyl-1,3-dioxane was added in one portion. The reaction mixture was then allowed to warm to room temperature and stirred for additional 120 min before it was then quenched by slow addition of 9.8 ml (ca. 0.12 mol) concentrated hydrochloric acid. The biphasic mixture was poured to 50 ml of water and extracted with 150 ml of ethyl ether. The aqueous layer was extracted thoroughly with additional ethyl ether. The ethereal extracts were combined, washed with 2x50 ml of saturated brine, dried over anhydrous magnesium sulfate, filtered and evaporated under reduced pressure to give an brown oily residue. The crude product was purified using silica gel flash chromatography eluting with 20% ether/hexane to give 5.40 g (20%) of b-keto ester as a light yellow oil.

Example 504C

trans,trans-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502 and substituting ethyl 5-(1,3-dioxolyl)-2-oxopentanoate for ethyl 3-methylhexanoate afforded the title compound. ¹H NMR (CDCl₃, 300 MHz) δ 0.93 (t, J = 7.2 Hz, 3H), 0.95 (t, J = 7.2 Hz, 3H), 1.23-1.38 (m, 4H), 1.52 (sextet, J = 7.9 Hz, 4H), 1.85-1.95 (m, 2H), 2.02-2.17 (m, 2H), 3.18 (dd, J = 6.0 Hz, 9.0 Hz, 2H), 3.30 (dd, J = 9.0 Hz, 18.0 Hz, 2H), 3.35 (m, 1H), 3.79 (dd, J = 3.6 Hz, 6.9 Hz, 1H), 3.83-3.88 (m, 3H), 3.97 (dd, J = 4.8 Hz, 6.0 Hz, 1H), 4.05 (q, J = 9.6 Hz, 2H), 4.30-4.40 (m, 1H), 4.37 (s, 2H), 4.87 (t, J = 3.6 Hz, 1H), 5.94 (s, 2H), 6.73 (d, J = 8.1 Hz, 1H), 6.79 (dd, J = 1.8 Hz, 8.1 Hz, 1H), 6.87 (d, J = 1.8 Hz, 1H). MS (APCI) (M+H)⁺ at m/e 505. Anal calcd for C₂₇H₄₀N₂O₇·1.2 TFA: C, 55.05; H, 6.47; N, 4.37. Found: C, 55.12; H, 6.44; N, 4.27.

Example 505

trans,trans-2-(2-(2-Tetrahydro-2H-pyran)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 505A

Ethyl 5-(2-tetrahydro-2H-pyran)-3-oxopentanoate

Using the procedure of Huckin and Weller, Tetrahedron Lett. 3927, (1971), the title compound was prepared from ethyl acetoacetate and 2-(bromomethyl)tetrahydro-2H-pyran as a light yellow oil.

5

Example 505B

trans,trans-2-(2-(2-Tetrahydro-2H-pyran)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

10

Using the procedures described in Example 502 and substituting ethyl 5-(2-tetrahydro-2H-pyran)-2-oxopentanoate for ethyl 3-methylhexanoate afforded the title compound as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) as a mixture of two diastereoisomers: δ 0.89 (t, J = 8.1 Hz, 3H), 0.89 (t, J = 8.1 Hz, 3H), 0.91 (t, J = 8.1 Hz, 3H), 0.91 (t, J = 8.1 Hz, 3H), 1.20-1.40 (m, 10H), 1.42-1.66 (m, 18H), 1.71 (brm, 2H), 1.85 (brm, 2H), 1.96-2.23 (brm, 4H), 3.10-3.29 (m, 8H), 3.29-3.52 (m, 6H), 3.54-3.81 (m, 6H), 4.01 (q, J = 9 Hz, 2H), 4.12-4.25 (m, 4H), 4.43 (d, J = 9 Hz, 2H), 4.50 (d, J = 2.7 Hz, 2H), 5.94 (s, 2H), 5.95 (s, 2H), 6.76 (s, 2H), 6.76 (s, 2H), 6.81 (s, 1H), 6.81 (s, 1H). MS (APCI) (M+H)⁺ at m/e 517. Anal calcd for C₂₉H₄₄N₂O₆·1.4 TFA: C, 56.48; H, 6.77; N, 4.14. Found: C, 56.46; H, 6.99; N, 3.83.

20

Example 506

trans,trans-2-(2,2,4-Trimethyl-3-pentenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

25

Example 506A

Methyl 3,3,5-trimethyl-4-hexenoate

30

To a slurry of isopropyltriphenylphosphonium iodide (20.5g, 47mmol) in 200mL of tetrahydrofuran was added n-butyllithium (27mL of a 1.6M solution in hexane, 43mmol), and the solution was briefly warmed to 0°C. After recooling, a solution of methyl 3,3-dimethyl-4-oxobutenoate (5.7g, 40mmol), prepared according to the procedure of Hudlicky *et al.*, Synth. Commun., **16** 169 (1986) in 10mL of tetrahydrofuran was added, and the reaction was warmed to 0°C for 30min. The reaction was quenched with dilute hydrochloric acid, and partitioned with ethyl acetate. The organic layer was washed with water, and brine, dried with anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure. The residue

35

was purified by flash chromatography on silica gel eluting with 10% ethyl acetate in hexanes to give 2.1g (30%) of the title compound.

Example 506B

- 5 *trans, trans*-2-(2,2,4-Trimethyl-3-pentenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502 and substituting methyl 3,3,5-trimethyl-4-hexenoate for ethyl 3-methylhexanoate afforded the title compound, which was isolated by lyophilization from dilute aqueous TFA/CH₃CN. ¹H NMR (CDCl₃, 300 MHz) δ 0.92 (t, J=7.2 Hz, 3H), 0.94 (t, J=7.2 Hz, 3H), 1.11 (s, 3H), 1.13 (s, 3H), 1.24-1.37 (m, 4H), 1.46-1.59 (m, 4H), 1.61 (d, J=1.2 Hz, 3H), 1.69 (d, J=1.2 Hz, 3H), 2.04-2.11 (m, 2H), 3.10-3.20 (m, 2H), 3.30-3.39 (m, 3H), 3.67-3.82 (m, 2H), 3.95-4.08 (m, 1H), 4.32 (m, 2H), 4.37-4.47 (m, 1H), 4.99 (s, 1H), 5.95 (s, 2H), 6.73 (d, J=7.8 Hz, 1H), 6.78 (dd, J=8.4, 1.2 Hz, 1H), 6.84 (d, J=1.2 Hz, 1H). MS (DCI/NH₃) m/e 515 (M+H)⁺. Anal calcd for C₃₀H₄₆N₂O₅•1.05 TFA: C, 60.77; H, 7.48; N, 4.42. Found: C, 60.83; H, 7.20; N, 4.43.

Example 507

trans, trans-2-(2,2-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 507A

Methyl 3,3-dimethyl-3-(1,3-dioxolan-2-yl)propanoate

Methyl 3,3-dimethyl-4-oxobutanoate (10g, 70mmol), prepared according to the procedure of Hudlicky *et al.*, Synth. Commun., **16** 169 (1986), was dissolved in 40mL of benzene, followed by addition of ethylene glycol (20mL), and *p*-toluenesulfonic acid monohydrate (1.3g). The reaction was refluxed with azeotropic removal of water for 1 hour. The reaction was poured into 200mL of ether, washed with saturated sodium bicarbonate, water and brine, dried with anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure to give 12.4g (94%) of the title compound.

Example 507B

trans, trans-2-(2,2-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502 and substituting methyl 3,3-dimethyl-3-(1,3-dioxolan-2-yl)propanoate for ethyl 3-methylhexanoate afforded the title compound, which was isolated by lyophilization from dilute aqueous TFA/CH₃CN. ¹H NMR (CDCl₃, 300 MHz) δ 0.82-1.00 (m, 12H), 1.24-1.40 (m, 4H), 1.43-1.64 (m, 5H), 1.76-1.84 (m, 1H), 2.93-3.00 (m, 1H), 3.15-3.47 (m, 6H), 3.60-3.70 (m, 3H), 3.74-3.95 (m, 5H), 4.48 (s, 1H), 5.94 (m, 2H), 6.72 (d, J=8.0 Hz, 1H), 6.83 (dd, J=8.0, 1.2 Hz, 1H), 6.94 (d, J=1.2 Hz, 1H). MS (DCI/NH₃) m/e 533 (M+H)⁺. Anal calcd for C₂₉H₄₄N₂O₇ • 1.1 TFA • 0.2 H₂O: C, 56.63; H, 6.93; N, 4.23. Found: C, 56.60; H, 6.9%; N, 4.25.

Example 508

trans,trans-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-((N-4-heptyl-N-(2-methyl-3-fluorophenyl)) amino carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 508A4-Heptanol

To an ice cooled solution of 1.14g (10.0 mmol) of 4-heptanone in 20 mL of diethyl ether was added 370 mg (10.0 mmol) of LiAlH₄, in portions to keep ether reflux at a minimum. After 45 minutes, the reaction was quenched by sequential dropwise addition of 0.4 mL H₂O, 0.4 mL 15% (w/v) NaOH(aq), and 1.2 mL H₂O. After stirring another 45 minutes, MgSO₄ was added until the salts were free flowing, then the reaction was filtered. The salts were washed with diethyl ether (3 x 5 mL), then the filtrate and washings were concentrated to a colorless oil. Yield 1.16g (100%).

Example 508B4-Methanesulfonyloxyheptane

To an ice cooled solution of 834 mg (7.19 mmol) of 4-heptanol in 35 mL of CH_2Cl_2 was added 1.5 mL of triethylamine. Next, 0.7 mL (9 mmol) of methanesulfonyl chloride was added, dropwise, over 1 minute. The mixture was stirred at 0 °C for 30 minutes, then extracted with H_2O (1 x 15 mL), 5% NH_4OH (2 x 15 mL), 1M HCl (2 x 15 mL), and brine (1 x 15 mL), dried over MgSO_4 , filtered, and concentrated to an oil. Yield 1.31g (94%). ^1H NMR (300 MHz, CDCl_3) δ 0.96 (t, 6, $J = 9$), 1.43 (m, 4), 1.64 (m, 4), 3.00 (s, 3), 4.73 (quintet, 1 $J = 5$).

Example 508C

4-Fluoro-3-methylaniline

To a solution of 20g (129 mmol) of 2-fluoro-5-nitrotoluene in 400 mL of ethanol was added 2g of 10% Pd-C. The mixture was shaken under 45 P.S.I. H_2 until hydrogen uptake ceased. The catalyst was filtered away and washed with ethanol, then the combined filtrate and washings were concentrated to 15.2 g (94%) of a colorless oil.

Example 508D

N-Heptyl-4-fluoro-3-methylaniline

To a solution of 4.10 g (3.28 mmol) of 4-fluoro-3-methylaniline in 30 mL of acetonitrile was added 7.64 g (3.93 mmol) of 4-methanesulfonyloxyheptane, and 3.4 g (4.1 mmol) of $\text{NaHCO}_3(\text{s})$. The mixture was stirred at reflux for 24 hours, then poured into 150 mL of H_2O and extracted with diethyl ether (2 x 30 mL). The combined ether layers were back extracted with brine (1 x 30 mL), dried over MgSO_4 , filtered, and concentrated to an oil. This was purified via silica gel chromatography, eluting with 97.5: 2.5 hexanes: ethyl acetate, to give 2.56g (35%) of a pale yellow oil.

Example 508E

N,N-(4-Heptyl)-(4-fluoro-3-methyl)phenylbromoacetamide

To an ice cooled solution of 4.88g (21.9 mmol) of N-(4-heptyl)-4-fluoro-3-methylaniline and 4.9 mL (61 mmol) of pyridine in 100 mL of toluene was added a solution of 4.90 mL (56.2 mmol) of bromoacetyl bromide in 7 mL of

toluene. The solution was stirred for 24 hours, gradually warming to 25 °C, then extracted with 1M HCl (1 x 100 mL). The aqueous layer was back extracted with diethyl ether (1 x 50 mL), then the combined organic layers were washed with H₂O (2 x 50 mL), saturated NaHCO₃(aq) (2 x 50 mL), and brine (1 x 50 mL), dried over MgSO₄, filtered, and concentrated *in vacuo* to an oil. This was purified via silica gel chromatography, eluting with 90:10 hexanes: ethyl acetate to give 7.48g (99%) of a light yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 0.94 (t, 6, J = 5), 1.33 (m, 4), 1.43 (m, 4), 2.30 (s, 1.5), 2.31 (s, 1.5), 3.54 (s, 2), 4.72 (quintet, 1, J = 5), 6.96-7.04 (m, 2), 7.07(d, 1, J = 7).

Example 508F

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-((*N*-4-heptyl-*N*-(2-methyl-3-fluorophenyl)) amino carbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, substituting ethyl 5-(1,3-dioxolyl)-2-oxopentanoate for ethyl 3-methylhexanoate and *N,N*-(4-heptyl)-(4-fluoro-3-methyl)phenyl-bromoacetamide for *N,N*-diethylbromoacetamide afforded the title compound as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.93 (brt, 6H), 1.23-1.47 (m, 8H), 1.67-2.10 (m, 4H), 2.32 (s, 3H), 3.16 (t, J = 9.0 Hz, 1H), 3.52-3.67 (brn, 2H), 3.73 (t, J = 9.0 Hz, 1H), 3.81-4.02 (m, 6H), 4.13 (brn, 1H), 4.72 (quintet, J = 6.9 Hz, 1H), 4.86 (t, J = 4.0 Hz, 1H), 5.93 (s, 2H), 6.72 (d, J = 8.1 Hz, 1H), 6.78 (dd, J = 1.8 Hz, 8.1 Hz, 1H), 6.85 (d, J = 1.8 Hz, 1H), 6.96 (m, 2H), 7.08 (t, J = 9.0 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 599. Anal Calcd for C₃₃H₄₃N₂O₇F·0.8 TFA: C, 60.24; H, 6.40; N, 4.06. Found: C, 60.21; H, 6.14; N, 3.86.

Example 509

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, substituting ethyl 5-(1,3-dioxolyl)-2-oxopentanoate for ethyl 3-methylhexanoate and 6-methoxyxypiperonal for piperonal afforded the title compound as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.93 (t, J = 7.8 Hz, 3H), 0.95 (t, J = 7.8 Hz, 3H), 1.31 (m, 4H), 1.53 (m, 4H), 1.90 (m, 2H), 2.09 (m, 2H), 3.19 (dd, J =

8.4 Hz, 8.4 Hz, 2H), 3.30 (q, J = 9.6 Hz, 2H), 3.25-3.42 (m, 1H), 3.73 (q, J = 10.5 Hz, 1H), 3.78-3.94 (m, 4H), 3.88 (s, 3H), 3.96 (dd, J = 5.1 Hz, 6.0 Hz, 1H), 4.03 (dd, J = 3.0 Hz, 6.3 Hz, 2H), 4.33 (m, 3H), 4.87 (t, J = 3.6 Hz, 1H), 5.94 (s, 2H), 6.53 (d, J = 1.8 Hz, 1H), 6.63 (d, J = 1.8 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 535. Anal calcd for C₂₈H₄₂N₂O₈·1.05 TFA: C, 55.25; H, 6.63; N, 4.28. Found: C, 55.39; H, 6.66; N, 4.26.

Example 510

trans,trans-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, substituting o-methoxyphenoxyacetic acid for 3-methylhexanoic acid, the above compound was prepared as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (t, J=7Hz, 3H), 0.90 (t, J=7Hz, 3H), 1.15-1.35 (m, 4H), 1.40-1.55 (m, 4H), 3.05-3.25 (m, 4H), 3.28-3.55 (m, 4H), 3.58-3.68 (m, 1H), 3.75-3.80 (m, 1H), 3.82 (s, 3H), 3.91 (d, J=14Hz, 1H), 4.05-4.15 (m, 1H), 4.23-4.33 (m, 1H), 5.91 (s, 2H), 6.70 (d, J=8Hz, 1H), 6.82-6.95 (m, 5H), 7.03 (s, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 541. Anal calcd for C₃₀H₄₀N₂O₇: C, 66.65; H, 7.46; N, 5.18. Found: C, 66.37; H, 7.61; N, 5.09.

Example 511

(2S,3R,4S)-2-(2-(2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 511A

trans, trans-N-tert-Butoxycarbonyl-2-(2,2-dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylic acid

Ethyl *trans, trans*-2-(2,2-dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate (2.5g, 6.9mmol), prepared according to Example 503, was dissolved in 50mL of methylene chloride and di-*tert*-butyldicarbonate (1.5g) was added. After stirring overnight at room temperature, the solvent was removed under reduced pressure and the residue was purified by flash chromatography on silica gel eluting with 10%

ethyl acetate/hexanes to give the ethyl ester of the title compound (2.8g) as a colorless oil. The ester was dissolved in 50mL of ethanol followed by addition of sodium hydroxide (10mL of a 5M aqueous solution). After stirring for 20 hours at room temperature, the solvent was removed under reduced pressure, and the residue was dissolved in 150mL of water, and acidified with concentrated phosphoric acid. The mixture was extracted with chloroform (3X50mL), and the organic layers were washed with brine, dried over anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure to give the title compound (2.4g) as a white foam.

Example 511B

Methyl trans, trans-2-(2,2-dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylate: As a single enantiomer

The product from Example 510A (1.97g, 4.5 mmol) was dissolved in 20mL of THF and cooled to 0°C, followed by addition of DMF (0.017mL, 5%), and oxalyl chloride (0.437mL, 5.00mmol). After 1 hour, solvent was removed at 0°C under a stream of nitrogen. The residue was dissolved in 5mL of benzene and evaporated. In a separate flask, (S)-4-benzyl-2-oxazolidinone (1.2g, 6.8mmol) was dissolved in 30mL of THF followed by addition of n-butyllithium (4.0mL of a 1.6M solution in hexanes) at 0°C, and the slurry was stirred for 15min. The acid chloride was dissolved in 20mL of THF and cooled to 0°C, followed by dropwise addition of the lithium oxazolidone suspension via cannula. After 30min, the reaction was partitioned between ether and saturated bicarbonate. The organic phase was washed with water then brine, dried over anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel eluting with 15% ethyl acetate/hexanes to give the undesired diastereomer (1.17g, 43%), then elution with 20% ethyl acetate/hexanes gave the desired diastereomer (1.04g, 38%).

The desired diastereomer of the N-acyloxazolidinone (0.84g, 1.42mmol) was dissolved in 2.5mL of dichloromethane, and 2.5mL of trifluoroacetic acid was added. After 30min, the volatiles were removed under a stream of nitrogen, and the residue was twice dissolved in 5mL of toluene and evaporated under reduced pressure.

The TFA salt was stirred with 4mL of acetonitrile followed by addition of diisopropylethyl amine (1.0mL, 5.7mmol), and N-4-heptyl-N-(4-fluoro-3-methylphenyl)bromoacetamide (589mg, 1.7mmol) as a solution in 2mL of acetonitrile. After 21 hours, the reaction was warmed to 50°C for 3.5 hours.

The reaction was cooled, the solvent removed under reduced pressure, and the residue was purified by flash chromatography on silica gel eluting with 20-30% ethyl acetate/hexanes to give 0.939g of amide as a colorless oil.

The above amide (200mg, 0.26mmol) was dissolved in 2.0mL of THF and 0.7mL of water. Solid lithium hydroxide monohydrate (22mg, 0.53mmol) was added at 0°C, followed by 30% hydrogen peroxide (0.050mL, 0.55mmol). After 1 hour, the reaction was warmed to room temperature. After an additional hour, the reaction was partitioned between 1:1 ethyl acetate:hexanes and water, 0.15g of sodium thiosulfate was added and the mixture was mixed thoroughly. The organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure. The crude residue was dissolved in 2mL of ether, and 1mL of methanol. A solution of (trimethylsilyl)diazomethane in hexanes was added dropwise until the yellow color remained. The reaction was quenched by addition of 2 drops of glacial acetic acid, and the solvent was removed under reduced pressure. The residue was purified by flash chromatography on 10g of silica gel eluting with 15-20% ethyl acetate/hexanes to give 70mg of the title compound as a crystalline solid (mp137.5°C).

Example 511C

(2S,3R,4S)-trans, trans-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylate

The product from Example 510B (65mg, 0.10mmol) was dissolved in 1.0mL of methanol and sodium hydroxide (0.1mL of a 5M aqueous solution) was added. After 2 hours, the reaction was warmed to reflux. After 6 hours, the reaction was cooled, and the solvent was removed under reduced pressure. The residue was dissolved in water and acidified with concentrated phosphoric acid. The aqueous solution was washed with chloroform (3X5mL), which was then washed with brine, dried with anhydrous magnesium sulfate, filtered and evaporated under reduced pressure. The title compound was

isolated by lyophilization from dilute aqueous TFA/CH₃CN. ¹H NMR (CDCl₃, 300 MHz) δ 0.78-0.95 (m, 15H), 1.04-1.46 (m, 12H), 1.76-2.95 (m, 2H), 2.31 (s, 3H), 3.23-3.33 (m, 1H), 3.47-3.58 (m, 1H), 3.6-3.75 (m, 2H), 3.80-3.95 (m, 2H), 4.05-4.15 (m, 1H), 4.73 (m, 1H), 5.94 (s, 2H), 6.70-6.80 (m, 2H), 6.82-6.93 (m, 2H), 6.96-7.14 (m, 2H). MS (DCI/NH₃) m/e 597 (M+H)⁺. Anal calcd for C₃₅H₄₉N₂FO₅ • 0.05H₂O • 0.8TFA: C, 63.81; H, 7.30; N, 4.07. Found: C, 63.84; H, 7.18; N, 3.94. $[\alpha]_D^{21} = +46^\circ$ (c 2.7g/L, CHCl₃)

Example 512

- 10 trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 512A

2-Oxopyrrolidin-1-ylpropionic acid

15

To a stirred solution of 5.0 mL (40.5 mmol) 2-oxopyrrolidin-1-ylpropionitrile in 15 mL of dioxane was added 8.1 mL of hydrochloric acid, a 6.0 M aqueous solution. The resulting mixture was then refluxed at 110 °C over night. The reaction mixture was then allowed to cool to room temperature, extracted with methylene chloride three times. The extracts were combined and washed with saturated brine solution once, dried over anhydrous sodium sulfate, filtered and evaporated under reduced pressure to give 1.60 g (25%) of acid as a brown oil.

20

25

Example 512B

Ethyl 5-(2-oxopyrrolidin-1-yl)-3-oxopentanoate

The title compound was prepared from the above acid by adapting the method of Bram and Vilkas, Bul. Chem. Soc. Fr., 945 (1964).

30

Example 512C

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

35

Using the procedures described in Example 502, substituting ethyl 5-(2-oxopyrrolidin-1-yl)-3-oxopentanoate for ethyl 3-methylhexanoate afforded the title compound as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.91

(t, J = 7.5 Hz, 3H), 0.94 (t, J = 7.5 Hz, 3H), 1.23-1.38 (m, 4H), 1.44-1.60 (m, 4H), 2.05 (t, J = 6.9 Hz, 2H), 2.12-2.25 (m, 1H), 2.38 (td, J = 4.2 Hz, 8.4 Hz, 2H), 2.47-2.61 (m, 1H), 3.17 (dd, J = 6.0 Hz, 8.7 Hz, 2H), 3.24 (t, J = 9 Hz, 1H), 3.32 (t, J = 7.8 Hz, 2H), 3.38-3.48 (m, 3H), 3.52 (t, J = 9 Hz, 1H), 3.66 (t, J = 6.9 Hz, 1H), 3.96 (m, 2H), 4.14 (m, 1H), 4.38 (brs, 2H), 5.93 (s, 2H), 6.74 (d, J = 8.1 Hz, 1H), 6.89 (dd, J = 1.8 Hz, 8.1 Hz, 1H), 6.87 (d, J = 1.8 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 516. Anal calcd for C₂₈H₄₁N₃O₆·1.4 TFA: C, 54.78; H, 6.33; N, 6.22. Found: C, 54.69; H, 6.33; N, 6.14.

Example 513

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, substituting ethyl 5-(1,3-dioxolyl)-2-oxopentanoate for ethyl 3-methylhexanoate, N-4-heptyl-N-(4-fluoro-3-methylphenyl) bromoacetamide for N,N-dibutyl bromoacetamide and 6-methoxypiperonal for piperonal afforded the title compound as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.93 (br t, 6H), 1.23-1.47 (m, 8H), 1.67-2.10 (m, 4H), 2.32 (s, 3H), 3.16 (t, J = 9 Hz, 1H), 3.60-4.03 (m, 8H), 3.88 (s, 3H), 4.21 (brs, 1H), 4.72 (quintet, J = 6.6 Hz, 1H), 4.86 (t, J = 3.6 Hz, 1H), 5.93 (s, 2H), 6.49 (s, 1H), 6.61 (s, 1H), 6.96 (m, 2H), 7.08 (t, J = 9 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 629. Anal calcd for C₃₄H₄₅N₂O₈F·1.0 TFA: C, 58.21; H, 6.24; N, 3.77. Found: C, 58.11; H, 6.11; N, 3.58.

Example 514

trans,trans-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, substituting ethyl 5-methyl-3-oxooctanoate for ethyl 3-methylhexanoate and 6-methoxypiperonal for piperonal afforded the title compound as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.81 (s, 3H), 0.84 (s, 3H), 0.86 (t, J = 6.9 Hz, 3H), 0.93 (t, J = 6.9 Hz, 3H), 0.96 (t, J = 6.9 Hz, 3H), 1.09-1.38 (m, 8H), 1.45-1.59 (m, 4H), 1.84-2.00 (m, 2H), 3.15 (dd, J = 6.9 Hz, 10.0 Hz, 2H), 3.30-3.42 (m, 3H), 3.72 (t, J = 10.5 Hz, 1H), 3.86 (t, J = 10.5 Hz, 1H), 3.88 (s, 3H), 4.02 (q, J =

10.0 Hz, 1H), 4.12 (d, J = 16.8 Hz, 1H), 4.29 (d, J = 16.8 Hz, 1H), 4.41 (brn, 1H), 5.94 (s, 1H), 6.52 (d, J = 1.8 Hz, 1H), 6.67 (d, J = 1.8 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 533. Anal calcd for C₃₀H₄₈N₂O₆·0.9 TFA: C, 60.12; H, 7.76; N, 4.41. Found: C, 60.18; H, 7.62; N, 4.33.

5

Example 515

trans,trans-2-(2,2-dimethylpentyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- 10 Using the procedures described in Example 502, substituting ethyl 3,3-dimethylhexanoate for ethyl 3-methylhexanoate and 2,3-dihydro-benzofuran-5-carbaldehyde for piperonal afforded the title compound as an amorphous solid by lyophilization with CH₃CN/TFA/H₂O. ¹H NMR (300 MHz, CDCl₃) δ 0.83 (s, 3H), 0.85 (s, 3H), 0.86 (t, J=7.2 Hz, 3H), 0.92 (t, J=7.2 Hz, 3H),
- 15 0.95 (t, J=7.2 Hz, 3H), 1.09-1.39 (m, 8H), 1.44-1.59 (m, 4H), 1.88 (dd, J=15.0, 7.2 Hz, 1H), 2.00 (d, J=15.0 Hz, 1H), 3.09 (m, 2H), 3.18 (t, J=9.0 Hz, 2H), 3.27-3.38 (m, 3H), 3.65-3.95 (m, 2H), 4.05 (q, J=10.0 Hz, 1H), 4.18 (d, J=16.8 Hz, 1H), 4.30-4.45 (m, 2H), 4.55 (t, J=9.0 Hz, 2H), 6.70 (d, J=8.4 Hz, 1H), 7.04 (dd, J=8.4, 2.1 Hz, 1H), 7.23 (brs, 1H). MS (DCI/NH₃) at m/e 501 (M+H)⁺. Anal calcd for
- 20 C₃₀H₄₈N₂O₄·1.05 TFA: C, 62.14; H, 7.97; N, 4.51. Found: C, 62.19; H, 8.00; N, 4.43.

Example 516

- trans,trans-2-(2,2-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid
- 25

- Using the procedures described in Example 502, substituting methyl 3,3-dimethyl-3-(1,3-dioxolan-2-yl)propanoate for ethyl 3-methylhexanoate and 6-methoxypiperonal for piperonal afforded the title compound as an amorphous solid by lyophilization with CH₃CN/TFA/H₂O. ¹H NMR (CDCl₃, 300 MHz) δ 0.93 (t, J=7.2 Hz, 3H), 0.94 (t, J=7.2 Hz, 3H), 0.95 (s, 3H), 0.96 (s, 3H), 1.31 (sextet, J=7.2 Hz, 4H), 1.45 (m, 4H), 1.93 (dd, J=15.9, 6.0 Hz, 1H), 2.13 (d, J=15.9 Hz, 1H), 3.20 (dd, J=7.7, 7.7 Hz, 1H), 3.26-3.40 (m, 3H), 3.60 (m, 1H), 3.75-
- 30 3.86 (m, 3H), 3.88 (s, 3H), 3.93-4.01 (m, 3H), 4.00-4.11 (m, 1H), 4.23 (d, J=15.9 Hz, 1H), 4.37-4.48 (m, 2H), 4.49 (s, 1H), 5.94 (s, 2H), 6.51 (d, J=2.1 Hz, 1H), 6.64 (d,
- 35

J=2.1 Hz, 1H). MS (DCI/NH₃) at m/e 563 (M+H)⁺. Anal calcd for C₃₀H₄₆N₂O₈·0.9 TFA: C, 57.41; H, 7.11; N, 4.21; found: C, 57.35; H, 6.86; N, 4.05.

5 Example 517

trans,trans-2-(2-(2-Methoxyphenyl)-ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, substituting o-methoxyphenylpropionic acid for 3-methylhexanoic acid, the above compound was prepared as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.85 (t, J=7Hz, 3H), 0.91 (t, J=7Hz, 3H), 1.10-1.27 (m, 4H), 1.42-1.60 (m, 4H), 1.72-1.89 (m, 1H), 1.91-2.02 (m, 1H), 2.55-2.77 (m, 2H), 2.94 (t, J=6Hz, 1H), 3.05-3.30 (m, 6H), 3.59-3.82 (m, 3H), 3.73 (d, J=14Hz, 1H), 3.77 (s, 3H), 5.91 (s, 2H), 6.70 (d, J=8Hz, 1H), 6.78-6.88 (m, 3H), 6.92 (d, J=2Hz, 1H), 7.08-7.19 (m, 2H). MS (DCI/NH₃) (M+H)⁺ at m/e 539. Anal calcd for C₃₁H₄₂N₂O₆: C, 69.12; H, 7.86; N, 5.20. Found: C, 68.89; H, 7.70; N, 4.99.

20 Example 518

trans,trans-2-(2,2-Dimethyl-3-(E)-pentenyl)-4-(1-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

25 Example 518A

4-Methyl-3-penten-2-ol

To a stirred solution of 3-methyl-2-butenal (8.7g, 103mmol) in 100mL of tetrahydrofuran under N₂ at 0 °C was added methylmagnesium bromide (38mL of a 3.0M solution in ethyl ether, 114mmol) dropwise. The resulting mixture was allowed to warm to room temperature slowly and stirred at room temperature for 1 hour before it was quenched with 25mL of saturated NH₄Cl. The resulting biphasic mixture was partitioned between ethyl ether and water. The organic layer was washed with brine, dried with anhydrous magnesium sulfate, filtered, and the solvent was removed under reduced pressure to give 8.4g (81%) of alcohol as a colorless oil.

Example 518Btrans-Ethyl 3,3-dimethyl-4-pentenoate

- A mixture of 4-methyl-3-penten-2-ol (7.4g, 74mmol), triethyl
 5 orthoacetate (13.6mL, 74mmol) and propionic acid (0.28mL, 3.7mmol) was
 heated at 150 °C for 7 hours. The product was then distilled under normal
 pressure (200-220 °C) to give 5.0g of crude ester as a colorless oil.

Example 518C

- 10 trans, trans-2-(2,2-Dimethyl-3-(E)-pentenyl)-4-(1-methoxy-1,3-benzodioxol-5-yl)-
1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- Using the procedures described in Example 502, substituting *trans*-ethyl
 3,3-dimethyl-4-pentenoate for ethyl 3-methylhexanoate and 6-
 15 methoxypiperonal for piperonal afforded the title compound as an
 amorphous solid by lyophilization from dilute aqueous TFA/CH₃CN. ¹H NMR
 (CDCl₃, 300 MHz) δ 0.92 (t, J=7.2 Hz, 3H), 0.95 (t, J=7.2 Hz, 3H), 0.97 (s, 3H), 0.99
 (s, 3H), 1.31 (sextet, J=7.2 Hz, 4H), 1.52 (quintet, J=7.2 Hz, 4H), 1.58 (d, J=5.4 Hz,
 3H), 1.92 (dd, J=15.0, 6.6 Hz, 1H), 2.04 (d, J=15.0 Hz, 1H), 3.15 (dd, J=7.8, 7.8 Hz,
 20 1H), 3.30-3.40 (m, 3H), 3.75 (m, 2H), 3.87 (s, 3H), 3.99 (q, J=9 Hz, 2H), 4.11-4.30
 (m, 3H), 5.29 (d, J=15.6 Hz, 1H), 5.38 (dd, J=15.6, 6 Hz, 1H), 5.94 (s, 2H), 6.50 (d,
 J=1.8 Hz, 1H), 6.63 (d, J=1.8 Hz, 1H). MS (DCI/NH₃) at m/e 531 (M+H)⁺.
 Analysis calc'd for C₃₀H₄₆N₂O₆·0.95 TFA: C, 59.95; H, 7.41; N, 4.38; found: C,
 60.00; H, 7.33; N, 4.35.

Example 519

trans, trans-2-(3-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-
butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

30

Example 519A3-(2-Pyridyl)-propionic Acid

- In a 50 mL round-bottomed flask equipped with a stirring bar was
 placed 3-(2-pyridyl)-propanol (1 g, 7.6 mmol), water (13 mL) and
 35 concentrated sulfuric acid (0.5 g, 5.1 mmol). To this stirred solution was
 added over a period of 30 min potassium permanganate (1.8 g, 11.3 mmol)

while the reaction temperature was maintained at 50 °C. After the addition was completed, the mixture was held at 50 °C until the color of the reaction mixture turned brown, then heated at 80 °C for 1 hour and filtered. The filtrate was evaporated to dryness to yield quantitatively the desired acid (1.14 g) suitable for next step without further purification. To prepare a pure acid, the residue thus obtained was boiled in ethanol (10 mL) in the presence of charcoal (0.1 g) for 5 min, filtered and cooled to give crystalline 3-(2-pyridyl)-propionic acid (0.88 g, 78%).

Example 519B

trans, trans-2-(3-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedure described in Example 502, the title compound was isolated by lyophilization from dilute aqueous TFA/CH₃CN as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 8.65 (d, J=6.0 Hz, 1H), 8.06 (t, J=6.91 Hz, 1H), 7.70 (d, J=9.0 Hz, 1H), 7.51 (t, J=6.91 Hz, 1H), 6.82-6.66 (m, 3H), 5.91 (s, 2H), 4.45 (s, 2H), 4.29-4.18 (m, 1H), 4.04 (dd, J=20.1, 10.5 Hz, 1H), 3.84 (t, J=12.6 Hz, 1H), 3.62 (dd, J=13.8, 9.6 Hz, 1H), 3.46-3.13 (m, 7H), 2.51 (broad s, 2H), 1.60-1.43 (m, 4H), 1.37-1.22 (m, 4H), 0.91 (t, J=8.4 Hz, 6H). MS (DCI/NH₃) m/e 510 (M+H)⁺. Anal calcd for C₂₉H₃₉N₃O₅ • 1.75 TFA: C, 55.04; H, 5.79; N, 5.92. Found: C, 55.08; H, 5.64; N, 5.81.

Example 520

(2S, 3R, 4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 520A

(2S, 3R, 4S)-Ethyl-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)pyrrolidine-3-carboxylate-(S)-Mandelate

The racemic amino ester from Example 512 (3.45g, 8.98mmol) in 10mL of ethyl acetate was treated with (S)-(+)-mandelic acid (0.75g, 4.93mmol). Upon the formation of the clear solution, hexane was dropped in slowly with stirring till the solution became light cloudy. The solution was left stirred at room temperature over night. The crystals was then collected by filtration,

recrystallized from ethyl acetate/hexane twice to give a yield of 800 mg (17%) of pure salt.

Example 520B

- 5 (2S, 3R, 4S)-Ethyl-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylate

To a stirred solution of pure mandelate (150 mg, 0.28 mmol) in CH₃CN was added *N,N*-dibutylbromoacetamide (84 mg, 0.34 mmol) and diisopropylethylamine (98 μ L, 0.56 mmol). The resulting mixture was stirred at room temperature over night. Solvent was then removed under reduced pressure and the crude product was purified by silica gel flash chromatography to give 140 mg (90% yield) of the title compound.

- 15 Example 520C

(2S, 3R, 4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedures described in Example 502, the title compound was prepared as an amorphous solid by lyophilization with CH₃CN/TFA/H₂O. ¹H NMR (CDCl₃, 300 MHz) δ 0.91 (t, J = 7.5 Hz, 3H), 0.94 (t, J = 7.5 Hz, 3H), 1.23-1.38 (m, 4H), 1.44-1.60 (m, 4H), 2.05 (t, J = 6.9 Hz, 2H), 2.12-2.25 (m, 1H), 2.38 (td, J = 4.2 Hz, 8.4 Hz, 2H), 2.47-2.61 (m, 1H), 3.17 (dd, J = 6.0 Hz, 8.7 Hz, 2H), 3.24 (t, J = 9 Hz, 1H), 3.32 (t, J = 7.8 Hz, 2H), 3.38-3.48 (m, 3H), 3.52 (t, J = 9 Hz, 1H), 3.66 (t, J = 6.9 Hz, 1H), 3.96 (m, 2H), 4.14 (m, 1H), 4.38 (brs, 2H), 5.93 (s, 2H), 6.74 (d, J = 8.1 Hz, 1H), 6.89 (dd, J = 1.8 Hz, 8.1 Hz, 1H), 6.87 (d, J = 1.8 Hz, 1H). MS (DCI/NH₃) (M+H)⁺ at m/e 516. Anal calcd for C₂₈H₄₁N₃O₆·0.85 TFA: C, 58.23; H, 6.89; N, 6.86. Found: C, 58.37; H, 6.90; N, 6.84.

- 30 Example 521

(2S, 3R, 4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

35 Using the procedures described in Example 520, substituting *N,N*-(4-heptyl)-(4-fluoro-3-methyl)phenyl-bromoacetamide for *N,N*-

dibutylbromoacetamide afforded the title compound as an amorphous solid by lyophilization with CH₃CN/TFA/H₂O. ¹H NMR (CDCl₃, 300 MHz) δ 0.85-0.98 (m, 6H), 1.22-1.55 (m, 8H), 2.04 (quintet, J=7.9 Hz, 4H), 2.32 (s, 3H), 2.36 (t, J=7.9 Hz, 2H), 2.61 (m, 1H), 3.14 (m, 1H), 3.25-3.61 (m, 5H), 3.66-3.77 (m, 1H), 3.79-3.90 (m, 2H), 3.92-4.03 (m, 1H), 4.69 (quintet, J=6.8 Hz, 1H), 5.95 (s, 2H), 6.71 (s, 2H), 6.78 (s, 1H), 6.93-7.13 (m, 3H); MS (DCI/NH₃) at m/e 610 (M+H)⁺. Anal calc'd for C₃₄H₄₄N₃O₆F₁·1.45 TFA: C, 57.18; H, 5.91; N, 5.42. Found: C, 57.20; H, 5.62; N, 5.52.

Example 522

trans, trans-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 522A

3-(1-Pyrazolyl)-propionic Acid

In a 10 mL round-bottomed flask equipped with a condenser and a stirring bar was placed pyrazole (0.50 g, 7.3 mmol), acrylic acid (0.50 mL, 7.3 mmol) and triethylamine (3 mL). The reaction mixture was refluxed for 6 hours. After removing triethylamine, the viscous oil was dried on high vacuo during 12 hours to yield quantitatively the desired acid (1.0 g) suitable for the next step without further purification.

Example 522B

trans, trans-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Using the procedure described in Example 502, the title compound was isolated by lyophilization from dilute aqueous TFA/CH₃CN as an amorphous solid. ¹H NMR (CDCl₃, 300 MHz) δ 7.56 (d, J=3.0 Hz, 1H), 7.50 (d, J=3 Hz, 1H), 6.83-6.66 (m, 3H), 6.28 (t, J=3 Hz, 1H), 5.91 (s, 2H), 4.55-3.98 (m, 6H), 3.83-3.72 (t, J=10.5 Hz, 1H), 3.61-3.40 (t, J=10.5 Hz, 1H), 3.36-3.12 (m, 5H), 2.69-2.43 (m, 2H), 1.59-1.42 (m, 4H), 1.38-1.21 (m, 4H), 0.91 (t, J=7.5 Hz, 6H). MS (DCI/NH₃) at m/e 499 (M+H)⁺. Anal calcd for C₂₇H₃₈N₄O₅•0.75 TFA: C, 58.60; H, 6.69; N, 9.59. Found: C, 58.53; H, 6.45; N, 9.67.

Example 523

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

5

Example 523AN-Butyl-N-(3-hydroxypropyl)-amine

To a solution of 15.9g (100 mmol) of methyl 3-*N*-(*n*-butyl)aminopropionate in 150 mL of diethyl ether at 0 °C was added 50 mL (0.35 mmol) of 1.0M LiAlH₄ in diethyl ether, keeping reflux at a minimum. The mixture was stirred at 0 °C for 2.25 hours, then quenched by sequential dropwise addition of 1.9 mL H₂O, 1.9 mL 15%w/v NaOH(aq), and 5.7 mL H₂O. After stirring for 30 min, the salts were filtered and washed with diethyl ether, then the filtrate was concentrated to 11.3 g (86%) of a light yellow oil.

15

Example 523BN-Butyl-N-(3-hydroxypropyl)-chloroacetamide

To an ice cooled solution of 1.31g (10.0 mmol) of *N*-butyl,*N*-(3-hydroxypropyl)amine in 20 mL of ethyl acetate was added a solution of 1.71g (10.0 mmol) of chloroacetic anhydride in 10mL of ethyl acetate. The mixture was stirred, and gradually warmed to room temperature over 18 hours. The reaction was extracted with H₂O (1 x 50 mL), saturated NaHCO₃ (aq) (2 x 50 mL), and brine (1 x 50 mL), dried over MgSO₄, filtered, and concentrated to an oil. The product was purified via silica gel chromatography, eluting with 80:20 hexanes:ethyl acetate to give 723 mg (35%) of a light yellow oil.

25

Example 523C

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

30

Using the procedures described in Example 1D, substituting *N*-butyl-*N*-(3-hydroxypropyl)-chloroacetamide for *N*-propyl bromoacetamide and adding DMSO as cosolvent, afforded the title compound, which was isolated by lyophilization from dilute aqueous TFA/CH₃CN. ¹H NMR (CD₃OD, 300 MHz) δ 0.78-0.95 (m, 3H), 1.00-1.80 (m, 4H), 2.80-3.65 (m, 15H), 3.80 (d, J=1.5 Hz, 2H), 5.93 (s, 2H), 6.72-7.05 (m, 5H), 7.33-7.40 (m, 2H). MS (DCI/NH₃) at m/e 513

35

(M+H)⁺. Anal calc'd for C₂₈H₃₆N₂O₇•1.6 H₂O: C, 62.12; H, 7.30; N, 5.17.
Found: C, 62.04; H, 7.21; N, 4.88.

Example 524

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propyl-N-propoxyamino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 524A

N-Boc-O-allylhydroxylamine

O-Allylhydroxylamine hydrochloride hydrate (5.0g) was dissolved in THF (15 mL). The solution was cooled to 0°C in an ice bath. Diisopropylethylamine (8mL) and di-*t*-butyldicarbonate (10.0g) were added. The mixture was stirred at 0°C for 1 hour at which point the bath was removed and the reaction allowed to warm to room temperature and stirred overnight. The THF was removed *in vacuo* and the residue taken up in EtOAc (25 mL), and washed with water (1 x 50 mL), saturated sodium bicarbonate solution (3 x 50 mL), 1N phosphoric acid (3 x 50 mL), and brine (1 x 50 mL). The organic layer was dried with sodium sulfate and evaporated to give a light yellow oil (6.5g) which was used without any further purification.

Example 524B

N-Boc-N-propyl-O-allylhydroxylamine

N-Boc-O-allylhydroxylamine (6.5g) from the above procedure was dissolved in dry THF (25 mL) and the solution cooled to 0°C in an ice bath. Sodium hydride (1.5g, 60% dispersion in oil) was added portionwise over 5 min. The resulting mixture was stirred for 30 min at 0°C. 1-Iodopropane (3.8mL) was added dropwise to the mixture. The reaction was stirred at 0°C for 1 hour, then stirred overnight at room temperature. The THF was removed *in vacuo* and the residue taken up in EtOAc (50 mL) and washed with water (1 x 50 mL), saturated sodium bicarbonate solution (3 x 50 mL), 1N phosphoric acid (3 x 50 mL), and brine (1 x 50 mL). The organic layer was dried with sodium sulfate and evaporated to give a light yellow oil, which was purified by flash chromatography on silica gel eluting with 5% EtOAc/hexanes to give the title compound as a colorless oil (6.0g).

Example 524CN-Boc-N-propyl-N-propoxyamine

- 5 N-Boc-N-propyl-O-allylhydroxylamine (6.0g) was dissolved in EtOAc (100 mL). 10% Palladium-on-carbon (0.5g) was added, and the mixture was purged with nitrogen. The nitrogen line was exchanged for a balloon of hydrogen, and the mixture was stirred at room temperature for 6 hours. The catalyst was removed by filtration through a pad of Celite and the solvents
- 10 were removed *in vacuo* to give a yellow oil which was purified by flash chromatography on silica gel eluting with 5% EtOAc/hexanes to give the title compound as a colorless oil (5.8g).

Example 524DN-Propyl-N-propoxyamine hydrochloride

- 15 N-Boc-N-propyl-N-propoxyamine (5.8g) was dissolved in 4N HCl/dioxane (10mL) and stirred at room temperature for 7 hours. The solvent was removed *in vacuo* and the residue triturated with diethyl ether. The
- 20 resulting yellow solid (2.1g) was collected by filtration and washed with diethyl ether.

Example 524EN-propyl-N-propoxy-bromoacetamide

- 25 N-Propyl-N-propoxyamine hydrochloride (0.30 g) was dissolved in acetonitrile and cooled to -20°C. Pyridine (0.2 mL) was added. Bromoacetyl bromide (0.15g) was added dropwise over 5 min. The solution was stirred at -20°C for 30 min. The bath was removed and the solution was stirred for 6
- 30 hours at room temperature. The solvent was removed *in vacuo* and the residue taken up in EtOAc (50 mL) and washed with water (1 x 25 mL), 1N phosphoric acid (3 x 25 mL), and brine (1 x 25 mL). The organic layer was dried with sodium sulfate and evaporated to give a dark orange oil (0.35g). The product is a mixture of chloro- and bromoacetamides in a ratio of ~3:1.

Example 524F

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-hydroxypropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedure of Example 523C, employing N-propyl-N-propoxy-bromoacetamide and ethyl 2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate. The crude product was purified by preparative HPLC (Vydac mC18) eluting with a 10-70% gradient of CH₃CN in 0.1% TFA. The appropriate fraction was lyophilized to give the product as a white solid. ¹H NMR (CDCl₃, 300 MHz) δ 0.87 (m, 6H, J=8Hz), 1.49 (m, 2H, J=8Hz), 1.61 (m, 2H, J=8Hz), 3.55 (m, 6H), 3.80 (m, 2H), 3.81 (s, 3H), 4.00 (m, 2H), 4.13 (d, 2H, J=17Hz), 5.96 (s, 2H), 6.77 (d, 1H, J=9Hz), 6.90 (m, 3H), 7.05 (d, 1H, J=1Hz), 7.44 (d, 2H, J=9Hz). MS (DCI/NH₃) m/e 499 (M+H)⁺. Anal calcd for C₂₇H₃₄N₂O₇ . 1.20 TFA: C, 55.57; H, 5.58; N, 4.41. Found: C, 55.59; H, 5.58; N, 4.55.

Example 525

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-propoxyamino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 525A

N-butyl-N-(2-hydroxyethyl)-amine

In a thick walled glass tube 5 ml (100 mmol) of ethylene oxide was condensed at -78°C. To this 12.5 ml (120 mmol) of butylamine was added and the tube was sealed. The resultant solution was allowed to heat in an oil bath at 50°C for 18 hours. Unreacted reagents were removed by evaporation to give the title compound.

Example 525B

N-Butyl-N-(2-azidoethyl)-chloroacetamide

To 500 mg of N-butyl-N-2-hydroxyethylamine was added 2 mL of thionyl chloride, dropwise. After the initial reaction had ceased, the reaction was stirred for 10 min, then concentrated to an oil. Diethyl ether was added and evaporated to aid in removal of the thionyl chloride. The residue was taken up in 10 mL of DMF, and 1.0g (16 mmol) of sodium azide was added. The

reaction was stirred at 75 °C for 2 hours, then poured into 50 mL of 0.6M NaHCO₃(aq.) and extracted with diethyl ether (3 x 15 mL). The combined ether layers were back extracted with brine (1 x 15 mL), dried over MgSO₄, and filtered. To the ether solution was added 850 mg (4.97 mmol) of chloroacetic anhydride. The reaction was stirred for 10 min, then concentrated to an oil. This was taken up in 10 mL of saturated NaHCO₃(aq.) and extracted with diethyl ether (3 x 5 mL). The combined ether layers were back extracted with brine (1 x 5 mL), dried over MgSO₄, filtered, and concentrated to an oil. This was purified via silica gel chromatography, eluting with 30% ethyl acetate: hexanes, to give 161 mg (17%) of an oil.

Example 525C

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(2-aminoethyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

According to the procedure of Example 523C, N-butyl-N-(2-azidoethyl)-chloroacetamide was coupled with ethyl 2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate. The crude product was chromatographed on silica, using 40% EtOAc in hexanes to elute. The product was dissolved in a solution of ethanol and aqueous 2.5 N sodium hydroxide and stirred for 3 hours at room temperature. The solution was concentrated *in vacuo* and water added. The mixture was extracted with ether; the aqueous layer was acidified to pH 4 with 1N H₃PO₄ and extracted with EtOAc. The latter organic extract was washed with brine and dried over Na₂SO₄. To 100 mg (0.10 mmol) of the azide was added 1mL of 1M HCl(aq.), 0.5 mL of dioxane, and 5 mg of 10% Pd-C. The suspension was stirred under 1 atm. of H₂ for 5 hours, then filtered and concentrated to a white solid. The product was purified via HPLC, eluting with a 0 to 70 CH₃CN in 0.1% aqueous TFA gradient to give the title compound as its TFA salt. ¹H NMR (CD₃OD, 300 MHz) δ 0.92 (t, J=7.0 Hz, 3H), 0.96 (t, rotamer), 1.23 (m, 2H), 1.41 (m, 2H), 3.06 (m, 4H), 3.39 (m, 2H), 3.69 (m, 2H), 3.84 (s, 3H), 3.94 (m, 3H), 4.18 (m, 2H), 5.05 (bd, J=10.7 Hz, 1H), 5.98 (s, 2H), 6.84 (d, J=7.7 Hz, 1H), 6.93 (dd, J=1.8, 8.1 Hz, 1H), 7.05 (m, 3H), 7.56 (m, 2H). MS (DCI/NH₃) at m/e 498 (M+H)⁺. Anal calcd for C₂₇H₃₅N₃O₆•3.15 TFA: C, 46.68. H, 4.49. N, 4.90. Found: C, 46.61; H, 4.73; N, 4.79.

Example 526

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-aminopropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

To and ice-cold solution of the compound of Example 523C (100 mg, 0.19 mmol) in 1 mL of dichloromethane was added 17mL of methanesulfonyl chloride, and 39 mL of triethylamine. The mixture was stirred for 20 min, then diluted with 1.5 mL of dichloromethane and extracted once with 5mL of water to which had been added 1 drop of 85% H₃PO₄, then 5% ammonium hydroxide (1 x 2.5 mL), and brine (1 x 2.5 mL), dried over MgSO₄, filtered, and concentrated to an oil. To a solution of 81 mg (0.13 mmol) of the mesylate in 1mL of DMF was added 65 mg (10 mmol) of sodium azide. The mixture was stirred for 1 hour at 50 °C, then poured into 10 mL of water and extracted with diethyl ether (3 x 5 mL). The combined ether layers were back extracted with brine (1 x 5 mL), dried over MgSO₄, filtered, and concentrated to an oil. This was purified via silica gel chromatography, eluting with 60:40 hexanes: ethyl acetate to give 57 mg of a colorless oil. The product was dissolved in a solution of ethanol and aqueous 2.5 N sodium hydroxide and stirred for 3 hours at room temperature. The solution was concentrated *in vacuo* and water added. The mixture was extracted with ether; the aqueous layer was acidified to pH 4 with 1N H₃PO₄ and extracted with EtOAc. The latter organic extract was washed with brine and dried over Na₂SO₄. To this azide was added 1mL of 1M HCl(aq.), 0.5 mL of dioxane, and 5 mg of 10% Pd-C. The suspension was stirred under 1 atm. of H₂ for 5 hours, then filtered and concentrated to a white solid. The product was purified via HPLC, eluting with a 0 to 70 CH₃CN in 0.1% aqueous TFA gradient to give the title compound as its TFA salt. ¹H NMR (D₆-DMSO, 300 MHz) δ 0.85 (apparent q, J=6.8 Hz, 3H), 1.17 (m, 2H), 1.30 (m, 2H), 1.67 (m, 2H), 2.71 (m, 2H), 3.04 (m, 1H), 3.21 (m, 3H), 3.45 (m, 1H), 3.75 (m, 3H), 3.97 (s, 3H), 3.85-4.80 (broad m, 3H), 6.03 (m, 2H), 6.87 (dd, J=1.4, 8.1 Hz, 1H), 6.92 (d, J=7.8 Hz, 1H), 7.01 (m, 2H), 7.16 (m, 1H), 7.55 (m, 2H), 7.72 (m, 2H), 7.85 (m, 1H); MS (DCI/NH₃) (M+H)⁺ at m/e 512. Anal calcd for C₂₈H₃₇N₃O₆•3.0 TFA: C, 47.84. H, 4.72. N, 4.92. Found: C, 47.86; H, 4.75; N, 4.97.

Example 527

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-dimethylaminopropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 527A

N-butyl-N-(3-bromopropyl)bromoacetamide

To 1.50g (11.4 mmol) of N-butyl-N-(3-hydroxy)propylamine was added 3 mL of 48% HBr(aq.), and 1.5 mL of conc. H₂SO₄. The reaction was stirred at reflux for 3 hours, then cooled to room temperature and stirred for 22 hours. The mixture was poured over 50 mL of ice, and the solution was treated with 50 mL of 2M NaOH(aq.). The basic solution was extracted with ethyl acetate (3 x 25 mL), then the combined ethyl acetate layers were back extracted with brine (1 x 25 mL), dried, and filtered. To the ice cooled ethyl acetate solution was added 3mL of triethylamine, then 1.5 mL of bromoacetyl bromide as a solution in 3.5 mL of ethyl acetate. The reaction was stirred at 0 °C for 30 min, then extracted with 1M HCl(aq.) (2 x 25 mL) saturated NaHCO₃(aq.) (1 x 25 mL) and brine (1 x 25 mL). The organic layer was dried over MgSO₄, filtered, and concentrated to an oil. This was purified via silica gel chromatography, eluting with 30% ethyl acetate in hexanes to give 1.47g of a colorless oil.

Example 527B

Ethyl *trans, trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-bromopropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

According to the procedure of Example 523C, N-butyl-N-(3-bromopropyl)bromoacetamide was coupled with ethyl 2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate. The crude product was chromatographed on silica, using 40% EtOAc in hexanes to elute.

Example 527C

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-dimethylaminopropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

To 400 mg (0.663 mmol) of the compound of Example 527B in 4 mL of absolute EtOH was added 1.2 mL of 2.0 M Me₂NH in THF. The reaction was heated at 50 °C for 3h, then stirred at room temperature for 18 hours. The

mixture was concentrated, then reconcentrated from CH₃CN to remove most of the trimethylamine. The product was purified via silica gel chromatography, eluting with 9:1 CH₂Cl₂: MeOH over about 20 mL of silica gel to give the ethyl ester. The product was dissolved in a solution of ethanol and aqueous 2.5 N sodium hydroxide and stirred for 3 hours at room temperature. The solution was concentrated *in vacuo* and water added. The mixture was extracted with ether; the aqueous layer was acidified to pH 4 with 1N H₃PO₄ and the product was purified by preparative HPLC. ¹H NMR (CD₃OD, 300 MHz) δ 0.92 (t, J=7.0 Hz, 3H), 1.22 (m, 2H), 1.39 (m, 2H), 1.90 (m, 2H), 2.87 (s, 6H), 3.07 (m, 4H), 3.24 (m, 1H), 3.43 (m, 1H), 3.62 (m, 1H), 3.84 (s, 3H), 3.88 (m, 3H), 4.07 (m, 1H), 4.17 (m, 1H), 4.97 (m, 1H), 5.97 (s, 2H), 6.83 (d, J=8.1 Hz, 1H), 6.93 (dd, J=1.7, 8.1 Hz, 1H), 7.05 (m, 3H), 7.53 (m, 2H). MS (DCI/NH₃) at m/e 540 (M+H)⁺. Anal calcd for C₃₀H₄₁N₃O₆•2.95 TFA: C, 49.22. H, 5.06. N, 4.80. Found: C, 49.16; H, 5.11; N, 4.62.

Example 528

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-trimethylammoniopropyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 527C, substituting aqueous Me₃N for Me₂NH. ¹H NMR (CD₃OD, 300 MHz) δ 0.91 (m, 3H), 1.24 (m, 2H), 1.40 (m, 2H), 1.99 (m, 2H), 3.13 (s, 9H), 3.18 (s, rotamer), 3.20 (m, 3H), 3.39 (m, 4H), 3.72 (m, 1H), 3.84 (s, 3H), 4.03 (m, 3H), 4.35 (m, 1H), 5.19 (m, 1H), 5.97 (s, 2H), 6.84 (d, J=8.1 Hz, 1H), 6.96 (dd, J=1.7, 7.9 Hz, 1H), 7.10 (m, 3H), 7.62 (m, 2H). MS (DCI/NH₃) at m/e 554 (M+H)⁺. Anal calcd for C₃₁H₄₄N₃O₆•0.1 H₂O•1.65 TFA: C, 47.25. H, 4.96. N, 4.32. Found: C, 47.25; H, 4.74; N, 4.75.

Example 529

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-aminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 529A

N-butyl-N-(4-hydroxybutyl)-amine

A solution of 8.1 g (110 mmol) of *n*-butylamine and 8.6 g of butyrolactone in 50 ml toluene was allowed to reflux under nitrogen atmosphere for 50 hours. Volatile solvents were removed *in vacuo*. To a solution of 3.18 gm (20 mmol) of the resultant N-butyl-4-hydroxybutyramide in 50 ml of toluene were added 120 ml (120 mmol) DIBAL(25%W). The solution was heated with stirring at 70 °C for 18 hours. After cooling to 0 °C, the reaction was quenched with methanol (1/3 amount of DIBAL solution was used) followed by addition of saturated solution of Rochelle's salt. The mixture was extracted twice with EtOAc; the organic extracts were washed with brine and dried over Na₂SO₄.

Example 529B

N-butyl-N-(4-hydroxybutyl)-chloroacetamide

Pyridine (2 ml) was added to an ice cold solution of 0.58 gm (4 mmol) of N-butyl-N-(4-hydroxybutyl)-amine in 10 ml of EtOAc. To this solution 0.769 gm (4.5 mmol) chloroacetic anhydride was added in small portions. The reaction mixture was allowed to stir for 5 hours at 0 °C, and then was allowed to warm to room temperature. Bicarbonate was added, and the resultant mixture was extracted with EtOAc. The organic layer was washed with water and brine. The crude material was purified by column chromatography.

Example 529C

Ethyl *trans, trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-hydroxybutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

According to the procedure of Example 523C, N-butyl-N-(4-hydroxybutyl)-chloroacetamide was coupled with ethyl 2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate. The crude product was chromatographed on silica gel.

Example 529D

Ethyl *trans, trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-bromobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

To the solution of 0.180 gm (0.33 mmol) of the compound of Example 529C in 2 ml DMF 0.086 gm (1 mmol) of lithium bromide and 0.120 ml (0.66

mmol) of PBr₃ was added. The reaction mixture was allowed to stir at 0°C for 2 hours and was slowly warmed to room temperature. Bicarbonate was added, and the resultant mixture was extracted with EtOAc. The organic layer was washed with water and brine. The crude material was purified by column chromatography.

Example 529E

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-aminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

To a solution of 0.135 gm (0.21 mmol) of the compound of Example 529D in 2 ml DMF was added 0.1 gm of sodium azide. Reaction was allowed to stir at room temperature for 18 hours under nitrogen atmosphere. After addition of water, the product was extracted into EtOAc. The crude product (117 mg) was dissolved in 10 ml ethanol under nitrogen atmosphere. To this 45 mgs of 10% Pd/C catalyst was added, the nitrogen from the reaction flask was evacuated and was flushed with hydrogen by placing a balloon filled with hydrogen.

The reaction was allowed to stir for 4 hours under hydrogen atmosphere, and was worked up by filtering through a Celite pad. The product was dissolved in a solution of ethanol and aqueous 2.5 N sodium hydroxide and stirred for 8 hours at room temperature. The solution was concentrated *in vacuo* and water added. The mixture was extracted with ether; the aqueous layer was acidified to pH 4 with 1N H₃PO₄, and the product was purified by

preparative HPLC. ¹H NMR (CD₃OD, 300 MHz) δ 0.90 (t, J=7 Hz, 3H), 1.10-1.65 (m, 6H), 2.85-2.95 (m, 2H), 3.00-4.10 (m, 14H), 5.50 (d, J=3 Hz, 2H), 5.97 (s, 2H), 6.82 (d, J=8 Hz, 1H), 6.91 (dd, J=7 Hz, 1H), 7.00-7.06 (m, 3H), 7.45-7.55 (m, 2H). MS (DCI/NH₃) at m/e 526 (M+H)⁺. Anal calc'd for C₂₉H₃₉N₃O₆·2.2 TFA: C, 51.75; H, 5.35; N, 5.41. Found: C, 51.75; H, 5.31; N, 5.30.

Example 530

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared from the compound of Example 529D, employing the procedures of Example 527C. ¹H NMR (CD₃OD, 300

MHz) δ 0.90 (dt, J=7Hz, 3H), 1.1-1.75 (m, 8H), 2.75 (d, J=7 Hz, 6H), 3.0-4.25 (m, 16H), 5.97 (s, 2H), 6.83 (d, J=8 Hz, 1H), 6.93 (dd, J=8 Hz, 1H), 7.02-7.08 (m, 3H), 7.49-7.56 (m, 2H). MS (DCI/NH₃) at m/e 554 (M+H)⁺. Anal calc'd for C₃₁H₄₃N₃O₆•2.1 TFA: C, 53.31; H, 5.73; N, 5.30. Found: C, 53.50; H, 5.38; N, 5.34.

Example 531

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-pyridyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 531A

N-butyl-N-(3-pyridyl)-amine

To a solution of 941 mg (10 mmol) of 3-aminopyridine and 0.9 mL of butyraldehyde in 30 mL of CH₃OH was added 10 mL of glacial acetic acid. The mixture was stirred at room temperature for 1 hour, then the reaction was cooled with an ice bath, and 650 mg (10.3 mmol) of sodium cyanoborohydride was added. The ice bath was removed, and the reaction was stirred for 4.5 hours at room temperature. The mixture was poured into 300 mL of 0.67M NaOH(aq.), and extracted with ethyl acetate (3 x 50 mL). The combined organic layers were back extracted with brine (1 x 50 mL), dried over MgSO₄, filtered, and concentrated to an oil. The product was isolated via silica gel chromatography, eluting with 3:1 ethyl acetate: hexanes to give 1.18g (79%) of a colorless solid.

Example 531B

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-pyridyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 531A was reacted according to the procedures of Example 523, to give the title compound. ¹H NMR (D₆-DMSO, 300 MHz) δ 0.80 (t, J=6.4 Hz, 3H), 1.15-1.99 (m, 4H), 2.59 (m, 1H), 3.05 (m, 2H), 3.26 (m, 2H), 3.49 (m, 2H), 3.56 (t, J=7.1 Hz, 2H), 3.73 (s, 3H), 6.00 (s, 2H), 6.80 (m, 3H), 6.85 (d, J=8.1 Hz, 1H), 6.98 (m, 2H), 7.04 (m, 1H), 7.41 (dd, J=1, 4.7 Hz, 8.1H), 7.58 (m, 1H), 8.36 (bs, 1H), 8.54 (bs, 1H), 12.24 (bs, 1H). MS (DCI/NH₃) at

m/e 532 (M+H)⁺. Anal calcd for C₃₀H₃₃N₃O₆•0.1 H₃PO₄: C, 66.55. H, 6.20. N, 7.76. Found: C, 66.59; H, 6.06; N, 7.60.

Example 532

5 *trans, trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-aminomethylphenyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 532A

N-butyl-N-(3-hydroxymethylphenyl)-amine

10

To a solution of 3.69 g (30 mmol) of 3-amino benzyl alcohol in 20 ml DMSO was added 3.78 g (45 mmol) solid NaHCO₃ and 2.91 ml (27 mmol) 1-bromobutane. The reaction was allowed to stir at 50 °C for 18 hours (overnight). Reaction was worked up by adding 250 ml water and product was extracted in ethyl acetate. Water was added, and the resultant mixture was extracted with EtOAc. The organic layer was washed with water and brine.

15

Example 532B

20

N-butyl-N-(3-hydroxymethylphenyl)-bromoacetamide

To a solution of 3.42 g (19.2 mmol) of the compound of Example 532A in 20 ml toluene, was added 2.42 ml (30 mmol) pyridine. The mixture was cooled to 0°C; 4.025 gm (20.0 mmol) of bromoacetyl bromide (diluted with 5 ml toluene) was added in a dropwise fashion. The reaction mixture was allowed to stir for 5 hours at 0°C and then was allowed to warm to room temperature. Saturated potassium carbonate solution was added, and the mixture was stirred vigorously for 2 hours. The mixture was extracted with EtOAc; the organic layer was washed with 1N H₃PO₄, water, and brine.

25

30

Example 532C

Ethyl *trans, trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-chloromethylphenyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

35

According to the procedure of Example 523C, N-butyl-N-(3-hydroxymethylphenyl)-bromoacetamide was coupled with ethyl 2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate. The crude product (129 mg) was dissolved in 0.5 ml of DMF and cooled to 0°C; 19 mg of
 5 LiCl was added, followed by 85 μ l of thionyl chloride. The mixture was allowed to stir for 30 min; water was added, and the mixture was extracted with EtOAc. The organic extracts were washed with water and brine, and dried over Na₂SO₄.

Example 532D

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-aminomethylphenyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 532C (182 mg) was dissolved in 1 mL of
 15 DMF. Two drops of water were added, followed by 126 mg (2.0 mmol, 6.5 eq) of sodium azide. The resultant solution was heated at 115 °C for 3 hours. Water was added, and the mixture was extracted with EtOAc. The organic extracts were washed with water and brine, and dried over Na₂SO₄.

Example 532E

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-aminomethylphenyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

In a 50 ml round bottom flask 0.090 gm Tin (II) chloride was suspended
 25 in 1 ml acetonitrile. Triethylamine (0.2 mL) was added, followed by 0.19 ml of thiophenol; the reaction mixture turned yellow. Reaction flask was cooled to 0°C in ice bath; a solution of 0.185 gm of the compound of Example 532D in 2 ml acetonitrile was added. The mixture was allowed to stir for 30 min. Ether (10 ml) was added, followed by addition of 10 ml 2N HCl. The aqueous
 30 extract was basified with 4N NaOH and extracted with dichloromethane. The organic layer was washed with water and brine. The crude product was dissolved in a solution of ethanol and aqueous 2.5 N sodium hydroxide and stirred for 8 hours at room temperature. The solution was concentrated *in vacuo* and water added. The mixture was extracted with ether; the aqueous
 35 layer was acidified to pH 4 with 1N H₃PO₄, and the product was purified by preparative HPLC. ¹H NMR (CD₃OD, 300 MHz) δ 0.88 (t, J=7 Hz, 3H), 1.15-1.45

(m, 4H), 3.40-4.20 (m, 14H), 5.97 (s, 2H), 6.82 (d, J=8 Hz, 1H), 6.88 (dd, J=8 Hz, 1H), 6.97-7.20 (m, 5H), 7.40 (d, J=9 Hz, 2H), 7.56 (d, J=5 Hz, 2H). MS (DCI/NH₃) at m/e 560 (M+H)⁺. Anal calcd for C₃₂H₃₇N₃O₆•4.2 TFA: C, 46.72; H, 4.00; N, 4.05. Found: C, 46.66; H, 4.06; N, 4.00.

Example 533

trans, trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(3-trimethylammoniomethylphenyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

To a stirred solution of 0.128 gm of the compound of Example 532C in 0.5 ml methanol, 0.25 ml of an aqueous solution of trimethylamine was added. The mixture was allowed to stir at room temperature under nitrogen atmosphere for 4 hours. 1N HCl was added; the aqueous was washed with ether to extract organic impurities. The aqueous layer was dried azeotropically with toluene, and the residue was dried under high vacuum. Yield 0.115 gm. ¹H NMR (300 MHz, D₆-DMSO) δ 0.83 (t, J=7 Hz, 3H), 1.15-1.40 (m, 4H), 2.62 (s, 2H), 3.35 (s, 9H), 3.40-3.80 (m, 10H), 4.47 (s, 2H), 6.00 (s, J=3 Hz, 2H), 6.75-6.90 (m, 3H), 7.25-7.37 (m, 2H), 7.45-7.60 (m, 3H). MS (DCI/NH₃) at m/e 602 (M+H)⁺.

Example 534

(2R,3R,4S)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

Example 534A

Ethyl (3-fluoro-4-methoxy)benzoylacetate

Sodium hydride (17g of a 60% suspension in mineral oil) is washed three times with toluene. The powder is suspended in 138 mL of toluene, and 35 mL of diethyl carbonate is added. The mixture is heated to 90 °C, and a solution of 25 g of 3-fluoro-4-methoxyacetophenone and 50 ml of diethyl carbonate in 50 ml of toluene was added portionwise. Heating is continued for 30 min, then the reaction is cooled to room temperature. A solution of 50 ml of concentrated HCl in 75 ml of ice water is added slowly, and the mixture is stirred. The mixture is extracted with toluene; the combined organic extracts

are washed with brine and bicarbonate solutions. The product is dried over Na_2SO_4 and decolorized with charcoal to give 34.5 g (97%) of the title compound.

5

Example 534B

Ethyl 2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

- The compound of Example 534A (12.5 g) and 5-(nitrovinyl)-1,3-benzodioxole (13.1 g, 20% excess) were suspended in a mixture of 75 ml of THF and 13 ml of $i\text{PrOH}$. DBU (0.25 g) was added, and the mixture was stirred at room temperature for 30 min. An additional 0.1 g of DBU was added, and the solution was stirred for 1 hour. The solvents were removed in vacuo; toluene was added, along with brine containing 3 ml of concentrated HCl.
- The mixture was extracted twice with toluene; the organics were dried over MgSO_4 . The residue was flashed on silica, using CH_2Cl_2 to elute. Yield 75%. This material (17.4 g) is combined with 35 g of Raney Nickel (washed) in 250 mL of EtOAc. The mixture is shaken under 4 atm of hydrogen for 18 hours. The solution is concentrated in vacuo; the residue is chromatographed on silica, eluting with 4% EtOAc in CH_2Cl_2 . Yield 10.13 g = 66%. The product is combined with 26 ml of THF and 50 ml of EtOH; 2.18 g of NaBH_3CN are added, along with a trace of bromocresol green as indicator. A solution of 1:2 concentrated HCl/EtOH is added dropwise to maintain pH at green-yellow; after color persists, the reaction mixture is stirred for an additional 20 min. The solvents are removed in vacuo; the residue is stirred with mixture of toluene and KHCO_3 solution. The organic phase is washed with water and brine, and dried over MgSO_4 . The crude product is purified by flash chromatography on silica, eluting with 2:1 EtOAc/hexanes. Yield 5.92 g (58%) of a 2:1 mixture of trans-trans and cis-trans isomers.

30

Example 534C

Ethyl (2R,3R,4S)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

35

To the racemic amino ester above (15.0 g, 38.8 mmol), dissolved in 75 ml methylene chloride and cooled in an ice bath, was added Boc anhydride (9.30 g, 42.7 mmol). After stirring 2 hours at room temperature, the solution

was concentrated *in vacuo*; the residue was dissolved in 50 ml ethanol and treated with a solution of 3.75 g sodium hydroxide in 19 ml water. The solution was warmed until all was soluble. After stirring for 2 hours at room temperature, the solution was concentrated and redissolved in 200 ml of water. This was extracted with 75 ml of diethyl ether. The ether layer was extracted with 40 ml of water. The combined aqueous phases were acidified with 7.5 g acetic acid; the mixture was stirred until a solid formed. The solid was filtered, washed with water and dissolved in methylene chloride. After drying with sodium sulfate, the solution was concentrated and the residue crystallized from 1:1 ether:hexane to get 15.99 g of product, m.p. 200-203 (90% yield). The crude acid was suspended in 80 ml ethyl acetate and treated with 4.00 g (33.1 mmol) of (S)-(-)- α -methylbenzylamine. After heating to dissolve the acid, 80 ml of ether was added. Scratching with a glass rod caused the product to crystallize. The solids were filtered and washed with ether-ethyl acetate solution to give 8.22 g (81% yield based on 50% maximum recovery) of salt, m.p. 165-168°C. After one recrystallization, chiral HPLC analysis, using a Regis Whelk-O column, indicated >99.5 % e.e. The salt was dissolved in 500 ml of 36% HCl in ethanol; a white solid forms. The resultant suspension was heated for 16 hours at 52°C. After concentrating *in vacuo*, the residue was combined with toluene and stirred with potassium bicarbonate in water for 30 minutes. The toluene was separated, dried (Na₂SO₄) and concentrated. The residue was chromatographed on silica gel, eluting with 33% hexane-67% ethyl acetate to get 6.9 g (99%) of the resolved amino ester.

Example 534D

Ethyl (2R,3R,4S)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propylamino)ethyl)-pyrrolidine-3-carboxylate

The compound of Example 534C was dissolved in 1,2-dibromoethane (10 mL per 1 g of starting material); diisopropylethylamine (1 mL per 1 g of starting material) and NaI (100 mg per 1 g of starting material) were added, and the mixture was stirred at 100°C for 1 hour. Toluene was added, and the mixture was washed with bicarbonate. The solvents were concentrated, and the resultant black residue was chromatographed on silica gel, eluting with 4:1 hexane-EtOAc to give the N-(2-bromoethyl)pyrrolidine (85-92%). This compound was combined with n-propylamine (3.5 eq.) and NaI (10% by

weight of bromide) in ethanol (5 mL per 1 g of bromide), and was heated at 80°C for 2 hours. Toluene was added, and the mixture was washed with bicarbonate, dried (Na₂SO₄), and concentrated. More toluene was added, and removed *in vacuo*, to get rid of the primary amine. The residue was dissolved in heptane and filtered to remove a small amount of insoluble material. Evaporation of the solvent gave the desired product (86-93% yield), which was used for the next step without further purification.

Example 534E

1-Pentanesulfonyl chloride

1-Pentanesulfonic acid, sodium salt (10 g, 57.5 mmol) was charged into a 250 ml round bottom flask (allow headroom). Thionyl chloride (20 mL) is added; gas evolves, and a white solid forms. The mixture is heated at 60 °C for 3 hours. The solvents are removed in vacuo; toluene is added and removed in vacuo to remove residue of SOCl₂. The residue is partitioned between CH₂Cl₂ and ice water; the organic layer is dried over Na₂SO₄. The crude product is purified by distillation (bp 54-56 °C @ 0.5 mm Hg) to give a clear oil, 61% yield.

Example 534F

(2*R*,3*R*,4*S*)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(*N*-propyl-*N*-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 534D (200 mg, 0.43 mmol) was dissolved in 5 mL of CH₃CN; 110 mg (2 eq) of *N,N*-diisopropylethylamine and 72.8 mg (1.2 eq) of 1-pentanesulfonyl chloride were added sequentially, the resultant solution was allowed to stir at room temperature for 30 min. The solvent was evaporated under reduced pressure and the residue was dissolved in EtOAc. The solution was washed with saturated NaHCO₃ solution, 1N H₃PO₄, and brine, dried over Na₂SO₄ and evaporated to give a yellowish oil which was purified by flash chromatography on silica gel eluting with 40% EtOAc/hexane to give 220 mg of product (85%). This ester was dissolved in 5 mL of EtOH, to which was added NaOH (46 mg, 3 eq) solution in 2 mL of H₂O. This mixture was stirred for 3 hours at room temperature. The solution was concentrated in vacuo using low (<40°C) heat. Water (10 mL) and ether (50 mL) were added; the ether layer was extracted with 5 mL of water. The

combined aqueous mixture was back-extracted with ether and then neutralized with acetic acid. This solution was extracted twice with ether. The ether was dried (Na_2SO_4) and concentrated in vacuo. EtOAc (1 mL) and ether (1 mL) were added to dissolve the product, and hexane was added dropwise to produce a white solid. The solid was collected and dried in vacuo to give 125 mg of the title compound.

Example 534H

(2R,3R,4S)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid, hydrochloride salt



The free amine is dissolved in iPrOH; a slight excess of HCl in iPrOH is added, and the solution is concentrated in vacuo. More IPA is added, and the solution is re-concentrated. The resultant sticky

material is stirred with ether overnight to give a white powder, which is collected by filtration and dried overnight in vacuo at 60 °C. Yield 95%.


Example 535

The compounds in Table 3C may be prepared using methods presented in the above Examples.


Table 3C.




3




4




5




6




7




8




9




10




11




12




13




14




15




16



17




18




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100


19




20




21




22




23




24




25




26




5 27




28



29




30




10


31




32




33




34



35




36





5

37



38

39




41


10

40


453




42




43




44




45




46




47




48




49




50




51




52




53




54




55




56




57




58




59




60




61




62




63




64




65




66




67




68




69




70




71




72




73




74




75




76




77




78




79




10 80




81




82




83




84




85




86




87




88




89




90




91




92




93




94



95




96




97


460




98




99




100




101




102





103



104





105





106

107





108

109




10 110

111




462

112




113




463


114




115




5 116




117



118




119




10

120


121




122




123




124




125




126





127




128




129




5 130




131



132




133




10


134




135



136




137




467


138




139




140




141




142




143



144



145




138
139
140
141
142
143
144
145

5


10

468


146




147




5 148




149



150




151




10

152



153





CC(C)=C[C@H](C(=O)O)[C@@H]1CN(C(=O)NCCN(C)CC)CC[C@H]1C2=CC=C3C(=C2)OCOC3CN(C)CCCCCNC(=O)CCN1[C@H](Cc2cc3c(cc2)OCO3)c([C@@H]1C[C@H](C=C(C)C)C(=O)O)CCCN(CC)C(=O)NCC[C@H](C=C(C)C)[C@@H](C(=O)O)[C@H]1c2cc(OC)c3ccccc23O1CCN(CC)C(=O)CN1CC[C@H](C1)[C@H](C(=O)O)[C@H](C=C(C)C)[C@H]2C=CC(OC)OC2[illegible]CC(C)C=C[C@H]1[C@@H](C(=O)O)[C@H](C2=CC(OC)=C3C2OC3)[C@@H](CN(CCN(C)C)C(=O)Nc4ccc(cc4)CN(C)C)[C@H]1C

10



161

470





162

163



164



165




10

166


167



168




169




472


170




171




5 172




173



174




175




10


176




177



178




179




5


181




180



182



183





184


185




475



192




193




476


194




195




5 196




197



198




199




10


200




201



202




203




5


204




205



10 206




207




208

209


478




210




211




212




213




214



215




216




217


479




218




219




220




221




222



223



224



225

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

5

10

$$\begin{array}{ccccccc} \text{C}_2\text{F}_6 & \text{C}_2\text{F}_4 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 \\ \text{C}_2\text{F}_6 & \text{C}_2\text{F}_4 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 \\ \text{C}_2\text{F}_6 & \text{C}_2\text{F}_4 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 & \text{C}_2\text{F}_2 \end{array}$$

5

10


CCN(CC)C(=O)CN1CC[C@H](C1[C@@H](C(=O)O)[C@H](C2=CC(OC)=C[C@H]2OCCO)C3=CC=C(C=C3)OCCO)C4=CC=C(C=C4)F

229


CCCCN(CCCO)C(=O)CN1[C@H](C[C@@H]1C[C@H](C(=O)O)C2=CC(OC)=C3C=C[C@H](C2)OC3)C4=CC(OC)=C5C=C[C@H](C4)OC5

233


481




234




235




236




237




238



239



240




241


[illegible]

10


483




250




251




5 252




253




254



255




256




257


484




258




259




260




261




262



263




264




265


485




266




267




268




269




270




271




272




273




274



276




278




280


486




275



277




279




281


487




282




283




284




285




286



287




288




289

10


488




290




291




292




293




294



295




296




297

10


489




298




299




300




301




302



303




304




305


490




306




307




308




309




310



311



312




313


10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782
 783
 784
 785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796
 797
 798
 799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000

10


491




314




315




316




317




318



319




320




321


492




322




323




324




325




326



327




328





329

493





330

331





332

333




334

335




336


337




494




338




339




5 340




341




342



343




344




345


495




346




347




348




349




350



351



352



353



5

5

10


[illegible]

10




498


362




363




364




365




366




367



10 368




369




370


371




372




373




374




375




376




377




378




379




380




381




382





383



384




385




501


386




387




5 388




389



390




391




10

392




393




394

395


502




396




397




398




399



400





401




402

403


503



404




405




5

406




407




10

408




409




504


410




411




5 412




413



414




415



416


417

505




418


419




421



423




422




506


424




425



426




427




5

428



429





10


430

431


507



432




433




5


434




435



10 436




437




508


438




439




5 440




441




442




443



10 444




445




509

446




447




5


448




449



450




451



10


452

453




511


460




461




462




463




464




465



466



467




Chemical structure 460: A complex molecule featuring a 7-membered lactam ring connected via a chiral center to a 5-membered ring. This 5-membered ring is further connected to a 3,4-methylenedioxyphenyl group and a carboxylic acid group. The 5-membered ring is also connected to a side chain containing a tertiary amine and a carbonyl group.

5


10

512

468




469




5


470




471



10 472





473




474

475


513




476




477



478




479




480



481

[illegible]

5




492




10


494




489




491



493





495




CCCCN(CCCC)C(=O)CN1[C@H](Cc2c(c3cc(OC)c4c3OCO4)cc2C1)C[C@H](C1CCN(C)CC1)C(=O)OCCCCN(CCCC(=O)CN1CC[C@H](C1)C[C@@H](C2=CC3=C(C=C2)OC3)C(=O)O)C(=O)OCCN4CCCC4=OCCCN(CC)CC(=O)CCN1C[C@H](C[C@H](C1)C[C@@H](C2=CC(OC)=C3C=C[C@H](C3OC2)C(=O)O)C4CCN(CC)CC4=O)C5CCN(CC)C(=O)N(CCC)C(=O)N1CC[C@H](C1)C2=C(C(=C(C=C2)OC)OCO)C[C@H](C2)C(=O)OCCOC(=O)N(CC)C(=O)CN1CC[C@H](C1)[C@H](C(=O)O)[C@H](c2cc3c(cc2)OCO3)CCCCCN(CCCCNC(=O)CN1CC[C@H](C1Cc2cc3cc(OC)c4ccccc34O2)C(=O)O)C(=O)O

503


517



504




505




5

506




507




10

508




509




518

510




511




5


512




513



514



515





10


516

517


519




518




519




520




521



10 522




523




520

524




525




5


526




527



528




529




10

521


530




531




5 532




533



534




535




10


536




537



538




539




5


540




541



10 542




543




15 544

545


523




546




547




548





549



550






551







553

5




555




525

560




561




5


562




563



564



565





10


566

567


526




5 568




569




570




571



10 572




573




527


574




575



576




577




5

578




579




580

581


528




582




583




5 584




585



586




587




588


589




590




591




592





593



594




595




CCN(C)CCCCNC(=O)CN1Cc2ccc3c(c2)OCO3[C@H]1C[C@@H](C(=O)O)C[C@H](N1C)CCN1C(=O)N(C)C1=OCCCN(C(=O)CCN(Cc1ccc2c(c1)OCO2)c1ccc(OC)cc1)C(=O)c1ccccn1CCCN(C(=O)CCN1C[C@H](C[C@@H]1C[C@H](C2=CC(=C(C=C2)OC3COC3)C(=O)O)C4=CC=CC=C4C5C(=O)NCC5=O)C6=CC=CC=C6CCCN(C(=O)CN(Cc1ccc(N)cc1)C(=O)N2CC[C@H](C2)C[C@@H](C(=O)O)[C@H](C3=CC=C4C(=C3)OCOC4=O)C5=CC=CC=C5)CC6=CC=CC=C6C7=CC(=CC=C7)C(=O)N(C)C(=O)N6CCN(CC)Cc1ccc(cc1)C(=O)NCCN2[C@H](Cc3c(cc4c3OCO4)OC)C[C@@H](C(=O)O)C2Cc5cnc(=O)n5=OCCCCNC(=O)CN(Cc1cc2ccccc2o1)C[C@H](Cc3cnc(=O)n3C)C(=O)OCCN(CC)C(=O)C1=CC=C(C=C1)C(F)=C1C=CC(=C1)C2=CC=C3C=C(C=C2)C(=C3)OC4=CC=CC=C4[C@H]5C[C@@H](C(=O)O)N(CCN5C6=CC=CC=C6)CC7=CC=CC=C7CCCCN(CCCC(=O)N[C@@H]1C[C@H](Cc2c(c3ccccc3O2)C[C@H]1C(=O)O)C[C@H]1CCN(C(=O)O1)C(=O)O

531


604




605




606




607




608




609





610



611





532



612



613



5



614

615



616


617




10

533


618




619




620




621




622




623



624



625



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100


5

10


02
 03
 07
 07
 08
 07
 07
 04
 1
 02
 03
 07
 14
 03
 03

535


626




627




628




629




630




631




632




633



634




635




5


636




637



10 638





639




15 640

641


537




642




643




644




645




646




647




648




649




5 650




651



652





653




654

655


539




656




657




658




659



10 660




661




540

662




663




5


664




665



666



667





10



668

669



541





670



672



674




10

CCCCN(CCCC(=O)CN1CC[C@H](C1)[C@H](Cc2c(c3cc(OC)c4c3OCO4)cc2)C[C@@H](C(=O)O)N5CC[C@H](C5)C(=O)N6C(C)CC6CCN(CC)C(=O)CN1CC[C@H](C1)[C@H](Cc2c(c3cc(OC)c4ccccc34o2)O)C(=O)OCCN(C)CCCN(C)C(=O)CN(C)[C@H](Cc1ccc2c(c1)OCO2)C[C@@H](C(=O)O)C[C@H](C)N3CCC(=O)N3CCN(C)CCCC(=O)N(CC[C@H]1CC[C@@H](C1Cc2cc3c(cc2)OCO3)C)CC[C@H](C(=O)O)C4CC(C)CC4=OCCN(CCC)C(=O)N(CCN(C)C)C[C@H](C(=O)O)[C@@H](C1=CC=C2C(=C1)OCOC2)C3=CC(OC)=CC=C3CCN(C)CCCCN(C)C(=O)C1CN(C[C@H]1[C@@H](C(=O)O)[C@H](C2=CC(=C(C=C2)OC3OC4C(=C(C=C3)OC4=O)C)C)C5C(C)CC(=O)N5C)C


CCN(C)CCCCN(CCCC)C(=O)N[C@@H](Cc1ccc2c(c1)OCO2)C[C@H](C(=O)O)[C@@H](C)N3CCCC3CCCN(CCC)C(=O)CN(CCC1C(CCN1C(=O)O)C2=CC=C3C(=C2)OCOC3)C4=CC=CC=C4NCCCN(Cc1cccnc1)C(=O)N[C@@H](Cc1ccc2c(c1)OCO2)C[C@H](C(=O)O)N3CC[C@H](C3)C(=O)NCCN(CC)C(=O)N(CC1CC(C1)C(C2=CC(OC)=C3C2OC3)C(=O)O)CCN4C(=O)C(C(C4)C)CCCN(CC)C(=O)N(CC1=CC=C(C=C1)CN(C)C)C(=O)N2CC[C@H](C[C@@H](C2)C[C@H](C3=CC=C4C(=C(C=C3)OC4OC5=CC=CC=C5C)C)C(=O)O)CC6(C)CCN(C6)C(=O)OCCCCN(CCCC)C(=O)CN1Cc2cc3ccccc3oc2[C@H]1Cc1ccc(cc1)C(=O)O

691




545


698




699




5 700




701



702




703



10


704

705


[illegible]

547


712




713



714




715




5


716




717




718




719




720



722




724




726


721





723



725




727




550


728




729




5 730




731



732



733




10


734

735


551




736




737




5 738




739



740




741




10

742




743


552




744




745




5 746





747




748




749




750




751




752




753



754




755




10


756



757



5



CCCN(C(=O)C[C@H]1N[C@@H](Cc2c3ccccc2O3)[C@H](C(=O)O)[C@@H]1Cn1cnc(C)c1=O)c1cccnc1CCCN(C)C(=O)N(Cc1ccc2c(c1)OCO2)C(=O)N(Cc1ccc(N)cc1)C(=O)O

10


CN(C)CCCCN(CCCC)C(=O)CN[C@H](Cc1c[nH]c2c1=O)C[C@@H](C(=O)O)[C@H](c1ccc2c(c1)OCO2)NCCCN(C(=O)CN1[C@H](Cc2c3ccccc2O3)C[C@@H]1C(=O)O)C(=O)Nc4ccc[n+]4CCN(C)Cc1ccc(cc1)N(CCC)C(=O)CCN2C[C@H](C3=CC=C4C(=C3)OCOC4=CC=C3C(=O)O)[C@@H](C2)CCN5C(=O)N(C)CC5

765


555




5 766




767



768




769




10


770




771



772




773




5

774




775




557


776




777




5 778




779



780





781



782



783

558





784

785



786



787




10

788


789




790




791




5 792




793



794




795




10 796


797




798




799




5 800





801



802




803




561


804




805




806




807




5 808




809



810




811




562


812




813




814




815




5 816




817



818




819




563

820




822


821




823




5 824





825



826




827




564


828




829



830




831




5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100


5 832




833



834




835




565


836




837




838




839




5 840




841



842




843




10 844

845


566



847




846




567


848




849



850




851




5


852




853



10 854




855




568


856




857




5 858




859



860



861




10


862

863


569




864




865




5 866




867



868




869




10

570


870




871



872




873




5


874




875



876




877




10

571


878




879



880




881




5


882




883



884




885




10


886




887




5 888




889



890




891




10

892




893




CCN(C)CCCCNC(=O)C[C@H]1N[C@@H](Cc2c(c3ccccc23)OCO3)[C@H](C(=O)O)[C@H]1Cc4ccccc4N5CCCC5S(=O)(=O)O

574


896




897




5 898




899



900




901




10


902




903



904




905




5


906




907



10 908





909




910

911


[illegible]




919



921




923




578


924




925




5 926




927



928




929




10

930




931




579

932




933




5


934




935



10 936





937




938

939


580




940




941



942



943





945

CCCCN(CCCC)C(=O)CN1CC[C@H](C1Cc2cc3OCOc3cc2C)C(=O)OCC(C)C(C)C(C)C(=O)N(c1ccc(F)cc1)C(=O)CN2C[C@H](C[C@@H](C2)C[C@H](C3=CC=C4C(=C3)OCOC4=CO)C(=O)O)C[C@H](C2)CN(C)C(=O)CCCCCN(CCCCOC1=CC=C2C(=C1)OCOC2)CC(=O)N[C@@H](C[C@H](C)C(=O)NCC)C[C@H](C(=O)O)CCCOC(=O)N(CC)C[C@H]1CN(C[C@@H]1Cc2cc3c(cc2)OCO3)c4ccc(OC)cc4CCN(C)C(=O)N[C@H]1C[C@@H](C(=O)O)[C@H](c2cc3c(cc2)OCO3)C[C@H]1C(=O)N(CC)CCNCCN(C)C(=O)C[C@H]1CN(C(=O)NCCCCN)C[C@@H]1[C@H](C(=O)O)[C@H]2C=CC(OC)OC2CCN(C)CCCN(C)C(=O)CN[C@H](Cc1ccc2c(c1)OCO2)C[C@H](C(=O)O)CN(C)C(=O)CCCN(C)CCCC(=O)CCN1[C@H](C[C@@H]1[C@H](C(=O)O)C2=CC=C3C=C(C=C2)OC3)C4=CC=CC=C4OC5=CC=CC=C5


CCN(C)C(=O)C[C@H](C(=O)O)[C@@H](Cc1ccc2c(c1)OCO2)c1ccc(OC)cc1CCN(C)CCCCN(C)C(=O)CCN1C[C@H](C[C@@H](C1)C(=O)O)[C@H](C2=CC3=C(C=C2)OCOC3)CCCN(C)CCCCNC(=O)CCN1C[C@H](C[C@@H]1C(=O)O)[C@H](C2=CC(=C3C=C(C=C2)OC3OC2)C)CCCN(CC)C(=O)CCN(C[C@H]1C[C@@H](C[C@H]1C[C@@H](C2=CC(=C(C=C2)OC3C=CC(OC3OC4C=CC=C4)C=C2)C(=O)O)C5=CC=CC=C5)C(=O)Nc6ccccn6CCN(CC)C(=O)NCCN(C(=O)O[C@H]1Cc2cc(OC)c3ccccc23O1)[C@@H](C(=O)O)CCN(CC)C(=O)NCCN(C(=O)O)[C@H]1Cc2cc(OC)c3ccccc23O1CCN(CC)C(=O)N(CC)C(=O)N1CC[C@H](C1)C2=CC=C3C(=C2)OC3C=C(C=C3)OC

959


583




960




961



962




963




10


964




965



966




967




585


968




969




5 970




971



972




973




10


974



975




976




5

978




10

980




15 982


977





979



981




983




588


984




985




5 986




987



988




989




10

990




991




50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

590


992




993




5 994




995



996




997




10


998



999




1000




5


1001




1002




1003



1004




1005




1006

1007


592




1008




1009




1010




1011




1012




1013




1014




1015



1016




1017




5

1018




1019




10

1020




1021




594

1022




1023




5


1024




1025



10 1026




1027




1028

1029


595



1031




1030




596


1032




1033




5 1034




1035



1036



1037




10


1038

1039


597




1040




1041




1042




1043




1044



1045




1046




1047


598




1048




1049




1050




1051




1052




1053



1054




1055




5


1056




1057



10 1058





1059




1060

1061


600



1062




1063




601


1064




1065




5 1066




1067



1068



1069





10

1070



1071

602





1072

1073





1074

1075



1076

1077




1072
1073
1074
1075
1076
1077
1078
1079

5


10

603


1078




1079




5 1080




1081



1082




1083




1084

1085




605


1092




1093




5 1094




1095



1096




1097




10


1098




1099



1100




1101




5

1102




1103




10

1104





1105




1106

1107


607




1108




1109



1110




1111




608


1112




1113




1114




1115




1116




1117




1118




1119



1120




1121




5

1122




1123




10

1124





1125



1126



1127

610





1128

1129





1130

1131



1132

1133




1128
1129
1130
1131
1132
1133
1134
1135


5

10


1134




1135




5 1136




1137



1138




1139




10


1140




1141



1142




1143




5


1144




1145



10 1146




1147




1148

1149


613




1150




1151



1152




1153




1154

1155


614




1156




1157




1158




1159




1160



1161





1162





1163

615





1164

1165





1166

1167




1168

1169




616


1170




1171




5 1172




1173



1174




1175




10 1176


1177




1178




1179




1180





1181



1182




1183




618


1184




1185




5 1186




1187



1188



1189




10


1190

1191


619




1192




1193




1194





1195




1196




1197




1198




1199




1200




1201




1202




1203




1204




1205




1206




1207




1208




1209




5 1210




1211



1212




1213




10


1214




1215



1216




1217




5


1218




1219



10 1220





1221




1222

1223




625


1224




1225




5 1226




1227



1228




1229




10

1230




1231




627


1232




1233




5 1234




1235




1236




1237




1238




1239



1240




1241




5


1242




1243



10 1244





1245



1246



1247

629





1248

1249





1250

1251



1252

1253




1247
1248
1249
1250
1251
1252
1253
1254


5

10

1254




1255




631


1256




1257




5 1258




1259



1260





1261



1262



1263

632





1264

1265





1266

1267



1268

1269




1263
1264
1265
1266
1267
1268
1269
1270

5


10

633


1270




1271




5 1272




1273



1274




1275




1276

1277


634




1278




1279




1280





1281



1282




1283




635


1284




1285




5 1286




1287



1288




1289




10


1290




1291



1292




1293




5

1294




1295




10

1296





1297




15 1298

1299


637




1300




1301



1302




1303




638


1304




1305




5 1306




1307



1308




1309




10


1310




1311



1312




1313




5


1314




1315



10 1316




1317



1318


1319




CCCCN(CCCCNC(=O)CNC1Cc2cc3ccccc3oc21)C(=O)N1CCC(=O)N1C(=O)OCCN(C)CCCCN(CCCC)C(=O)CN(CCC1=CC=C2C(=C1)OC2)C[C@H](C(=O)O)CC3=CC(=O)N(C3=O)C=O

642


1328




1329




5 1330




1331



1332




1333




10


1334




1335




1336




1337




5 1338




1339



1340





1341




1342

1343


644




1344




1345




1346




1347



1348



1349




1343
1344
1345
1346
1347
1348
1349
1350


5

10

1350




1351




646


1352




1353




5 1354




1355



1356




1357




1358


1359




1360




1361




1362




1363



1364




1365




1366


1366




1367




5 1368




1369



1370





1371




1372

1373


649



1374




1375




650


1376




1377




5 1378




1379



1380



1381




10


1382

1383


651




1384




1385




1386




1387




10 1388




1389




1390




1391




5 1392




1393



1394




1395




1396

1397


653



1398




1399




654


1400




1401




5 1402




1403



1404




1405




1406

1407


655




1408




1409




1410





1411



1412




1413




5

656


1414




1415




1416




1417



1418




1419




1420

1421


657




1422




1423




1424




1425



1426




1427




CN(C)CCCN(C)C(=O)CN[C@H](Cc1ccc2c(c1)OCO2)[C@@H](Cc1cccnc1)C(=O)OCC[N+](C)(C)CCCN(C)C(=O)CC[C@H]1CN(C[C@@H]1Cc2cc3occc3cc2)Cc4cccnc4C(=O)OCCCCCN(CCCC)C(=O)N[C@@H](Cc1ccc2c(c1)OCO2)[C@H](Cc1cncn1)C(=O)OCCN(C)CCCCN(CCCC)C(=O)N[C@@H](Cc1ccc2c(c1)OCO2)[C@H](Cc1cccnc1)C(=O)OCCN(C)CCCCN(CCCC)C(=O)N1CC[C@H](C1Cc2cc3ccccc3o2)C[C@H](C1=CC=CC=C1)C(=O)OCCCN(C(=O)CN2[C@H](Cc3c[nH]cn3)[C@@H](C4=CC5=C(C=C4)OCO5)C2)Cc6ccccn6CCCN(Cc1cccnc1)C(=O)CN2[C@H](Cc3cccnc3)[C@@H](C(=O)O)[C@H](C4=CC5=C(C=C4)OCO5)N2CCN(CC)C(=O)N(CC1=CC=C(C=C1)CN)CC2CN(C2[C@H](C3=CC4=C(C=C3)OCOC4)C[C@@H](C5=CC=NC=C5)C)C(=O)O

1436




1437




5


1438




1439



10 1440








1441




1442

1443




661


1448




1449




5 1450




1451



1452




1453




1454

1455


662




1456




1457




1458




1459




1460



1461





1462




1463

CCN(C)CCCCN(C)C(=O)N(CCN(C)C[C@H](Cc1ccc2c(c1)OCO2)[C@@H](C(=O)O)c3ccncc3)CCCN(C(=O)Nc1cccnc1)CNC(Cc2cccnc2)C3Cc4ccccc4O3C(=O)OCCCN(C(=O)CN[C@@H]1[C@H](Cc2c(c3ccccc2O3)CO)C[C@H]1Cc4cccnc4)[C+]1=CC=CC=C1CCN(CC)C(=O)N[C@@H](Cc1ccc2c(c1)OCO2)[C@H](Cc1cncn1)CC(=O)OCCN(C)CC1=CC=C(C=C1)N(CCC)C(=O)CCN2C[C@H](C[C@@H](C2)c3cc4c(cc3)OCO4)CC[C@H](C2)c5ccnnc5CCCCN(CCCC)C(=O)CN1[C@H](Cc2c(c3cc4c(c3)OCO4)ccccc2C(=O)O)[C@H](C1)C(=O)N1CCCC1


664



1470




1471




[illegible]CCCCN(CCCCNC(=O)CN1CC[C@H](C1)[C@H](C(=O)O)[C@@H](C1=CC=C2C(=C1)OCOC2)C3=CC=CC=C3)C(=O)OCN(C)CCCCNC(=O)CN1CC[C@H](C1)[C@H](C(=O)O)[C@@H](C1=CC=C2C(=C1)OCOC2)C3CCCC3=OCCN(C)CCCC(=O)N[C@@H]1CC[C@H](C1Cc2cc3ccccc3o2)C[C@H](C1CCNC1=O)C(=O)OCCCCNCCCNC(=O)CN[C@H](Cc1c[nH]c1)[C@H](Cc2c3ccccc3oc2)[C@@H](C(=O)O)NCCN(C)CCCCN(CCCC(=O)N1CC[C@H](C1Cc2cc3ccccc3o2)C[C@H](C1CCC(=O)N1)C(=O)O)C(=O)OCCN(C)CCCCN(CCCC)C(=O)CN1Cc2cc3c(cc2O1)OCO3[C@H]1C[C@@H](C(=O)O)N1Cc2ccccc2CCCN(C(=O)CN1[C@H](Cc2cc3ccccc2O3)C[C@@H](C(=O)O)[C@H]1N2CCCC2=O)c3cc[n+](c3)C


1480




1481




1482




1483




1484




1485



1486




1487




668


1488




1489




1490




1491




1492




1493




1494




1495




1496




1497




5 1498




1599



1500




1501




10

1502




1503




15

671


1504




1505




1506




1507




1508




1509



1510




1511

[illegible]


5

10


1512




1513




5 1514




1515



1516




1517




10

1518




1519




673


1520




1521




1522




1523




1524




1525



1526



1527




1520
 1521
 1522
 1523
 1524
 1525
 1526
 1527

5


10

674


1528




1529




5 1530




1531




10 1532




1533



1534




1534




62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80


1536




1537




5 1538




1539



1540




1541




10

1542




1543




15

678


1544




1545




1645




1647




1548




1549



1550




1551




679


1552




1553




5 1554




1555



1556




1557




10

1558




1559




681


1560




1561




5 1562




1563



1564



1565




10


1566

1567


$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) = \frac{\partial L}{\partial x}$	$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{y}} \right) = \frac{\partial L}{\partial y}$
$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{z}} \right) = \frac{\partial L}{\partial z}$	$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\theta}} \right) = \frac{\partial L}{\partial \theta}$
$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\phi}} \right) = \frac{\partial L}{\partial \phi}$	$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\psi}} \right) = \frac{\partial L}{\partial \psi}$




1568




5




1572




10




1569



1571




1573




CCCCN(CCCC(=O)N[C@@H]1[C@H](Cc2c(c3cc(OC)c4c3OCO4)cc2C(=O)O)C[C@H]1CN1CCCC1=O)CCOCCOC(=O)C(C(=O)O)N(CCC1CCCC1=O)C2Cc3cc(OC)c4ccccc34O2

684


1576




1577



1578




1579




5

1580




1581




10

1582




1583




685

1584




1585




5


1586




1587



10 1588




1589




1590


1591




1592




1593




1594




1595




1596




1597




1600




1601




1602




1603




1604



1605




1606




1607







1608




1609




1610





1611




1612




1613




1614




1615



1616




1617




5


1618




1619




10 1620




1621




1622




1623




1624




1625




1626




1627




10 1628




1629




1630




1631




5 1632




1633



1634




1635




10

1636


1637




1638




1639




1641



1642




1643




694


1644




1645




1646



1647




1648




695


1649




1650




1651



1652




1653




1654

1655


696




1656




1657




1658




1659




1660




1661




1662




1663




1664




1665




1666




1667




1668




1669



1670




1671




699


1672




1673




1674




1675



1676




1677




1678

1679


700




1680




1681




1682




1683




1684




1685




1686




1687




1688




1689




1690




1691



10 1692




1693




CCCCN(CCCC(=O)N[C@@H]1C[C@H](Cc2c3ccccc2O3)C[C@H]1Cc4c5ccccc4O5)C(=O)OCCOC(=O)CN(C[C@H](C1=CC=C2C=C(C=C1)OCO2)C[C@@H](C3=CC=C4C=C(C=C3)OCO4)C(=O)O)CC

703


1696




1697




5 1698




1699



1700




1701




1702

1703


704




1704




1705




1706





1707




1708




1709




1710




1712




1714




1716




1711




1713




1715




1717



1718




1719




707


1720




1721




1722




1723




1724




1725



1726




1727




1720
1721
1722
1723
1724
1725
1726
1727

1728




1729




5


1730




1731



10 1732





1733




1734


1735




1736




1737




1738




1739




10 1740



1741




1742





710



1743





1745





1747



1749




1751




713


1752




1753




5 1754




1755



1756




1757




10


1758




1759



1760




1761




5

1762




1763




10

1764





1765




15 1766

1767




716


1768




1769




5 1770




1771



1772




1773




10

1774




1775




00
12
074
077
13
070
077
00
" "
00
00
00
00
00
00
00

718

1776




1777




5


1778




1779



10 1780



1781



Example 536

(2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(*N*,*N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

15

Example 536A

Ethyl 5,5-dimethyl-3-oxooctanoate

Ethyl 3,3-dimethylhexanoate was prepared using the general procedure of Cahiez *et al.*, *Tetrahedron Lett.*, **31**, 7425 (1990). To a solution of

63.8 g (370 mmol) of this compound in 400 mL of ethanol, cooled to 0°C, was added a solution of 30 g of NaOH in 150 mL of water. The resultant solution was warmed to ambient temperature and stirred overnight. Solvents were removed in vacuo; the residue was taken up in 700 mL of water, and extracted twice with 1:1 ether/hexanes. The aqueous layer was acidified to pH3 with 1N HCl and extracted twice with hexanes. The combined hexane extracts were washed with brine, dried over sodium sulfate, filtered and concentrated. A 20.2 g (150 mmol) sample of the crude product is dissolved in 150 mL of THF; 27.3 g of 1,1'-carbonyldiimidazole is added portionwise, to control gas evolution. In meantime, 33.4 g of potassium ethylmalonate and 13.4 g of magnesium chloride are combined in 350 mL of THF (overhead mechanical stirring) and warmed to 50°C for 3 hrs. This mixture is cooled to ambient temperature, and the above acid imidazolid solution is added. The resultant slurry is stirred overnight. Ether (600 mL), hexanes (600 mL) and aqueous 1N phosphoric acid (500 mL) are added, and the mixture is stirred for 30 min. The aqueous layer is separated; the organics are washed sequentially with bicarb (2X), water and brine. The organics are dried over sodium sulfate, filtered and concentrated to give 30.2 g (95% yield) of a colorless liquid.

Example 536B

4-Methoxy-6-(2-nitrovinyl)-1,3-benzodioxole

3-Methoxypiperonal (50.0 g) is combined with 71.9 mL of nitromethane in 250 mL of acetic acid; 36 g of ammonium acetate is added, and the mixture is heated to 50°C for 4 hrs. Solvents are removed in vacuo; the residue is taken up in water and stirred for 20 min. The solution is filtered; the filtrate is washed with water, then ether, to give 51.8 g of a yellow solid.

Example 536C

Ethyl *trans, trans*-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The compound of Example 536A (6.42 g, 30 mmol) was combined with 5.79 g of the compound of Example 536B in 40 mL of THF. DBU (0.5 mL) was added, and the mixture was stirred at ambient temperature for 6 hrs, during which time it turns reddish brown, and homogeneous. The solvents were removed in vacuo; the residue was taken up in EtOAc and washed

sequentially with aqueous 1N phosphoric acid and brine. The organic phase was dried over sodium sulfate, filtered and concentrated. The residue was dissolved in 50 mL of THF; 12 g of Raney Nickel catalyst (washed sequentially with water and ethanol) was added, followed by 10 mL of acetic acid. The resultant mixture was hydrogenated under 4 atmospheres of hydrogen until hydrogen uptake ceased (~ 3 hrs). The catalyst was removed by filtration; solvents were removed in vacuo. The residue was dissolved in 90 mL of 2:1 ethanol/THF; 30 mg of bromocresol green indicator was added, followed by 30 mL of 1N sodium cyanoborohydride in THF. Concentrated HCl was added dropwise to maintain pH at the indicator point, over 1 hr. The resultant solution was stirred overnight at ambient temperature. Bicarb was added, and the solvents were removed in vacuo; the residue was partitioned between water and EtOAc. The organic material was washed with water (2X) and brine. The organic phase was dried over sodium sulfate, filtered and concentrated. The crude product was dissolved in 100 mL of acetonitrile; 10 mL of Hünig's base was added, and the solution was warmed to 40°C overnight. Removal of solvents in vacuo provided 5.0 g of a yellowish oil.

Example 536D

Ethyl (2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The crude compound of Example 536C (2.0 g) was combined with 4 mL of triethylamine in 40 mL of THF; 2.0 g of di-*tert*-butyldicarbonate was added, and the mixture was stirred at ambient temperature for 5 hrs. Solvents were removed in vacuo, and the residue was taken up in 60 mL of ethanol. Aqueous sodium hydroxide (10 mL of 2.5 N solution) was added, and the resultant solution was stirred overnight. Solvents were removed in vacuo; the residue was taken up in water and extracted with ether. The aqueous phase was acidified with aqueous 1N phosphoric acid and extracted with EtOAc. The organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to give 1.0 g of a colorless oil. A sample of this material (0.734 g, 1.58 mmol) was combined with 0.35 g of pentafluorophenol and 0.364 g of EDAC in 5 mL of DMF. The resultant solution was stirred at ambient temperature for 1 hr, then was poured onto 50 mL of 0.6M sodium bicarbonate solution and extracted (3 X 15 mL) with ether. The combined ether extracts were washed with brine, dried over magnesium sulfate, filtered,

and concentrated in vacuo to give a foam, which was dissolved in 5 mL of THF and cooled to 0°C. Simultaneously, 0.418 g (2.37 mmol) of R-4-benzyl-2-oxazolidinone was combined with ~0.1 mg of pyreneacetic acid in 5 mL of THF and cooled to 0°C. N-butyllithium (1.6M in hexanes) was added to a red endpoint (persists ~10 sec), and the solution was stirred for 10 min. The solution was transferred into the solution of the pentafluorophenyl ester, and the resultant solution was stirred at 0°C for 40 min. Solvents were removed in vacuo; the residue was taken up in bicarb and extracted with ether (3 X 10 mL). The combined ether extracts were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The crude mixture of diastereomeric products was separated by flash chromatography on silica gel, eluting with a gradient from 4:1->3:1->2:1 hexanes/EtOAc, giving 423 mg of the faster-moving and 389 mg of the slower-moving diastereomer, respectively. The faster-moving diastereomer was dissolved in 2 mL of a 2.0M solution of sodium methoxide in methanol (freshly prepared, containing 5% methyl formate by volume) and stirred at ambient temperature for 16 hrs. Solvents were removed in vacuo, and the residue was partitioned between ether and aqueous 1N sodium hydroxide. The ether layer was washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The residue was purified by flash chromatography on silica gel, eluting with 4:1 hexanes/EtOAc. The resultant material was dissolved in 5 mL of TFA and stirred at ambient temperature for 1 hr. Solvents were removed in vacuo; the residue was suspended in bicarb and extracted with EtOAc. The organic phase was washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo to give 98 mg of product.

Example 536E

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 536D (48 mg) was combined with 35 mg of the compound of Example 501A in 3 mL of acetonitrile; 0.5 mL of Hünig's base was added, and the solution was allowed to stir overnight at ambient temperature. Solvents were removed in vacuo; the residue was partitioned between EtOAc and aqueous 1N phosphoric acid. The organic layer was washed with bicarb and brine, then dried over sodium sulfate, filtered and concentrated. The residue was purified by flash chromatography on silica

gel, eluting with 2:1 hexanes/EtOAc. The product was dissolved in 4 mL of ethanol; 1 mL of 2.5N aqueous sodium hydroxide was added, and the resultant solution was stirred overnight at ambient temperature. Solvents were removed in vacuo; the residue was taken up in water and extracted with ether. The aqueous phase was acidified to pH 3 with aqueous 1N phosphoric acid and extracted with EtOAc. The organic extracts were washed with brine, dried over sodium sulfate, filtered and concentrated to give a colorless oil. Lyophilization from acetonitrile/0.1% aqueous TFA gave 56 mg of a white solid.

^1H NMR (CDCl_3 , 300 MHz) δ 0.81 (s, 3H), 0.84 (s, 3H), 0.86 (t, $J = 6.9$ Hz, 3H), 0.93 (t, $J = 6.9$ Hz, 3H), 0.96 (t, $J = 6.9$ Hz, 3H), 1.09-1.38 (m, 8H), 1.45-1.59 (m, 4H), 1.84-2.00 (m, 2H), 3.15 (dd, $J = 6.9$ Hz, 10.0 Hz, 2H), 3.30-3.42 (m, 3H), 3.72 (t, $J = 10.5$ Hz, 1H), 3.86 (t, $J = 10.5$ Hz, 1H), 3.88 (s, 3H), 4.02 (q, $J = 10.0$ Hz, 1H), 4.12 (d, $J = 16.8$ Hz, 1H), 4.29 (d, $J = 16.8$ Hz, 1H), 4.41 (brn, 1H), 5.94 (s, 1H), 6.52 (d, $J = 1.8$ Hz, 1H), 6.67 (d, $J = 1.8$ Hz, 1H). MS (ESI) $(\text{M}+\text{H})^+$ at m/e 533. Anal calcd for $\text{C}_{30}\text{H}_{48}\text{N}_2\text{O}_6 \cdot 0.7$ TFA: C, 61.57; H, 8.01; N, 4.57. Found: C, 61.59; H, 8.20; N, 4.63.

Example 537

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 537A

Ethyl *trans, trans*-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

Prepared according to the procedures of Example 536C above, substituting the compound of Example 501B (5-(2-nitrovinyl)-1,3-benzodioxole) for 4-methoxy-6-(2-nitrovinyl)-1,3-benzodioxole.

Example 537B

Ethyl (2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The compound of Example 537A (6.8 g) was dissolved in 100 mL of ether; a solution of 1.6 g of (S)-(+)-mandelic acid in 60 mL of ether was added, the total volume was made up to ~200 mL, and the solution was seeded. The mixture was stirred slowly overnight. The resultant crystals were collected by

filtration and recrystallized from ether/EtOAc to give 1.8 g of a white solid. This material was partitioned between bicarb and ether; the ether layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo to give the enantiomerically pure product (>98% e.e.).

Example 537C

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

- 5 Prepared from the compound of Example 537B according to the procedures of Example 536E. ¹H NMR (CDCl₃, 300 MHz) δ 0.80-0.99 (m, 15H), 1.10-1.37 (m, 8H), 1.43-1.58 (m, 4H), 1.77-1.97 (m, 2H), 3.48-3.12 (m, 5H), 3.60-3.69 (m, 1H), 3.75-3.86 (m, 1H), 3.95-4.16 (m, 2H), 4.28-4.4 (m, 2H), 5.94 (s, 2H), 6.74 (d, J=7.8 Hz, 1H), 6.8 (dd, J=8.1, 1.5 Hz, 1H), 6.87 (d, J=1.8 Hz, 1H). MS (APCI+) m/e 503 (M+H)⁺.

Example 538

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

15

Example 538A

N-Boc-N-butyl-O-allylhydroxylamine

- 20 (15 mL). The solution was cooled to 0°C in an ice bath. Diisopropylethylamine (8mL) and di-*t*-butyldicarbonate (10.0g) were added. The mixture was stirred at 0°C for one hour at which point the bath was removed and the reaction allowed to warm to room temperature and stirred overnight. The THF was removed *in vacuo* and the residue taken up in EtOAc (25 mL), and washed
- 25 with water (1 x 50 mL), saturated sodium bicarbonate solution (3 x 50 mL), 1N phosphoric acid (3 x 50 mL), and brine (1 x 50 mL). The organic layer was dried with sodium sulfate and evaporated to give a light yellow oil (6.5g). This crude product was dissolved in dry THF (25 mL) and the solution cooled to 0°C in an ice bath. Sodium hydride (1.5g, 60% dispersion in oil) was added portionwise
- 30 over five minutes. The resulting mixture was stirred for 30 minutes at 0°C. 1-Iodobutane (4.1 mL) was added dropwise to the mixture. The reaction was stirred at 0°C for one hour, then stirred overnight at room temperature. The THF was removed *in vacuo* and the residue taken up in EtOAc (50 mL) and washed with water (1 x 50 mL), saturated sodium bicarbonate solution (3 x 50
- 35 mL), 1N phosphoric acid (3 x 50 mL), and brine (1 x 50 mL). The organic layer was dried with sodium sulfate and evaporated to give a light yellow oil, which

was purified by flash chromatography on silica gel eluting with 5% EtOAc/hexanes to give the title compound as a colorless oil (6.0 g).

Example 538B

N-butyl-N-propoxyamine trifluoroacetate

The compound of Example 538A (6.0 g) was dissolved in EtOAc (100 mL). 10% Palladium-on-carbon (0.5 g) was added, and the mixture was purged with nitrogen. The nitrogen line was exchanged for a balloon of hydrogen, and the mixture was stirred at room temperature for 6 hours. The catalyst was removed by filtration through a pad of Celite and the solvents were removed *in vacuo* to give a yellow oil which was purified by flash chromatography on silica gel eluting with 5% EtOAc/hexanes to give a colorless oil (5.8 g). A sample of the resultant material (1.15 g) was dissolved in CH₂Cl₂ (5 mL) and cooled in an ice bath. Trifluoroacetic acid (3mL) was added and the solution stirred cold for two hours. The solvent was removed *in vacuo*, care being taken not to allow the solution to warm above room temperature. The residue contained considerable TFA and was used without further purification.

Example 538C

N-butyl-N-propoxy-bromoacetamide

The salt of Example 538B (0.60 g) was dissolved in acetonitrile (5 mL) and cooled to -20°C. Hünig's base (5.5 mL) was added slowly. Bromoacetyl bromide (0.5 mL) was added dropwise over five minutes. The solution was stirred at -20°C for 30 minutes. The bath was removed and the solution was stirred for six hours at room temperature. The solvent was removed *in vacuo* and the residue taken up in EtOAc (50 mL) and washed with water (1 x 25 mL), 1N phosphoric acid (3 x 25 mL), and brine (1 x 25 mL). The organic layer was dried with sodium sulfate and evaporated to give a dark orange oil (0.65 g) which was used without further purification.

Example 538D

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 537B was reacted with the compound of Example 538C according to the procedures of Example 536E.

Example 539

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-propyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 539A

N-propyl-N-propoxy bromoacetamide

Prepared according to the procedures of Example 538A-C, substituting iodopropane for iodobutane in Example 538A.

Example 539B

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-propyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 540

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 537B was reacted with the compound of Example 539A according to the procedures of Example 536E.

Example 541

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-propyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 536D was reacted with the compound of Example 539A according to the procedures of Example 536E.

Example 542

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 542A

trans-Ethyl 3,3-dimethyl-4-hexenoate

A mixture of 4-methyl-3-penten-2-ol (7.4 g, 74 mmol), triethyl orthoacetate (13.6 mL, 74mmol) and propionic acid (0.28 mL, 3.7 mmol) was heated at 150°C for 7 hr. The product was then distilled under normal pressure (200-220 °C) to give 5.0 g of crude ester as a colorless oil.

Example 542B

Ethyl trans,trans-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The title compound is prepared according to the procedures of Examples 536A and 536C, substituting the compound of Example 542A for ethyl 3,3-dimethylhexanoate in Example 536A and the compound of Example 501B (5-(2-nitrovinyl)-1,3-benzodioxole) for 4-methoxy-6-(2-nitrovinyl)-1,3-benzodioxole in Example 536C.

Example 542C

Ethyl (2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The compound of Example 542B was resolved according to the procedure described in Example 537B.

Example 542D

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 542C was reacted with the compound of Example 538C according to the procedures of Example 536E.

5

Example 543

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-propyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

10

The compound of Example 542C was reacted with the compound of Example 539A according to the procedures of Example 536E.

Example 544

15

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 544A

20

Ethyl *trans,trans*-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The title compound is prepared according to the procedures of Examples 536A and 536C, substituting the compound of Example 542A for ethyl 3,3-dimethylhexanoate in Example 536A.

25

Example 544B

Ethyl (2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

30

The compound of Example 544A was resolved according to the procedure described in Example 536D.

Example 544C

35

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-butyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 544B was reacted with the compound of Example 538C according to the procedures of Example 536E.

Example 545

(2S,3R,4S)-2-(2-(2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-propoxy, N-(n-propyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 544B was reacted with the compound of Example 539A according to the procedures of Example 536E.

Example 546

(2S,3R,4S)-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-((N-4-heptyl-N-(2-methyl-3-fluorophenyl)) amino carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 546A

Ethyl trans,trans-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The title compound is prepared according to the procedures of Examples 536A and 536C, substituting the compound of Example 519A for 3,3-dimethylhexanoic acid in Example 536A.

Example 546B

Ethyl (2S,3R,4S)-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate

The compound of Example 546A (1.5 g) was dissolved in CH₂Cl₂ (25 mL). Di-*t*-butylidicarbonate (0.9 g) was added and the solution stirred overnight at room temperature. The solvent was evaporated *in vacuo* and the residue taken up in EtOAc (50 mL), washed with water (1x50 mL), saturated sodium bicarbonate solution (3x50 mL), and brine (1x50 mL). The organic layer was dried with sodium sulfate and evaporated *in vacuo* to give an oil with was purified by flash chromatography on silica gel eluting with 1/10/10 EtOH/EtOAc/hexanes to give a colorless oil (1.5 g). The oil was dissolved in EtOH (10 mL) and 50% NaOH solution (0.5 mL) and water (5 mL)

were added. The mixture was stirred overnight at room temperature. The solvents were evaporated *in vacuo* and the residue taken up in EtOAc (25 mL) and acidified with 1 N H₃PO₄ (10 mL). The layers were separated and the organic layer dried with sodium sulfate and evaporated to give a white semi-solid (1.3 g). A sample of the resultant Boc-protected amino acid (0.9 g) was dissolved in DMF (5 mL). (S)-Phenylalaninol (0.32 g), HOObt (0.33 g), and EDCI (0.40 g) were added and the solution stirred overnight at room temperature. Water (50 mL) was added and the mixture extracted with EtOAc (3x25 mL). The organic layers were combined, washed with water (2x50 mL), saturated sodium bicarbonate solution (3x50 mL), and brine (1x50 mL), and evaporated to give a yellow oil; tlc indicated the presence of two diastereomeric products. The diastereomeric amides were separated by flash chromatography on silica gel eluting with 1/12/12 EtOH/EtOAc/hexanes to give faster- (450 mg) and slower-moving isomers (400 mg). The faster-moving diastereomer (400 mg) was taken up in 6N HCl and heated at reflux overnight. The solvent was evaporated and the residue was taken up in toluene (75 mL) and evaporated. This was repeated two additional times to give a brown solid, which was dissolved in EtOH (50mL). 4N HCl/dioxane (10 mL) was added and the solution heated at reflux overnight. The EtOH was evaporated and the residue taken up in EtOAc which was treated with saturated sodium bicarbonate solution (3x50 mL), and brine (1x50 mL), and evaporated to give a brown solid. Flash chromatography on silica gel eluting with 30% EtOH/EtOAc gave a mixture of products (130mg) which was approximately 70% desired material. This product was carried forward without additional purification.

Example 546C

(2S,3R,4S)-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-((N-4-heptyl-N-(2-methyl-3-fluorophenyl)) amino carbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 546B was reacted with the compound of Example 508E according to the procedures of Example 536E.

Example 547

(2S,3R,4S)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-

carboxylic acidExample 547AN-butyl-4-hydroxybutyramide

To 30 mL (390 mmol) of γ -butyrolactone was added 45 mL (455 mmol) of *n*-butylamine. The solution was heated at 85°C for 1.5 hr, then the excess *n*-butylamine was removed *in vacuo*. The product crystallized on standing to give about 62 g of a colorless, low melting solid.

Example 547BN-butyl-4-hydroxybutyl chloroacetamide

To an ice cooled solution of 3.40 g (91.9 mmol) of LiAlH_4 in 90 mL of THF was added 2.4 mL of 98% H_2SO_4 , dropwise, with stirring. After bubbling had ceased, a solution of 4.7 g of the compound of Example 547A in 10 mL of THF was added. The mixture was stirred at reflux for 24 hr, then cooled with an ice bath and quenched by sequential dropwise addition of 1.7 mL H_2O , and 17 mL of 25% w/v aqueous NaOH. The white precipitate was filtered, and washed with about 50 mL of THF. The combined filtrate and washings were concentrated to 3.85 g of an oil. To an ice cooled solution of this material in 35 mL of ethyl acetate was added a solution of 5.0 g (29.2 mmol) of chloroacetic anhydride in 10 mL of ethyl acetate. The solution was stirred at 0°C for 30 min, then extracted with saturated aqueous NaHCO_3 solution (1 x 25 mL), 2M NaOH (1 x 25 mL), 5% NH_4OH (1 x 25 mL), 1M HCl (1 x 25 mL), and brine (1 x 25 mL), dried over MgSO_4 , filtered, and concentrated *in vacuo* to an oil. The product was purified *via* silica gel chromatography, eluting with 98:2 diethyl ether: methanol, to give 1.52 g (31%) of a colorless oil.

Example 547C

Ethyl (2*S*,3*R*,4*S*)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-hydroxybutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

To 1.52 g (6.85 mmol) of the compound of Example 547B was added 2.75 g (7.44 mmol) of the ethyl (2*S*,3*R*,4*S*)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate (prepared by neutralization of the

compound of Example 501G), 10 mL of DMSO, and 2 mL of N,N-diisopropylethylamine. The solution was stirred at ambient temperature for 22 h, then poured into 100 mL of water and extracted with diethyl ether (3 x 25 mL). The combined ether layers were washed with water (1 x 25 mL), 4% (v/v) H₃PO₄ (1 x 25 mL), saturated aqueous NaHCO₃ solution (1 x 25 mL), and brine (1 x 25 mL), dried over MgSO₄, filtered, and concentrated to an oil. This was purified via silica gel chromatography, eluting with 98:2 diethyl ether: methanol to give 3.0g (79%) of a colorless oil.

Example 547D

Ethyl (2S,3R,4S)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-bromobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

To an ice cooled solution of 2.80 g (5.05 mmol) of the compound of Example 547C in 27 mL of diethyl ether was added 1.4 mL (10 mmol) of triethylamine, then 0.58 mL of methanesulfonyl chloride. A white precipitate formed, and the suspension was stirred at 0 °C for 20 min. The reaction was diluted with 75 mL of diethyl ether, then extracted with saturated aqueous NaHCO₃ solution (2 x 25 mL), 5% NH₄OH (2 x 25 mL), and brine (1 x 25 mL), dried over MgSO₄, filtered, and concentrated to 3.0 g of a colorless oil. To this material in 45 mL of DMF was added 6.0 g (69 mmol) of LiBr. The reaction warmed to about 50 °C, then gradually cooled. The solution was stirred at ambient temperature for 4h, then poured into 450 mL of water, and extracted with diethyl ether (3 x 100 mL). The combined ether layers were back extracted with water (1 x 100 mL), and brine (1 x 100 mL), dried over MgSO₄, filtered, and concentrated *in vacuo* to an oil. The product was purified via silica gel chromatography, eluting with 3:1 diethyl ether: petroleum ether, to give 2.65 g (90%) of a colorless oil.

Example 547E

(2S,3R,4S)-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

To a solution of the compound of Example 547D (0.825 g, 1.34 mmol) in 3 mL of ethanol was added 5 mL of 4.07M dimethylamine in ethanol; the

resultant solution was heated at reflux for 75 min. Solvents were removed in vacuo. The residue was purified by flash chromatography on silica gel, eluting with 9:1 dichloromethane/methanol. The resultant material was taken up in 5 mL of 1.4N NaOH in 5:1 ethanol/water and stirred at ambient temperature for 14 hrs. Solvents were removed in vacuo; the residue was taken up in water, then adjusted to pH 6-7 with 1M HCl (~7 mL required). The mixture was extracted with EtOAc (3X); the aqueous layer was concentrated in vacuo. The residue was washed 3X with acetonitrile; the combined washes were filtered through Celite and concentrated to give 596 mg of a white foam.

Example 548

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 547, substituting the compound of Example 537B (ethyl (2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate) in Example 547C.

Example 549

(2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 547, substituting the compound of Example 536D (ethyl (2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate) in Example 547C.

Example 550

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 547, substituting the compound of Example 542C (ethyl (2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate) in Example 547C.

Example 551

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 547, substituting the compound of Example 544A (ethyl (2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate) in Example 547C.

Example 552

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-((N,N-di(nbutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 1, substituting the compound of Example 541C (ethyl(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate).

Example 553

(2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-((N,N-di(n-butyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

Prepared according to the procedures of Example 1, substituting the compound of Example 544B (ethyl (2S,3R,4S)-2-(2,2-Dimethylpent-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-pyrrolidine-3-carboxylate).

Example 554

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-((bis-(o-tolyl)methyl)amino)carbonylmethyl)-(pyrrolidine-3-carboxylic acid

Examples 554A through 554E

These compounds were prepared in Examples 501 B-F.

Example 554F

Bis-(o-tolyl)methylamine

In a 50 mL round-bottom flask were placed 2,2'-dimethylbenzophenone (prepared from commercially available methyl-2-methylbenzoate according to the procedure in *J. Chem. Soc.*, 1929, 1631) (2.50 g, 10 mmol), hydroxylamine hydrochloride (0.76 g, 11 mmol), pyridine (5 mL) and ethanol (5 mL). The mixture was stirred under reflux for 8 h, cooled to r.t., diluted with EtOAc (25 mL) and transferred into a separatory funnel. The aqueous layer was removed, and organic layer was washed in turns with CuSO₄ (25 mL), water (25 mL) and brine (25 mL). After concentration of the organic phase, the residual oil obtained was purified by column chromatography (elution with 10 % EtOAc in Hexanes) to give 1.31 g (73%) of oxime as a white crystalline solid.

To 55 mL of ammonia cooled in a dry ice-acetone bath was added 130 mg (6 mmol) of sodium metal. To the resulting blue solution at -78 °C was added slowly 650 mg (3 mmol) of the above oxime in 25 mL of anhydrous THF. The solution was stirred for 1 h followed by the addition of 1 g of ammonium chloride. The resulting colorless reaction mixture was warmed to r.t., transferred to a separatory funnel, diluted with 50 mL of water and extracted with dichloromethane (3x50 mL). The combined organic layers were dried (Na₂SO₄) and concentrated to give a yellowish oil. The residue was purified by a column chromatography (elution with 60% EtOAc in hexanes, followed by elution with 2% Et₃N in EtOAc) to give 500 mg (66%) of the pure amine.

Example 554G

N-(Bis-(o-tolyl)methyl) bromoacetamide

The compound of Example 554F (100 mg, 0.47 mmol) was dissolved in 2 mL of 1,2-dichloroethane. To this solution at -78 °C was added Et₃N (0.05 mL) and then dropwise bromoacetyl bromide (40 mL, 0.47 mmol in 1 mL of 1,2-dichloroethane). The reaction mixture was stirred at -78 °C for 10 min, then at r.t. for 2 h., diluted with water (10 mL), and extracted with 1,2-dichloroethane (2x25 mL). The combined organic layers were concentrated to give the bromoacetamide as a white solid (184 mg, 96%) suitable for further use without additional purification.

Example 554H

Ethyl trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-((bis-(o-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylate

The compound of Example 554G was dissolved in 5 mL of CH₃CN and added to a solution of 0.20 g (0.54 mmol) of the compound of Example 1E, *N,N*-diisopropylethylamine (0.1 mL) and CH₃CN (10 mL). The reaction mixture was stirred overnight at r.t., diluted with H₂O (25 mL) and extracted with EtOAc (2x25 mL). The combined organic fractions were

concentrated to give a yellow oil, which was purified by a column chromatography (elution with 40% of EtOAc in Hexanes) to give 250 mg (73%) of the title compound.

Example 554J

5 *trans,trans*-2-[4-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-((*o*-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The compound of Example 554H was dissolved in a solution of 50 mL of ethanol and 10 mL of aqueous sodium hydroxide (6N) and stirred overnight at room temperature. The solution was then diluted with 30 mL of water, transferred to a separatory funnel and extracted with a mixture of 20% Hexanes in EtOAc (2x50mL). The aqueous phase was treated with hydrochloric acid (3N) until pH=4 and extracted with chloroform (3x50mL). The combined organic fractions containing the acid product were concentrated to get a yellow viscous oil. The title compound was then isolated by lyophilization from dilute CH₃CN/TFA/H₂O as an amorphous solid: ¹H NMR (300 MHz, CDCl₃) δ 2.14 (s, 3H), 2.20 (s, 3H), 3.02-3.33 (m, 2H), 3.40-3.72 (m, 3H), 3.80 (s, 3H), 4.16-4.24 (broad s, 1H), 5.92 (m, 2H), 6.36-6.42 (m, 1H), 6.58-6.67 (m, 2H), 6.81 (t, *J* = 9 Hz, 4H), 6.88-7.00 (m, 2H), 7.05-7.27 (m, 8H). MS (ESI⁺) *m/e* 593 (M+H⁺). Anal. Calc for C₃₆H₃₆N₂O₆* 0.4 TFA: C, 69.25 H, 5.75 N, 4.39. Found: C, 69.20 H, 5.68 N, 4.22.

Example 555

20 *trans,trans*-2-[4-(2-Methoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-(2,2-dimethyl-1-phenylpropyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by using the procedures of Example 554, substituting 2,2-dimethyl-1-phenylpropan-1-one for 2,2'-dimethylbenzophenone in Example 554F and ethyl [4-(2-methoxyethoxy)benzoyl] acetate for ethyl (4-methoxybenzoyl) acetate in Example 1C. ¹H NMR (300 MHz, CDCl₃) δ 0.85 (s, 9H, minor diastereomer), 0.88 (s, 9H, major diastereomer), 3.44 (s, 2H), 3.10-3.42 (m, 3H), 3.46 (s, 3H), 3.51-3.75 (m, 4H), 4.06-4.13 (m, 2H), 4.72 (m, 1H), 5.97 (m, 2H), 6.77-7.45 (m, 12H). MS (ESI⁺) *m/e* 589 (M+H⁺). Anal. Calc for C₃₄H₄₀N₂O₇* 0.75 TFA: C, 63.24 H, 6.09 N, 4.15. Found: C, 63.33 H, 6.18 N, 4.05.

Example 556

35 *trans,trans*-2-[4-(2-Methoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-((*o*-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by using the procedures of Example 554, substituting ethyl [4-(2-methoxyethoxy)benzoyl] acetate for ethyl (4-methoxybenzoyl) acetate in Example 554C. ¹H NMR (300 MHz, CDCl₃) δ 2.13 (s, 3H), 2.20 (s, 3H), 2.94-3.23 (m,

3H), 3.32-3.51 (m, 2H), 3.47 (s, 3H), 3.58-3.69 (broad s, 1H), 3.76 (dd, $J = 6, 1.5$ Hz, 4H), 4.09 (t, $J = 4.5$ Hz, 1H), 5.93 (m, 2H), 6.34-6.41 (d, $J = 7.5$ Hz, 2H), 6.58 (broad s, 2H), 6.72-6.98 (m, 3H), 7.05-7.28 (m, 8H). MS (ESI+) m/e 637 ($M+H^+$). Anal. Calc for $C_{38}H_{40}N_2O_7^*$ 0.2 TFA: C, 69.93 H, 6.14 N, 4.25. Found: C, 70.03 H, 6.08 N, 4.21.

5

Example 557

trans,trans-2-[4-(2-Isopropoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-(2,2-dimethyl-1-phenylpropyl)-1-amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared by using the procedures of Example 554,

- 10 substituting 2,2-dimethyl-1-phenylpropan-1-one for 2,2'-dimethylbenzophenone in Example 554F and ethyl [4-(2-isopropoxyethoxy)benzoyl] acetate for ethyl (4-methoxybenzoyl) acetate in Example 1C. 1H NMR (300 MHz, $CDCl_3$) δ 0.83 (s, 9H, major diastereomer), 0.88 (s, 9H, minor diastereomer), 1.19 (d, $J = 7$ Hz, 6H), 3.14-3.83 (m, 9H), 4.07 (p, $J = 10.5, 4.5$ Hz, 2H), 4.27-4.47 (m, 1H), 4.70 (t, $J = 9$ Hz, 1H), 5.93-6.00 (m, 2H), 6.73-7.38 (m, 12H). MS (ESI+) m/e 617 ($M+H^+$). Anal. Calc for $C_{36}H_{44}N_2O_7^*$ 0.6 TFA: C, 65.21 H, 6.56 N, 4.09. Found: C, 65.15 H, 6.59 N, 4.01.

Example 558

trans,trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,3-dimethyl-1-phenylbutyl)-1-amino)carbonylmethyl]-pyrrolidine-3-carboxylic acid

The title compound was prepared according to the procedures of Example 554, substituting 3,3-dimethyl-1-phenylbutan-1-one (prepared from commercially available 3,3-dimethyl-butyl chloride according to the procedure in *J. Amer. Chem. Soc.*, **72**, 1950, 222-227) for 2,2'-dimethylbenzophenone in Example 554F. 1H NMR (300 MHz, $CDCl_3$) δ 0.84 (s, 9H, minor diastereomer), 0.89 (s, 9H, major diastereomer), 1.68 (t, $J = 7.5$ Hz, 2H), 3.05-3.30 (m, 2H), 3.34-3.53 (m, 2H), 3.62-2.74 (m, 1H), 3.77 (s, 2H), 3.80 (s, 3H), 4.92-5.02 (m, 1H), 5.97-6.01 (m, 2H), 6.77 (t, $J = 6$ Hz, 2H), 6.88 (q, $J = 18, 7.5$ Hz, 2H), 6.97 (d, $J = 6$ Hz, 1H), 7.10-7.40 (m, 7H). MS (ESI+) m/e 559 ($M+H^+$). Anal. Calc for $C_{33}H_{38}N_2O_6^*$ 0.45 TFA: C, 66.75 H, 6.35 N, 4.59. Found: C, 66.69 H, 6.32 N, 4.46.

30

Example 559

trans,trans-2-[4-(2-Isopropoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-((1-(o-tolyl)-1-(o-ethylphenyl)-methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by using the procedures of Example 554,

- 35 substituting ethyl [4-(2-isopropoxyethoxy)benzoyl] acetate for ethyl (4-methoxybenzoyl) acetate in Example 554C and 2-ethyl-2'-methylbenzophenone (prepared from commercially available methyl-2-methylbenzoate according to the procedure in *J. Chem. Soc.*, 1929, 1631)

for 2,2'-dimethylbenzophenone in Example 554F. ^1H NMR (300 MHz, CDCl_3) δ 1.04-1.28 (m, 9H) 2.20 (s, 3H), 2.53 (broad s, 2H), 2.94-3.23 (m, 3H), 3.35-3.87 (m, 6H), 4.08 (broad s, 3H), 4.52 (broad s, 1H), 5.96 (s, 2H), 6.48-7.35 (m, 15H). MS (ESI+) m/e 678 ($\text{M}+\text{H}^+$).

Anal. Calc for $\text{C}_{41}\text{H}_{46}\text{N}_2\text{O}_7$ * 0.95 TFA: C, 65.46 H, 6.01 N, 3.46. Found: C, 65.47 H, 6.00 N, 3.10.

Example 560

trans,trans-2-(4-(2-(2-Propoxy)ethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-N-phenyl-N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid

Example 560A

N-phenyl-t-butylamine

t-Butylamine(2.0 eq.) was stirred in dry THF at r.t. and n-butyllithium(1.2 eq.) was added slowly. The resulting mixture was stirred for 30min. and then bromobenzene(1.0 eq.) added, refluxed for 4 hr. After work-up, it was purified and separated by a column(silica) to elute with hexane and ethyl acetate(9:1). Yield 50%.

Example 560B

Phenyl-t-butyl nitrosoamine

Phenyl-t-butylamine (6g, 0.04mol) was treated with conc.HCl (5ml) and a solution of NaNO_2 (6.4g, 2.4eq.) in 20ml of water was added slowly. The resulting mixture was stirred for 2hr at r.t. to produce an oily layer which was extracted with EtOAc, washed with brine, and dried over Na_2SO_4 . Yield 6g (85%). This nitrosoamine was used for the next step without further purification.

Example 560C

N-phenyl-t-butylhydrazine

To a stirred suspension of zinc dust(5.14g, 0.079mol) in water(15ml) was added dropwise a solution of the compound of Example 560B (3.5g, 0.02mol) in acetic acid(9ml) and the resultant mixture was stirred for 1hr at r.t. Dichloromethane(20ml) was added, the mixture was adjusted to pH 8-9 with 15% NaOH, The zinc dust was removed by filtration, and the crude reaction mixture was extracted with dichloromethane. The combined organic layers were dried(MgSO_4). Yield 1.97g(60%).

Example 560D

N-phenyl,N-t-butyl-N'-(bromoacetyl) hydrazine

The title compound was prepared by the procedure described Example 554G, substituting the compound of Example 560C.

Example 560E

- 5 *trans,trans*-2-(4-(2-(2-Propoxyethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-phenyl-N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Example 554. ¹H NMR (300MHz, DMSO) δ 0.92(d, J=24Hz, 6H), 1.10(s, 9H), 2.59-2.80(m, 2H), 2.95-3.10(m, 1H), 3.25-3.51(m, 3H), 3.58-3.70(m, 3H), 3.73-3.88(m, 1H), 4.02-4.07(m, 2H), 5.97-6.0(m, 2H), 6.78-6.93(m, 4H), 7.02-7.26(m, 7H), 7.35(d, J=8Hz, 1H). MS(ESI+) m/e 618(M+H+). Anal. Calc for C₃₅H₄₃N₃O₇: C, 67.07 H, 7.08 N, 6.70. Found: C, 67.21 H, 6.61 N, 6.40.

Example 561

- 15 *trans,trans*-2-(4-(2-Methoxyethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-phenyl-N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid

The title compound was prepared by the procedures described in Examples 554 and 560. ¹H NMR (300MHz, CD₃OD) δ 1.22(s, 9H), 2.87(d, J=15Hz, 1H), 2.98-3.07(m, 2H), 3.27(bs, 1H), 3.42(s, 3H), 3.60-3.68(m, 2H), 3.72-3.76(m, 2H), 3.92(d, J=9Hz, 1H), 4.10-4.14(m, 2H), 5.95(dd, J=2Hz, 4Hz, 2H), 6.82(d, J=8Hz, 1H), 6.90(dd, J=2Hz, 9Hz, 1H), 7.96(d, J=8Hz, 2H), 7.07(d, J=2Hz, 1H), 7.10-7.23(m, 5H), 7.42(d, J=8Hz, 2H). MS (ESI+) m/e 590 (M+H⁺). Anal. Calc for C₃₃H₃₉N₃O₇: C, 64.70, H, 6.67, N, 6.47. Found: C, 64.40, H, 6.40, N, 6.70.

- 25 As an indication that the compounds described herein act through binding to endothelin receptors, the compounds have been evaluated for their ability to displace endothelin from its receptor.

- 30 As an indication that the compounds described herein act through binding to endothelin receptors, the compounds have been evaluated for their ability to displace endothelin from its receptor.

Binding Assay

ET_A Receptor

- 35 Preparation of membranes from MMQ cells:

MMQ (MacLeod/MacQueen/Login cell line (prolactin secreting rat pituitary cells)) cells from 150 mL culture flasks were collected by centrifugation (1000xg for 10 min) and then homogenized in 25 mL of 10 mM Hepes (pH 7.4) containing 0.25 M sucrose and protease inhibitors (3 mM EDTA, 0.1 mM PMSF, and 5 µg/mL Pepstatin A) by a micro ultrasonic cell disruptor (Kontes). The mixture was centrifuged at 1000xg for 10 min. The supernatant was collected and centrifuged at 60,000xg for 60 min. The precipitate was resuspended in 20 mM Tris, pH 7.4 containing the above protease inhibitors and centrifuged again. The final pellet was resuspended in 20 mM Tris, pH 7.4 containing protease inhibitors and stored at -80°C until used. Protein content was determined by the Bio-Rad dye-binding protein assay.

(¹²⁵I)ET-1 binding to membranes:

Binding assays were performed in 96-well microtiter plates pretreated with 0.1% BSA. Membranes prepared from cells were diluted ~100 fold in Buffer B (20 mM Tris, 100 mM NaCl, 10 mM MgCl₂, pH 7.4, with 0.2% BSA, 0.1 mM PMSF, 5 µg/mL Pepstatin A, 0.025% bacitracin, and 3 mM EDTA) to a final concentration of 0.2 mg/mL of protein. In competition studies, membranes (0.02 mg) were incubated with 0.1 nM of (¹²⁵I)ET-1 in Buffer B (final volume: 0.2 mL) in the presence of increasing concentrations of unlabeled ET-1 or a test compound for 4 hours at 25 °C. After incubation, unbound ligands were separated from bound ligands by a vacuum filtration method using glass-fiber filter strips in PHD cell harvesters (Cambridge Technology, Inc., MA), followed by washing the filter strips with saline (1 mL) for three times. Nonspecific binding was determined in the presence of 1 µM ET-1. The data are shown in Table 4. The per cent inhibition at a concentration of 1 µM is shown. The data show that the compounds of the invention bind to the endothelin receptor.

Table 4
Binding Data

Example	% Inhibition of ETA at 1 µM	Example	% Inhibition of ETA at 1 µM
1D	96.4	34	95.5
2	58.4	35	91.8
3	42.2	36	94.5
4	78.2	37	47.9

5	95.1	38	100.0
6B	34.9	39	83.6
7	63.4	40	94.8
8	53.7	41	89.9
9	69.2	42	95.2
10	66.1	43	99.2
14	86.6	44	91.3
15	84.8	45	85.4
16	96.0	46	90.4
17	73.9	47	95.1
18	97.3	48	96.3
19	90.3	52	84.0
20	80.9	54	64.6
21	56.3	55	50.5
22	86.3	56	34.3
23	85.9	57	93.2
26	83.0	58	81.9
27	61.2	59	70.8
28	63.8	60	42.8
29	85.3	61C	90.6
30	80.0	62	94.1
31B	93.6	63	92.0

Example	% Inhibition of ET _A at 1 μ M	Example	% Inhibition of ET _A at 1 μ M
64	95.0	98	86.8
65	82.8	99	92.1
66	87.7	100	76.8
67	96.3	101	89.2
68	84.6	102	75.2
69D	37.4	103	69.0
70	62.7	104	98.0
71	81.4	105	98.6
72C	80.7	106	90.0

73C	96.3	107	97.2
74	95.6	109	96.8
75C	95.3	110	94.4
76	93.1	111	101.8
79	100.4	112	94.9
80	89.4	113	94.3
82	90.3	114	86.2
83	85.0	115	88.4
84	65.3	116	79.3
86	52.6	117	95.2
87	62.4	118	93.2
88	84.3	119	86.6
89	84.6	120	99.5
91C	91.6	121	98.6
92C	107.4	122	95.3
93C	59.2	125	97.2
95D	82.1	126	91.7
96	86.1	127	91.4
97	89.0	128	95.4

Example	% Inhibition of ET _A at 1 μ M	Example	% Inhibition of ET _A at 1 μ M
123	89.7	156	92.6
124	91.0	157	83.8
129	100.1	158	91.8
130	91.0	159	36.2
131	89.5	160B	80.3
132	90.0	161	93.6
133	88.6	162B	91.5
134	92.2	163	90.6
135B	77.7	164	98.6
136	79.4	165	54.1
138	83.0	166	91.6
139	98.6	167	94.4
140	106.3	291	100.0
141	92.8	293	89.8
142B	78.7	294	77.7
143	20.6	295	93.0
144	78.2	296	87.1
145	32.4	297	84.4
146	25.0	298	93.3
147	73.0	299	90.4
148	94.7	300	96.1
149	84.6	301	96.7
150	93.6	302	86.6
151	80.5	303	87.2
152	86.9	304	89.7
153	97.1	305	87.4
154	80.2	306	93.3
155	92.7	307	92.2

Example	% Inhibition of ETA at 1 μ M	Example	% Inhibition of ETA at 1 μ M
308	93.0	351	99.0
309	80.7	352	96.2
310	87.1	353	73.7
311	92.3	354	79.3
312	88.2	355	100
313	96.3	356	93.5
314	86.0	357	96.3
315	82.7	358	62.7
316	74.0	359	94.7
317	68.5	360	93.7
318	79.0	361	92.8
319	79.0	362	94.1
320	82.2	363	82.3
322	95.6	365	59.2
323	91.3	366	91.5
324	95.0	367	71.0
334	88.0	368	94.6
335	84.1	370	84.3
340	94.0	371	97.2
341	87.4	372	91.6
342	89.9	373	92.9
343	98.7	374	91.4
344	95.6	375	97.8
345	86.6	376	90.2
346	88.9	377	85.6
348	91.3	378	91.1
349	73.0	379	90.7
350	92.1	380	99.0

Example	% Inhibition of ETA at 1 μ M	Example	% Inhibition of ETA at 1 μ M
381	95.7	408	100
382	96.8	409	89.4
383	91.4	410	91.4
384	79.4	411	93.5
385	86.2	412	86.4
386	47.8	413	99.5
387	98.7	414	91.4
388	69.2	415	87.3
389	100	416	86.4
390	98.2	417	98.7
391	45.6	418	100
392	93.7	420	100
393	100	421	100
394	97.8	422	96.6
395	79.8	423	89.1
396	98.7	424	85.8
397	100	425	90.8
398	90.0	426	97.2
399	59.9	427	100
400	93.0	428	100
401	96.5	429	100
402	80.5	430	94.1
403	96.1	431	99.1
404	95.4	432	95.5
405	86.4	433	99.6
406	94.5	434	100
407	100	435	97.8

Example % Inhibition of
ET_A at 1 μ M

436 100
437 100
438 94.3
439 94.3
440 100
441 98.3
442 100
443 100
444 100
445 98.1
446 97.8
447 96.9
448 97.4
449 100.0
450 99.7
451 100
452 100
453 94.4
454 96.8
455 99.1
456 95.3
457 88.9
458 93.4

Example % Inhibition of
ET_A at 1 μ M

459 97.4
460 91.6
461 99.6
462 98.3
463 96.1
464 97.1
465 95.1
466 94.2
467 93.6
468 88.7
469 98.7
470 100
471 100
475 91.6
476 82.3
477 80.1
479 96.5
495 95.9
496 92.7
497 83.7
498 81.6
499 68.5
500 55.7

Example.	% Inhibition of ETA at 1 μ M
502	95.7
503	97.0
504	97.1
505	95.8
506	99.7
507	99.3
508	97.6
509	100
510	100
511	99.2
512	98.9
513	98.0
514	100
515	99.1
516	99.7
517	94.1
518	96.3
519	99.1
520	97.4
521	100
523	99.0
524	99.2
525	100
526	100
527	96.6
528	98.3
529	98.1
531	99.8
532	100
533	97.9
536	100
537	97.2
554	58.2
555	66.7

556	24
557	72.2
558	79.8
559	5.8
560	0
561	0

As further demonstration of the efficacy of the described compounds as functional antagonists of endothelin, the ability of the described compounds to inhibit ET-1-induced phosphatidylinositol hydrolysis was measured.

Determination of Phosphatidylinositol (PI) Hydrolysis

MMQ cells (0.4×10^6 cells/mL) were labeled with $10 \mu\text{Ci/mL}$ of (^3H)myo-inositol in RPMI for 16 hours. The cells were washed with PBS, then incubated with Buffer A containing protease inhibitors and 10 mM LiCl for 60 minutes. The cells were then incubated with test compounds for 5 minutes, and then challenged with 1 nM ET-1 . ET-1 challenge was terminated by the addition of 1.5 mL of $1:2 \text{ (v/v)}$ chloroform-methanol. Total inositol phosphates were extracted after adding chloroform and water to give final proportions of $1:1:0.9 \text{ (v/v/v)}$ chloroform-methanol-water as described by Berridge (Biochem. J. 206 587-595 (1982)). The upper aqueous phase (1 mL) was retained and a small portion ($100 \mu\text{L}$) was counted. The rest of the aqueous sample was analyzed by batch chromatography using anion-exchange resin AG1-X8 (Bio-Rad). The IC_{50} is the concentration of test compound required to inhibit the ET-induced increase in PI turnover by 50%. The results of the above study clearly indicate that the compounds act as functional ET antagonists.

5

10

Table 5
Phosphatidylinositol Hydrolysis

Example	IC ₅₀ μ M
1D	0.025
14	0.017
15	0.010
16	0.009
18	0.009
19	0.024
30	0.001
31B	0.002
43	0.0001
46	0.002
47	0.0005
48	0.0004
291	0.0098
300	0.0012
534	0.05
553	0.0004

Table 6
ET_A/ET_B Selectivity

15

MMQ cells, porcine cerebellar tissues (known to contain ET_B receptors) and chinese hamster ovary cells (CHO) permanently transfected with the human ET_A or ET_B receptor were homogenized in 25 ml of 10 mM Hepes (pH

7.4) containing 0.25 M sucrose and a protease inhibitor (50 mM EDTA , 0.1 mM PMSF, 5 μ g/ml Pepstatin A, and 0.025% Bacitracin) using a micro ultrasonic cell disruptor. The mixture was centrifuged at 10000xg for 10 min. The supernatant was collected and centrifuged at 60,000xg for 60 min. The precipitate was resuspended in 20 mM Tris, pH 7.4 containing protease inhibitor and centrifuged again. The final membrane pellet was resuspended in 20 mM Tris, pH 7.4 containing protease inhibitors and stored at -80 °C until used. Protein content was determined by the Bio-Rad dye-binding protein assay.

Binding assays were performed in 96-well microtiter plates pretreated with 0.1% BSA. Membranes prepared from cells were diluted ~100 fold in Buffer B (20 mM Tris, 100 mM NaCl, 10 mM MgCl₂, pH 7.4, with 0.2% BSA, 0.1 mM PMSF, 5 μ g/mL Pepstatin A, 0.025% bacitracin, and 50 mM EDTA) to a final concentration of 0.2 mg/mL of protein. In competition binding studies, membranes (0.02 mg) were incubated with 0.1 nM of (125I)ET-1 (for ETA assay in MMQ or CHO cells transfected with human ET_A receptor) or (125I)ET-3 (for ET_B assay in porcine cerebellum or CHO cells transfected with human ET_B receptor) in Buffer B (final volume: 0.2 mL) in the presence of increasing concentrations of the test compound for 3 hours at 25 °C. After incubation, unbound ligands were separated from bound ligands by a vacuum filtration method using glass-fiber filter strips in PHD cell harvesters (Cambridge Technology, Inc., MA), washing the filter strips three times with saline (1 mL). Nonspecific binding was determined in the presence of 1 μ M ET-1. IC₅₀ values are calculated using an average of at least two separate determinations. The data shows the selectivity of the compounds of the invention in binding to the endothelin receptors.

Table 6

EXAMPLE NO.	rET-A (% @ 1 μ M)	rET-A IC ₅₀ (nM)	pET-B IC ₅₀ (nM)	Selectivity (rA/pB ratio)	hET-A IC ₅₀ (nM)	hET-B IC ₅₀ (nM)	Selectivity (hA/hB ratio)
502	95.7	3.0	71,000	23,000			
503	97.0	1.4	50,000	35,000	0.92	52,000	56,000
504	97.1	3.1	>100,000	>32,000	4.6	>100,000	>21,000

505	95.8	2.0	60,000	30,000	5.7	68,000	12,000
506	99.7	3.2	>100,000	>31,000	3.0	61,000	20,000
507	99.3	3.0	>100,000	>33,000	1.63	>100,000	>60,000
508	97.6	1.9	45,000	23,000	2.1	51,000	24,000
509	100	0.56	30,000	53,000	0.51	23,000	45,000
510	100	0.50	35,000	68,000	1.0	11,000	11,000
511	99.2	0.81	N.D.	---	0.60	15,000	25,000
512	98.9	0.42	>80,000	>190,000	0.58	60,000	>102,000
513	98.0	0.30	8,800	29,000	0.36	14,000	37,000
514	100	1.0	26,000	26,000	0.36	9,800	29,000
515	99.1	1.6	>62,000	>37,000	6.7	>100,000	>15,000
516	99.7	0.71	29,000	40,000	1.8	37,000	21,000
517	94.1	1.0	30,000	30,000	0.43	12,000	29,000
518	96.3	1.3	85,000	63,000	0.31	38,000	124,000
519	99.1	0.38	14,000	36,000	0.23	19,000	83,000
520	97.4	0.20	28,000	130,000			
521	100	0.67	37,000	54,000			
523	99.0	0.42	360	880	0.33	290	880

524	99.2	0.79	1,700	2,100	0.82	890	1,100
525	100	8.2	560	70			
526	100	42	---	---	17	7,400	440
527	96.6	7.9	10,000	1,300			
528	98.3	11	43,000	3,800			
529	98.1	3.6	6,300	1,700			
531	99.8	1.2	---	---	0.71	870	1,200
532	100	5.1	3,200	630			
533	97.9	76	7,900	100	40	22,000	560
534		0.12	0.36	3.0	0.08	0.28	3.5
536	100	0.52	17,000	33,000	0.92	52,000	56,000
537	97.2	0.96	5,900	6,200	0.23	1,900	8,200
552	97.3	0.78	7100,000	7125,000	1.0	>96,000	>96,000
553	100	0.26	42,400	160,000	0.29	39,500	136,000

Determination of Plasma Protein Binding

5 A stock solution of the test compound in 50% ethanol (2 mg/mL) was diluted 10X into PBS. A 0.4 mL sample of this secondary stock solution was added to 3.6 mL of fresh plasma, and incubated at room temperature for 1 hour. A 1 mL sample of this incubation mixture was transferred to a Centrifree ultrafiltration tube. The sample was centrifuged in a fixed-bucket rotor for approximately 2 min and the filtrate was discarded. The sample was

centrifuged for another 15-30 min. A 100 μ L sample of the ultrafiltrate was transferred to a micro HPLC sample vial containing 150 μ L of HPLC mobile phase and mixed thoroughly. A 50 μ L sample was injected and the concentration of drug in the ultrafiltrate was determined by HPLC analysis compared against a standard sample prepared identically in the absence of plasma. Ultrafiltrate concentrations are calculated from a calibration curve. Protein binding is calculated according to the equation:

$$\%PB = (1 - (Cu/Ci)) * 100\%$$

where Cu is the ultrafiltrate concentration and Ci is the initial plasma concentration. The percent of bound compound is listed in Table 7.

Table 7.

Example #43	> 99.5 % bound
Example #530	78% bound
Example #531	92% bound
Example #532	96.8% bound
Example #533	82.6% bound

It has been demonstrated in the literature (Wu-Wong, et al., Life Sci. 1996, 58, 1839-1847, and references contained therein) that compounds which are highly protein bound show decreased potency *in vitro* in the presence of plasma proteins. A decrease in *in vitro* potency may correspondingly result in reduced *in vivo* potency. An endothelin antagonist which has reduced protein binding might be expected to be less susceptible to this effect, and thus be more potent as an *in vivo* agent.

The ability of "reduced protein binding" endothelin antagonists to exhibit enhanced activity in the presence of serum albumin has been demonstrated through the following study: A series of binding curves is recorded for a given antagonist, each experiment performed in the presence of increasing concentrations of serum albumin.

Protocol for Albumin-induced binding shift studies: Binding assays were performed in 96-well microtiter plates precoated with 0.1% BSA unless otherwise indicated.

Membranes were diluted in Buffer B (20mM Tris, 100mM NaCl, 10mM MgCl₂, pH 7.4, 0.1 mM PMSF, 5mg/mL Pepstatin A, 0.025% bacitracin and 3 mM EDTA) to a final concentration of 0.05 mg/ml of protein. Varying concentrations of human serum

albumin (HSA) were added as indicated. In competition studies, membranes were incubated with 0.1 nM of [125 I]ET in Buffer B (final volume: 0.2 ml) in the presence of increasing concentrations of unlabeled test ligands for 4 hours at 25°C. After incubation, unbound ligands were separated from bound ligands by vacuum filtration using glass-fiber filter strips in PHD cell harvesters (Cambridge Technology, Inc., Watertown, MA), followed by washing the filter strips with saline (1 ml) for three times. Nonspecific binding was determined in the presence of 1 μ M ET-1.

Figure 1A.




Figure 1B.




Figure 1C.




Figure 1

Inhibition of [125 I]ET-1 binding to human ET_A receptor by ET_A antagonists. Each curve was determined in the presence of either 0%, 0.2%, 1%, or 5% HSA, and assays were performed as described above. The results are expressed as % of control binding, with [125 I]ET-1 binding in the absence of antagonist defining 100%. Each point represents the mean (\pm S.D.) of three determinations.

As observed in Figure 1A, a compound which is highly protein bound (Example 43, >98% bound) shows a rightward shift of the binding curve (toward decreasing potency) in the presence of increasing albumin levels. The compound of Example 531 (Figure 1B), in which protein binding is reduced to 92%, shows a substantial diminution of this rightward shift; the shift is completely eliminated with the compound of Example 530 (Figure 1C), in which protein binding is reduced to 78%. This experiment demonstrates that a reduction in protein binding translates into increased potency in the presence of plasma proteins, and suggests that such compounds may exhibit enhanced *in vivo* activity.

The observed reduction in protein binding, in compounds which retain high affinity for endothelin receptors, appears linked to the placement of "basic" functionality (groups which carry a positive charge at physiological pH).


Such compounds also exhibit improved solubility in aqueous solutions, as demonstrated below (Table 1) in an experiment in which maximum solubility was measured in aqueous media at varying pH at about 25°C. These results indicate that compounds that contain charged groups on the amide sidechain exhibit increased solubility over a significant range of pH. Such increased aqueous solubility, coupled with the enhanced potency resulting from decreased protein binding, might make such compounds preferred for development as parenteral agents. Table 8 presents the pH-Solubility profiles for representative compounds of the present invention.

Table 8.

pH	[Example 43] (mg/mL	[Example 531] (mg/ml
5.1	0.08	>3.3
6.5	0.51	>3.4
7.1	0.99	3.54
7.6	1.14	3.55

The present invention provides less protein bound compounds having improved *in vitro* and *in vivo* activity as pharmaceutical agents. The present invention also provides compounds that show that the affinity of hydrophobic acids for plasma protein may be reduced by attaching a counterbalanced charge at a biologically acceptable site. For example, protein binding is reduced by attaching a "basic" functionality (groups which carry a positive charge at physiological pH) on the amide sidechain (see Formula XII wherein R₃ has an amide sidechain).

A particularly preferred compound of formula I is a compound of formula IIIa, also known as ABT-627:



IIIa

Other suitable endothelin ET-A receptor antagonist may be used, such as those disclosed in U.S. Patent Nos. 6,048,893, 6,017,951, and 5,998,468.

The term "inhibit" is defined to include its generally accepted meaning which includes preventing, prohibiting, restraining, and slowing, stopping or reversing progression, or severity, and holding in check and/or treating existing characteristics. The present method includes both medical therapeutic and/or prophylactic treatment, as appropriate.

The methods of the present invention are useful in men as well as in women. Preferably, however, the methods of the present invention are useful in men, more preferably men with prostate cancer.

The ability of the compounds of the invention to treat cancers can be demonstrated according to the method described in J. Clin. Invest. 87 1867 (1991). Types of cancer includes primary cancer such as breast, prostate, lung, kidney, thyroid, myeloma, lymphoma, sarcoma, osteosarcoma, and ovarian.

The ability of the compounds of the invention to treat nociception can be demonstrated according to the method described in J. Pharmacol. Exp. Therap. 271 156 (1994).

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as loweralkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like

dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

Basic addition salts can be prepared *in situ* during the final isolation and purification of the compounds of formula I, or separately by reacting the carboxylic acid function with a suitable base such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia, or an organic primary, secondary or tertiary amine. Such pharmaceutically acceptable salts include, but are not limited to, cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, aluminum salts and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. Other representative organic amines useful for the formation of base addition salts include diethylamine,

ethylenediamine, ethanolamine, diethanolamine, piperazine and the like.

The compounds of the invention are useful for antagonizing endothelin in humans or other mammals. Total daily dose administered to a host in single or divided doses may be in amounts, for example, from 0.001 to 1000 mg/kg body weight daily and more usually 0.1 to 100 mg/kg for oral administration or 0.01 to 10 mg/kg for parenteral administration. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose.

Pharmaceutical formulations may be prepared by procedures known in the art. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination, and the severity of the particular disease undergoing therapy.

The compounds of the present invention may be administered orally, buccally, parenterally, sublingually, by inhalation spray, rectally, or topically in dosage unit formulations

containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, transcutaneous, intradermal, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleagenous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-propanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which

are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

The compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and

metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and phosphatidyl cholines (lecithins), both natural and synthetic.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

A representative solid dosage form, for example, a tablet or a capsule, comprises:

Compound of the invention:	35% w/w
Starch, Pregelatinized, NF	50% w/w
Microcrystalline Cellulose, NF	10% w/w
Talc, Powder, USP	5% w/w

While the compounds of the invention can be administered as the sole active therapeutic agent, they can also be used in combination with one or more co-therapeutic agents, such as anticancer drugs or methods including, but not limited to, hormonal agents, such as leuprolide (Lupron®); gonadorelin antagonists, such as goserelin (Zoladex®) and abarelix; bicalutamide; nilutamide; flutamide; vitamin D; vitamin D analogues; estrogen and estrogen analogues, such as diethylstilbestrol; prednisone; hydrocortisone; ketoconazole; cyproterone acetate; progesterone; 5-alpha reductase

inhibitors, such as finasteride; bone-seeking radionuclides, such as samarium (Quadramet®), strontium (Metastron®), and ¹⁸⁶rethium; external beam radiation, including three

dimensional conformal radiation; brachytherapy, which is the

- 5 implantation of radioactive seeds directly into the prostate; monoclonal antibodies such as trastuzumab (Herceptin®); anti-angiogenic agents such as thrombospondin peptide or kringle 5; matrix metalloproteinase inhibitors; farnesyl transferase

inhibitors; lycopenes; urokinase; plasminogen activator

- 10 inhibitors; plasminogen activator receptor blockers; apoptosis inducers; selective and non-selective alpha blockers; platinum agents, such as cis-platinum and carbo-platinum; taxane class agents, such as docitaxil and paclitaxil; estramustine; gemcytabine; adriamycin; doxorubicin; daunorubicin;

- 15 mitoxantrone; vinblastine; vincristine; capecitabine; irinotecan; topotecan;

5-fluorouracil; interferons; cytoxan; methotrexate;

cytokines, such as IL-2; PPAR agonists, such as thiazolidine diones; retinoid-type agents, 5-lipoxygenase inhibitors, such

- 20 as zyfo (Zilueton®), COX-2 inhibitors; gene-therapy based therapeutics, including sense and anti-sense genes; cholesterol lowering drugs, such as lovastatin, pravastatin, and simvastatin; bisphosphonates; osteoprotegrin; and antibodies, both monoclonal and polyclonal; antibody-coupled

radionucleotides; antibody-coupled cytotoxic agents; antibody-coupled radionucleotides; viral-vector delivered agents; vaccines directed at protein, carbohydrate, or nucleic acid targets; aminoglutethimide; and suramin.

5 These combinations can be administered as separate compositions or as a single dosage form containing both or all agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions, which are given at the same time or different times, or the therapeutic
10 agents can be given as a single composition.

In addition, the compounds invention can be used in combination with one or more co-therapeutic agents which impede net bone loss, such as estrogens, bisphosphonates, and estrogen receptor modulators, such as raloxifene, and calcitonin.

15 The compounds of the invention can additionally be administered in combination with surgery, such as radical prostatectomy, cryotherapy, transurethral resection of the prostate as an adjuvant, and the like, or prior to surgery as a neoadjuvant agent.

20 The current major diseases or conditions of bone which are of public concern include, but are not limited to, post-menopausal osteoporosis, ovariectomy patients, senile osteoporosis, patients undergoing long-term treatment of corticosteroids, side effects from glucocorticoid or steroid
25 treatment, patients suffering from Cushings's syndrome, gonadal

dysgenesis, periarticular erosions in rheumatoid arthritis, osteoarthritis, Paget's disease, osteomalacia, hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, 5 osteoporosis from Lupron therapy, and starvation. All of these conditions are characterized by bone loss, resulting from an imbalance between the degradation of bone (bone resorption) and the formation of new healthy bone. This turnover of bone continues normally throughout life and is the mechanism by 10 which bone regenerates. However, the conditions stated above will tip the balance towards bone loss such that the amount of bone resorbed is inadequately replaced with new bone, resulting in net bone loss.

Examples

15 Studies were performed on male subjects with asymptomatic hormone refractory prostate cancer with rising PSA levels and on male subjects with symptomatic hormone refractory prostate cancer with rising PSA levels and pain. Subjects in the phase 20 II studies had castrate levels of testosterone, either due to pharmacologic intervention, via leuprolide (Lupron®) or goserelin (Zoladex®), or via surgical castration. Subjects received ABT-627 or placebo. The following tests were conducted:

ABT-627 was formulated in 2.5 and 10 mg doses. An oral liquid formulation of ABT-627 was also prepared as follows: 1 mg/ml ABT-627, 50% glycerin, 14% alcohol, and water. Matching placebos were also provided.

5 A number of recognized or putative biochemical markers of disease progression have been used to monitor treatment of individuals with prostate cancer. Among these markers are serum Prostate Specific Antigen (PSA), serum acid Phosphatase, Interleukin-6, and Chromagranin-A. As currently accepted,
10 favorable treatment is marked by a decrease or slower rate of increase for PSA, acid phosphatase, and Interleukin-6, while a favorable response is marked by an increase in Chromagranin-A.

Serum samples were obtained from subjects during treatment with the ET antagonist ABT-627 in order to determine PSA, acid
15 phosphatase, IL-6, and Chromagranin-A values.

Prostate Specific Antigen Level Assay

The effect of ABT-627 administration on prostate specific antigen (PSA) levels in human subject serum samples was
20 determined using the procedure described in the Chiron Diagnostics ACS: Centaur PSA2 Assay. This assay is a two-site sandwich immunoassay which uses direct chemiluminescence and constant amounts of two antibodies. The first antibody, the Lite Reagent, is an affinity purified polyclonal sheep anti-PSA
25 antibody labeled with acridinium ester. The Lite Reagent is

purchased as a 5.0 mL reagent pack comprising the polyclonal sheep anti-PSA antibody (3.1 μ g) in buffered saline with sodium azide (0.1%). The second antibody, the Solid Phase, is a monoclonal mouse anti-PSA antibody covalently coupled to

5 paramagnetic particles. The Solid Phase is purchased as a 25.0 mL reagent pack comprising the covalently coupled monoclonal mouse anti-PSA antibody (316 μ g) in buffered saline with sodium azide (0.1%). The assay was performed at Quintiles Laboratories (Smyrna, GA) using Chiron Diagnostics ACS:

10 Centaur® Automated Chemiluminescence Systems.

Briefly, a subject population was treated with a placebo or 2.5 mg or 10 mg of ABT-627. Blood samples were collected, allowed to adequately clot, centrifuged at 1000 x g for 15-20 minutes, and stored at -20 °C if not assayed within 48 hours.

- 15 A cuvette was charged sequentially with serum, Lite Reagent (50 μ L), and Solid Phase (250 μ L). The resulting mixture was incubated for 7.5 minutes at 37 °C, separated, and treated with the solution of Acid Reagent and Base Reagent to initiate the chemiluminescent reaction. A direct relationship exists
- 20 between the amount of PSA present in the patient sample and the RLU's (relative light units) detected. As shown by the area under the curve (AUC) in Figure 2, the rate of increase of PSA in the serum samples decreases after the administration of ABT-

627, demonstrating the effectiveness of ABT-627 as an agent for treating prostate cancer.

Acid Phosphatase Levels

5 The effect of ABT-627 administration on Acid Phosphatase levels in human subject serum samples was determined at Quintiles Laboratories using the chemical test described in Sigma Diagnostics Acid Phosphatase (ACP) Procedure No. 435. The enzyme Acid Phosphatase (ACP) catalyzes the hydrolysis of
10 alpha-naphthyl phosphate to alpha-naphthol and inorganic phosphate. The alpha-naphthol immediately reacts with fast red TR salt to produce a yellow chromophore with an absorbance maximum at 405 nm. The rate of increase in absorbance at 405 nm is directly proportional to ACP activity in the sample. ACP
15 activity was determined in the presence and absence of L-tartrate, the difference being attributed to prostatic acid phosphatase activity.

Briefly, a subject population was treated with a placebo or 2.5 mg or 10 mg of ABT-627. Blood samples were collected,
20 allowed to adequately clot, centrifuged at 1000 x g for 15-20 minutes, and stored at -20 °C if not assayed within 48 hours. Assays were performed on a Hitachi Spectrophotometer. A cuvette was charged sequentially with ACP reagent (1 mL), prepared as described in the assay protocol, and serum (0.1 mL). The
25 mixture was agitated and incubated for 5 minutes, and an

absorbance (A) at 405 nm (versus water as a reference) was read to provide an initial absorbance. The mixture was incubated for another 5 minutes, and a second absorbance was read to provide a final absorbance. A change A/5 minute value was obtained by subtracting the initial absorbance from the final absorbance and was used to calculate total ACP activity.

To provide the tartrate-resistant acid phosphatase activity, the above procedure was repeated with the addition of ACP tartrate reagent (0.01 mL) to the cuvette containing the ACP reagent and mixing before adding the serum. Prostatic acid phosphatase activity was calculated by subtracting the the tartrate-resistant acid phosphatase activity from the ACP activity. As shown shown by the (AUC) in Figure 7, the rate of increase and the average change from baseline for acid phosphatase was decreased in those subjects treated with ABT-627, again demonstrating the effectiveness of ABT-627 as an agent for treating prostate cancer.

Chromagranin-A Levels

The effect of ABT-627 administration on Chromagranin-A levels in human serum samples was determined by proprietary assay conducted at the Nichols Institute. The procedure is a two site chemiluminescence assay (ICMA) using one monoclonal antibody conjugated with biotin, another monoclonal antibody

labeled with an acridinium ester, and an avidin-coated solid phase. The antibody/Chromagranin-A/antibody complex is bound to the solid phase by the avidin-biotin interaction and unbound materials are removed by washing. The bound, acridinium-labeled material produces light that is detected in a luminometer after addition of triggering agents. The Limit of Detection (LOD) for the assay was 0.07 ng/mL. As shown by the AUC in Figure 8, the average change from baseline for Chromagranin-A was higher for subjects treated with 2.5 mg/day of ABT-627, again demonstrating the effectiveness of ABT-627 as an agent for treating prostate cancer.

Interleukin-6 Levels

The effect of ABT-627 administration on Interleukin-6 levels in human serum samples was determined at Quintiles Laboratories using a sandwich immunoassay. Human serum samples and standards were incubated in microtiter plate wells coated with a monoclonal anti-IL-6 antibody, in the presence of a second monoclonal anti IL-6 antibody, linked to acetylcholinesterase. After incubation, the wells were washed, and the bound enzymatic activity was measured using a chromogenic substrate. The intensity of the color was proportional to the concentration of IL-6 in the sample or standard. As shown by the AUC Figure 1, the average change in baseline for Interleukin-6 was lower in those subjects treated

with ABT-627, demonstrating the effectiveness of ABT-627 as an agent for reducing inflammation and ameliorating pain.

Bone Scan Methodology

5 Bone scans were performed with an NDA approved, Tc-99m phosphonate type radiopharmaceutical. This technique uses whole body format (skull to feet) so that anterior and posterior images are presented when using a 510 K-approved gamma camera. Alternatively, spot views covering both anterior
10 and posterior projections of the total body can be obtained. Interpretation was performed according to standard nuclear medicine criteria, on a bone by bone basis, by recording the number of lesions at each site. Each site was evaluated against a confidence score of 1 to 5, where 1 is negative, 2 is
15 probably negative, 3 is equivocal, 4 is probably positive, and 5 is definitely positive. The MSKCC (Clin. Can. Res. 1998; 4:1765-1772) was used to record these findings. For the purposes of scoring the extent of disease or the response to treatment, lesions with a confidence score of 4 and 5 were
20 considered positive, and all other lesions were considered negative. In addition, in a blinded study, a reference nuclear medicine physician interpreted the bone scans quantitatively as follows: the percent of involved bone was estimated for each individual bone, and the individual bone involvement was summed
25 to calculate a global percent bone scan index (BSI). More

specifically, the bone scan was separated into three indices. The first was the appendicular scan which involved arms and legs (i.e. the humerus and all bones distal to the humerus and the femur and everything distal to the femur). The second
 5 was the axial (everything but the arms and the legs). The results of these scans were combined to provide the total BSI.

Bone scans were conducted on each subject on day one of the study, and on the final day of the study, and the changes from baseline in bone scan index scores were analysed
 10 by mean change and mean percent change, adjusting for baseline characteristics as co-variables using SAS version XXX software.

As shown in Figure 6, bone scans indicated a decrease in the proportion of total skeletal involvement in those subjects receiving ABT-627 versus placebo, demonstrating the
 15 effectiveness of ABT-627 as an agent for reducing the fraction of total skeletal involvement by tumor.

VAS Methodology/Administration/Analysis

The Visual Analog Scale (VAS) is a common instrument of
 20 pain assessment performed by having a subject draw a vertical line on a 10 cm scale at the point that best describes his or her pain on average in the last 24 hours. A diagram of the scale is shown below:

(not to scale)

During the course of the study, pain assessments were done daily, at bedtime, by the subject. If the subject was unable to maintain the log, a caregiver could complete the log on his or her behalf. The log also contained a table on which was recorded all daily pain medication consumed by the patient. The logs of daily VAS scores and analgesic consumption were collected at biweekly visits of the subject to the clinic when a new log was distributed. Clinical personnel who received the logs measured the score by measuring the distance (in mm) from the "no pain" end mark to the point where the subject's line crossed the VAS line. The number was written into the case report form next to the date the subject completed that page of the logbook.

Subjects with pain were initially stabilized in their pain so that their pain was treated to a tolerable and constant level. For this study, "tolerable and constant" refers to a pain score less than or equal to 5 cm on the VAS for an average of seven successive days while using four or less rescue doses of pain medication per day. A rescue medication dose refers to a dose equal to one single dose a patient used for common timed pain relief.

The weekly VAS scores were calculated excluding the lowest and highest score for each week and averaging the remaining

five scores. If there were two days with the same VAS score, the day with the highest analgesic use was discarded.

The weekly mean VAS score was used to define subjects as responders or non-responders. A subject was considered a

5 responder based on the reduction in the pain intensity: a weekly VAS score reduction of greater than or equal to 25% during at least two consecutive weeks without an increase of analgesic use during the same period (compared to baseline). Alternatively, a subject was considered a responder if his pain
10 analgesic consumption was reduced by at least 25% during at least two consecutive weeks without a concomitant increase in VAS score.

The percentage of responders in each treatment group was compared to evaluate drug efficacy. The comparison was
15 subjected to an adjustment for baseline characteristics and prognostic factors as co-variates, and the analysis was performed using the Cochran-Mantel-Haenszel test or a generalized linear model.

Weekly VAS scores are examined using a longitudinal
20 analysis method to explore trends over time. The duration of the response, defined as the time from baseline to the last weekly assessment for which the responder definition was satisfied, was analyzed using the Kaplan-Meier methodology and logrank test. Cox proportional hazard models were used as
25 needed (see U.S. Department of Health and Human Services.

Management of Cancer Pain Clinical Practice Guidelines. AHCPR Publication #94-0592, Rockville, MD (1994). As shown by the AUC in Figure 3, VAS scores showed a decrease in pain, independent of the effects of morphine, after treatment with 5 with ABT-627, demonstrating the effectiveness of ABT-627 as an agent for ameliorating pain.

Osteoblastic Activity and Bone Markers

Markers of osteoblastic activity were assessed using urine 10 samples. Bone markers include bone alkaline phosphatase (BAP), deoxypridinoline, and N-telopeptide of Type I collagen. Blood samples were collected prior to dosing on Day 1, Day 42, Day 84, Day 168, and every 28 days after Day 168, with a final collection on the last day of the study.

Bone Alkaline Phosphatase


Bone Alkaline Phosphatase levels were determined using the bone-specific Alkphase-B® assay published by Metra Biosystems (Mountain View, CA). As shown by the AUC in Figure 5, BAP 20 levels decreased in subjects treated with ABT-627, demonstrating the effectiveness of ABT-627 as an agent for inhibiting abnormal bone remodeling.

Crosslinked N-Telopeptide Levels:

Cross-linked N-telopeptide levels were determined using the DiaSorin (Stillwater, MN) assay for the quantitative determination of carboxyterminal cross-linked telopeptide of type I collagen (ICTP) in human serum by equilibrium

radioimmunoassay (RIA). Briefly, samples were incubated with the ^{125}I ICTP tracer and ICTP primary antibody for 2 hours at 37 °C. Following the 2 hour incubation, a pre-precipitated second antibody complex was added to separate the bound from free tracer. The assay was then centrifuged and decanted after a 30 minute incubation at room temperature. The bound tracer in the pellet was counted with a gamma counter. Counts were inversely proportional to the amount of ICTP present in each sample. As shown by the AUC in Figure 4, Crosslinked N-telopeptide levels decreased in subjects treated with ABT-627, demonstrating the effectiveness of ABT-627 as an agent for inhibiting the bone remodeling associated with bone diseases.

The present invention covers compounds having the formula XII:



XII

wherein

Z is $-C(R_{18})(R_{19})-$ or $-C(O)-$ wherein R_{18} and R_{19} are independently selected from hydrogen and loweralkyl;

n is 0 or 1;

R is $-(CH_2)_m-W$ wherein m is an integer from 0 to 6 and W is

- (a) $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,
- (b) $-PO_3H_2$,
- (c) $-P(O)(OH)E$ wherein E is hydrogen, loweralkyl or arylalkyl,
- (d) $-CN$,
- (e) $-C(O)NHR_{17}$ wherein R_{17} is loweralkyl,
- (f) alkylaminocarbonyl,
- (g) dialkylaminocarbonyl,
- (h) tetrazolyl,
- (i) hydroxy,
- (j) alkoxy,
- (k) sulfonamido,
- (l) $-C(O)NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl, aryl or dialkylamino,
- (m) $-S(O)_2NHC(O)R_{16}$ wherein R_{16} is defined as above,



(n) ,




(o) ,



(p) ,



(q) ,




- 5 R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl,
- 10 hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen;
- 15 R_3 is (a) $R_4-C(O)-R_5$, $R_4-C(O)-R_5-N(R_6)-$, wherein R_5 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene, (iv) $-N(R_{20})-R_8$ or $-R_{8a}-N(R_{20})-R_8$ wherein R_8 and R_{8a} are independently selected from the group consisting of alkylene and alkenylene and R_{20} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl or cycloalkylalkyl or (v) $-O-R_9$ or
- 20 $-R_{9a}-O-R_9$ wherein R_9 and R_{9a} are independently selected from alkylene; R_4 and R_6 are $(R_{11})(R_{12})N$ - wherein R_{11} and R_{12} are independently selected from
- (1) hydrogen,
 - (2) loweralkyl,
 - (3) haloalkyl,
 - (4) alkoxyalkyl,
 - (5) haloalkoxyalkyl,
 - (6) alkenyl,
 - (7) alkynyl,
















- 5 (8) cycloalkyl,
 (9) cycloalkylalkyl,
 (10) aryl,
 (11) heterocyclic,
 (12) arylalkyl,
 (13) (heterocyclic)alkyl,
 (14) hydroxyalkyl,
 (15) alkoxy,
 (16) aminoalkyl,
 10 (17) trialkylaminoalkyl,
 (18) alkylaminoalkyl,
 (19) dialkylaminoalkyl,
 (20) carboxyalkyl,
 (21) (cycloalkyl)aminoalkyl,
 15 (22) (cycloalkyl)alkylaminoalkyl,
 (23) (heterocyclic)aminoalkyl, and
 (24) (heterocyclic)aminoalkyl, with the proviso that at least one of
 R₁₁ and R₁₂ is selected from heterocyclic, aminoalkyl,
 alkylaminoalkyl, dialkylaminoalkyl, trialkylaminoalkyl,
 20 alkylaminoalkyl, dialkylaminoalkyl, carboxyalkyl,
 (cycloalkyl)aminoalkyl, (cycloalkyl)alkylaminoalkyl,
 (heterocyclic)aminoalkyl, and (heterocyclic)alkylaminoalkyl;
 or a pharmaceutically acceptable salt thereof.

25

Preferred compounds having reduced protein binding are shown in Table 9A wherein R may be selected from the substituents shown in Table 9B.

Table 9A.







































Table 9B.





The term “(heterocyclic)aminoalkyl” as used herein refers to a heterocyclic moiety attached to the parent compound through an aminoalkyl. Examples of (heterocyclic)aminoalkyl include (pyridine)aminopropyl, (benzofuran)aminopropyl, (tetrahydropyran)aminoethyl, and the like.

The term “(heterocyclic)alkylaminoalkyl” as used herein refers to a heterocyclic moiety attached to the parent compound through an alkylaminoalkyl. Examples of (heterocyclic)alkylaminoalkyl include (pyridine)ethylaminopropyl, (benzofuran)methylaminoisobutyl, (tetrahydropyran)methylaminoethyl, and the like.

The ability of the compounds of the invention to lower blood pressure can be demonstrated according to the methods described in Matsumura, et al., Eur. J. Pharmacol. 185 103 (1990) and Takata, et al., Clin. Exp. Pharmacol. Physiol. 10 131 (1983).

5 The ability of the compounds of the invention to treat congestive heart failure can be demonstrated according to the method described in Margulies, et al., Circulation 82 2226 (1990).

The ability of the compounds of the invention to treat myocardial ischemia can be demonstrated according to the method described in
10 Watanabe, et al., Nature 344 114 (1990).

The ability of the compounds of the invention to treat coronary angina can be demonstrated according to the method described in Heistad, et al., Circ. Res. 54 711 (1984).

The ability of the compounds of the invention to treat cerebral
15 vasospasm can be demonstrated according to the methods described in Nakagomi, et al., J. Neurosurg. 66 915 (1987) or Matsumura, et al., Life Sci. 49 841-848 (1991).

The ability of the compounds of the invention to treat cerebral ischemia can be demonstrated according to the method described in Hara et al.,
20 European. J. Pharmacol. 197: 75-82, (1991).

The ability of the compounds of the invention to treat acute renal failure can be demonstrated according to the method described in Kon, et al., J. Clin. Invest. 83 1762 (1989).

The ability of the compounds of the invention to treat chronic renal
25 failure can be demonstrated according to the method described in Benigni, et al., Kidney Int. 44 440-444 (1993).

The ability of the compounds of the invention to treat gastric ulceration can be demonstrated according to the method described in Wallace, et al., Am. J. Physiol. 256 G661 (1989).

30 The ability of the compounds of the invention to treat cyclosporin-induced nephrotoxicity can be demonstrated according to the method described in Kon, et al., Kidney Int. 37 1487 (1990).

The ability of the compounds of the invention to treat endotoxin-induced toxicity (shock) can be demonstrated according to the method
35 described in Takahashi, et al., Clinical Sci. 79 619 (1990).

The ability of the compounds of the invention to treat asthma can be demonstrated according to the method described in Potvin and Varma, *Can. J. Physiol. and Pharmacol.* 67 1213 (1989).

The ability of the compounds of the invention to treat transplant-induced atherosclerosis can be demonstrated according to the method described in Foegh, et al., *Atherosclerosis* 78 229-236 (1989).

The ability of the compounds of the invention to treat atherosclerosis can be demonstrated according to the methods described in Bobik, et al., *Am. J. Physiol.* 258 C408 (1990) and Chobanian, et al., *Hypertension* 15 327 (1990).

The ability of the compounds of the invention to treat LPL-related lipoprotein disorders can be demonstrated according to the method described in Ishida, et al., *Biochem. Pharmacol.* 44 1431-1436 (1992).

The ability of the compounds of the invention to treat proliferative diseases can be demonstrated according to the methods described in Bunchman ET and CA Brookshire, *Transplantation Proceed.* 23 967-968 (1991); Yamagishi, et al., *Biochem. Biophys. Res. Comm.* 191 840-846 (1993); and Shichiri, et al., *J. Clin. Invest.* 87 1867-1871 (1991). Proliferative diseases include smooth muscle proliferation, systemic sclerosis, cirrhosis of the liver, adult respiratory distress syndrome, idiopathic cardiomyopathy, lupus erythematosus, diabetic retinopathy or other retinopathies, psoriasis, scleroderma, prostatic hyperplasia, cardiac hyperplasia, restenosis following arterial injury or other pathologic stenosis of blood vessels.

The ability of the compounds of the invention to treat acute or chronic pulmonary hypertension can be demonstrated according to the method described in Bonvallet et al., *Am. J. Physiol.* 266 H1327 (1994). Pulmonary hypertension can be associated with congestive heart failure, mitral valve stenosis, emphysema, lung fibrosis, chronic obstructive pulmonary disease (COPD), acute respiratory distress syndrome (ARDS), altitude sickness, chemical exposure, or may be idiopathic.

The ability of the compounds of the invention to treat platelet aggregation, and thrombosis, can be demonstrated according to the method described in McMurdo et al. *Eu. J. Pharmacol.* 259 51 (1994).

The ability of the compounds of the invention to treat cancers can be demonstrated according to the method described in Shichiri, et al., *J. Clin. Invest.* 87 1867 (1991).

The ability of the compounds of the invention to treat IL-2 (and other cytokine) mediated cardiotoxicity and vascular permeability disorders can be demonstrated according to the method described in Klemm et al., Proc. Nat. Acad. Sci. 92 2691 (1995).

- 5 The ability of the compounds of the invention to treat nociception can be demonstrated according to the method described in Yamamoto et al., J. Pharmacol. Exp. Therap. 271 156 (1994).

- The ability of the compounds of the invention to treat colitis can be demonstrated according to the method described in Hogaboam et al (EUR. J. Pharmacol. 1996, 309, 261-269).

The ability of the compounds of the invention to treat ischemia-reperfusion injury in kidney transplantation can be demonstrated according to the method described in Aktan et al (Transplant Int 1996, 9, 201-207).

- 15 The ability of the compounds of the invention to treat angina, pulmonary hypertension, raynaud's disease, and migraine can be demonstrated according to the method described in Ferro and Webb (Drugs 1996, 51, 12-27).

- The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as loweralkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

- 35 Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric

acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

Basic addition salts can be prepared *in situ* during the final isolation and purification of the compounds of formula (I), or separately by reacting the carboxylic acid function with a suitable base such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia, or an organic primary, secondary or tertiary amine. Such pharmaceutically acceptable salts include, but are not limited to, cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, aluminum salts and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. Other representative organic amines useful for the formation of base addition salts include diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like.

The compounds of the invention are useful for antagonizing endothelin in a human or other mammal. In addition, the compounds of the present invention are useful (in a human or other mammal) for the treatment of hypertension, acute or chronic pulmonary hypertension, Raynaud's disease, congestive heart failure, myocardial ischemia, reperfusion injury, coronary angina, cerebral ischemia, cerebral vasospasm, chronic or acute renal failure, non-steroidal antiinflammatory drug induced gastric ulceration, cyclosporin induced nephrotoxicity, endotoxin-induced toxicity, asthma, fibrotic or proliferative diseases, including smooth muscle proliferation, systemic sclerosis, cirrhosis of the liver, adult respiratory distress syndrome, idiopathic cardiomyopathy, lupus erythematosus, diabetic retinopathy or other retinopathies, psoriasis, scleroderma, prostatic hyperplasia, cardiac hyperplasia, restenosis following arterial injury or other pathologic stenosis of blood vessels, LPL-related lipoprotein disorders, transplantation-induced atherosclerosis or atherosclerosis in general, platelet aggregation, thrombosis, cancers, prostate cancer, IL-2 and other cytokine mediated cardiotoxicity and permeability disorders, and nociception, especially treatment of bone pain associated with bone cancer.

Total daily dose administered to a host in single or divided doses may be in amounts, for example, from 0.001 to 1000 mg/kg body weight daily and more usually 0.1 to 100 mg/kg for oral administration or 0.01 to 10 mg/kg for

parenteral administration. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host
5 treated and the particular mode of administration.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion,
10 drug combination, and the severity of the particular disease undergoing therapy.

The compounds of the present invention may be administered orally, parenterally, sublingually, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically
15 acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleagenous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension
20 in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-propanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids
25 such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release
30 the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active

compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

The compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and phosphatidyl cholines (lecithins), both natural and synthetic.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

A representative solid dosage form, for example, a tablet or a capsule, comprises:

Compound of the invention:	35% w/w
Starch, Pregelatinized, NF	50% w/w
Microcrystalline Cellulose, NF	10% w/w
Talc, Powder, USP	5% w/w

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more cardiovascular agents independently selected from diuretics, adrenergic blocking agents, vasodilators, calcium channel blockers, renin inhibitors, angiotensin converting enzyme (ACE) inhibitors, angiotensin II antagonists, potassium channel activators and other cardiovascular agents.

Representative diuretics include hydrochlorothiazide, chlorothiazide, acetazolamide, amiloride, bumetanide, benzthiazide, ethacrynic acid, furosemide, indacrinone, metolazone, spironolactone, triamterene, chlorthalidone and the like or a pharmaceutically acceptable salt thereof.

- 5 Representative adrenergic blocking agents include phentolamine, phenoxybenzamine, prazosin, terazosin, tolazine, atenolol, metoprolol, nadolol, propranolol, timolol, carteolol and the like or a pharmaceutically acceptable salt thereof.

- 10 Representative vasodilators include hydralazine, minoxidil, diazoxide, nitroprusside and the like or a pharmaceutically acceptable salt thereof.

Representative calcium channel blockers include amrinone, bencyclane, diltiazem, fendiline, flunarizine, nicardipine, nimodipine, perhexilene, verapamil, gallopamil, nifedipine and the like or a pharmaceutically acceptable salt thereof.

- 15 Representative renin inhibitors include enalkiren, zankiren, RO 42-5892, PD-134672 and the like or a pharmaceutically acceptable salt thereof.

Representative angiotensin II antagonists include DUP 753, A-81988 and the like.

- 20 Representative ACE inhibitors include captopril, enalapril, lisinopril and the like or a pharmaceutically acceptable salt thereof.

Representative potassium channel activators include pinacidil and the like or a pharmaceutically acceptable salt thereof.

- 25 Other representative cardiovascular agents include sympatholytic agents such as methyl dopa, clonidine, guanabenz, reserpine and the like or a pharmaceutically acceptable salt thereof.

- 30 The compounds of the invention and the cardiovascular agent can be administered at the recommended maximum clinical dosage or at lower doses. Dosage levels of the active compounds in the compositions of the invention may be varied so as to obtain a desired therapeutic response depending on the route of administration, severity of the disease and the response of the patient. The combination can be administered as separate compositions or as a single dosage form containing both agents.


- 35 When administered as a combination, the therapeutic agents can be formulated as separate compositions which are given at the same time or different times, or the therapeutic agents can be given as a single composition.

The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed compounds, processes, compositions and methods. Variations and changes which are obvious to one skilled in the art are intended to be within the scope and nature of the invention which are
5 defined in the appended claims.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

What is claimed is:

1. A compound of the formula:



wherein

Z is $-C(R_{18})(R_{19})-$ or $-C(O)-$ wherein R_{18} and R_{19} are independently selected from hydrogen and loweralkyl;

n is 0 or 1;

R is $-(CH_2)_m-W$ wherein m is an integer from 0 to 6 and W is

(a) $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,

(b) $-PO_3H_2$,

(c) $-P(O)(OH)E$ wherein E is hydrogen, loweralkyl or arylalkyl,

(d) $-CN$,

(e) $-C(O)NHR_{17}$ wherein R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,

(i) hydroxy,

(j) alkoxy,








(k) sulfonamido,

(l) $-C(O)NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl, aryl or dialkylamino,

(m) $-S(O)_2NHC(O)R_{16}$ wherein R_{16} is defined as above,



(n)



R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl,

- 10 haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl,
- 15 alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and (R_{aa})(R_{bb})N-R_{cc}- wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R₁ and R₂ is other than hydrogen;

R₃ is (a)R₄-C(O)-R₅-, R₄-R₅-, R₄-C(O)-R₅- N(R₆)-, R₆-S(O)₂-R₇- or R₂₆-S(O)-R₂₇-

wherein R_5 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene, (iv) $-N(R_{20})-R_8-$ or $-R_{8a}-N(R_{20})-R_8-$

wherein R_8 and R_{8a} are independently selected from the group consisting of alkylene and alkenylene and R_{20} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl or cycloalkylalkyl or (v) $-O-R_9-$ or $-R_{9a}-O-R_9-$ wherein R_9 and R_{9a} are independently selected from alkylene;

R_{5a} is (i) alkylene or (ii) alkenylene;

R_7 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene or (iv) $-N(R_{21})-R_{10}-$ or $-R_{10a}-N(R_{21})-R_{10}-$ wherein R_{10} and R_{10a} are independently selected from the group consisting of alkylene and alkenylene and R_{21} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl;

R_4 and R_6 are independently selected from the group consisting of

(i) $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from

- (1) hydrogen,
- (2) loweralkyl,
- (3) haloalkyl,
- (4) alkoxyalkyl,
- (5) haloalkoxyalkyl,
- (6) alkenyl,
- (7) alkynyl,
- (8) cycloalkyl,
- (9) cycloalkylalkyl,
- (10) aryl,
- (11) heterocyclic,
- (12) arylalkyl,
- (13) (heterocyclic)alkyl,
- (14) hydroxyalkyl,
- (15) alkoxy,
- (16) aminoalkyl,
- (17) trialkylaminoalkyl,
- (18) alkylaminoalkyl,
- (19) dialkylaminoalkyl, and
- (20) carboxyalkyl


(ii) loweralkyl,

(iii) alkenyl,

(iv) alkynyl,

(v) cycloalkyl,

- (vi) cycloalkylalkyl,
- (vii) aryl,
- (viii) arylalkyl,
- (ix) heterocyclic,
- (x) (heterocyclic)alkyl,
- (xi) alkoxyalkyl,
- (xii) hydroxyalkyl,
- (xiii) haloalkyl,
- (xiv) haloalkenyl,
- (xv) haloalkoxyalkyl,
- (xvi) haloalkoxy,
- (xvii) alkoxyhaloalkyl,
- (xviii) alkylaminoalkyl,
- (xix) dialkylaminoalkyl,
- (xx) alkoxy, and



wherein z is 0-5 and R_{7a} is alkylene;

R_{26} is (i) loweralkyl, (ii) haloalkyl, (iii) alkenyl, (iv) alkynyl, (v) cycloalkyl, (vi) cycloalkylalkyl, (vii) aryl, (viii) arylalkyl, (ix) heterocyclic, (x) (heterocyclic)alkyl, (xi) alkoxyalkyl or (xii) alkoxy-substituted haloalkyl; and R_{27} is alkylene or alkenylene;

(b) $R_{22}-O-C(O)-R_{23}-$ wherein R_{22} is a carboxy protecting group or heterocyclic and R_{23} is (i) a covalent bond, (ii) alkylene, (iii) alkenylene or (iv) $-N(R_{24})-R_{25}-$ wherein R_{25} is alkylene and R_{24} is hydrogen or loweralkyl,

- (c) loweralkyl,
- (d) alkenyl,
- (e) alkynyl,
- (f) cycloalkyl,
- (g) cycloalkylalkyl,
- (h) aryl,
- (i) arylalkyl,
- (j) aryloxyalkyl,

- (k) heterocyclic,
- (l) (heterocyclic)alkyl,
- (m) alkoxyalkyl,
- (n) alkoxyalkoxyalkyl, or
- (o) $R_{13}-C(O)-CH(R_{14})-$

wherein R_{13} is amino, alkylamino or dialkylamino and R_{14} is aryl or $R_{15}-C(O)-$ wherein R_{15} is amino, alkylamino or dialkylamino; or a pharmaceutically acceptable salt thereof.

2. The compound according to Claim 1 wherein n is 0 and Z is $-CH_2-$.

3. The compound according to Claim 1 wherein n is 1 and Z is $-CH_2-$.

4. The compound according to Claim 1 wherein n is 0, Z is $-CH_2-$, and R_3 is $R_4-C(O)-R_5-$, $R_6-SO_2-R_7-$ or $R_{26}-S(O)-R_{27}-$ wherein R_4 , R_5 , R_6 , R_7 , R_{26} and R_{27} are as defined therein.

5. The compound according to Claim 1 wherein n is 0, Z is $-CH_2-$, and R_3 is alkoxyalkyl or alkoxyalkoxyalkyl.

6. The compound according to Claim 1 wherein n is 0, Z is $-CH_2-$, and R_3 is $R_4-C(O)-R_5-$ wherein R_4 is $(R_{11})(R_{12})N-$ as defined therein and R_5 is alkylene or R_3 is $R_6-S(O)_2-R_7-$ or $R_{26}-S(O)-R_{27}-$ wherein R_7 is alkylene, R_{27} is alkylene and R_6 and R_{26} are as defined therein.

7. The compound according to Claim 1 wherein n is 0, Z is $-CH_2-$ and R_3 is $R_4-C(O)-N(R_{20})-R_8-$ or $R_6-S(O)_2-N(R_{21})-R_{10}-$ wherein R_8 and R_{10} are alkylene and R_4 , R_6 , R_{20} and R_{21} are as defined therein.

8. The compound according to Claim 1 wherein n is 0, R is tetrazolyl or $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group or R is tetrazolyl or R is $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 and R_2 are independently selected from (i) loweralkyl, (ii) cycloalkyl, (iii) substituted and unsubstituted aryl wherein aryl is phenyl substituted with one, two or three substituents independently selected from

loweralkyl, alkoxy, halo, alkoxyalkoxy and carboxyalkoxy, (iv) substituted or unsubstituted heterocyclic, (v) alkenyl, (vi) heterocyclic (alkyl), (vii) aryloxyalkyl, (viii) arylalkyl, (ix) (N-alkanoyl-N-alkyl)aminoalkyl, and (x) alkylsulfonylamidoalkyl, and R₃ is R₄-C(O)-R₅- wherein R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, heterocyclic, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, aryl and arylalkyl and R₅ is alkylene; or R₃ is R₄-C(O)-N(R₂₀)-R₈- or R₆-S(O)₂-N(R₂₁)-R₁₀- wherein R₄ is loweralkyl, aryl, alkoxy, alkylamino, aryloxy or arylalkoxy and R₆ is loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl, R₈ and R₁₀ are alkylene and R₂₀ and R₂₁ are loweralkyl; or R₃ is R₆-S(O)₂-R₇- or R₂₆-S(O)-R₂₇- wherein R₆ is loweralkyl or haloalkyl, R₇ is alkylene, R₂₆ is loweralkyl and R₂₇ is alkylene.

9. The compound according to Claim 1 wherein n is 0, R is -C(O)₂-G

wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl, haloalkyl or aryl, Z is -CH₂-, R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-t-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonylamidoalkyl, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl, or difluorophenyl, and R₃ is R₄-C(O)-N(R₂₀)-R₈- or R₆-S(O)₂-N(R₂₁)-R₁₀- wherein R₈ and R₁₀ are alkylene, R₂₀ and R₂₁ are loweralkyl, R₄ is loweralkyl, aryl, alkoxy, alkylamino, aryloxy or arylalkoxy and R₆ is loweralkyl, haloalkyl, alkoxyalkyl, aryl or arylalkyl.

10. The compound according to Claim 1 wherein n is 0, R is -C(O)₂-G

wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl, haloalkyl or aryl, Z is -CH₂-, R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl,

- (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl or (xiii) alkylsulfonylamidoalkyl, R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl, or difluorophenyl, and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl, arylalkyl, heterocyclic, hydroxyalkyl, alkoxy, aminoalkyl, and trialkylaminoalkyl.

11. The compound according to Claim 1 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i) loweralkyl (ii) alkenyl, (iii) arylalkyl, (iv) aryloxyalkyl, (v) heterocyclic, (vi) heterocyclic (alkyl), (vii) aryl, (viii) (N-alkanoyl-N-alkyl)aminoalkyl, or (viii) alkylsulfonylamidoalkyl, R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} is loweralkyl, and R_{12} is aryl or arylalkyl.

12. The compound according to Claim 1 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i) phenyl or (ii) substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy, and carboxyalkoxy, R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-

benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R₃ is R₆-S(O)₂-N(R₂₁)-R₁₀- wherein R₁₀ is alkylene, R₆ is loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl and R₂₁ is loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl.

13. The compound according to Claim 1 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl or aryl, Z is -CH₂-, R₁ is substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl or 1,4-benzodioxanyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy and alkoxyalkoxy, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R₃ is alkoxycarbonyl or R₆-S(O)₂-N(R₂₁)-R₁₀- wherein R₁₀ is alkylene, R₆ is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R₂₁ is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl.

14. The compound according to Claim 1 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl or haloalkyl, Z is -CH₂-, R₁ is substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl, aryl, arylalkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

15. The compound according to Claim 1 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$

wherein R_{16} is loweralkyl or haloalkyl, Z is $-CH_2-$, R_1 is loweralkyl, alkoxyalkyl, or alkenyl, R_2 is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$

wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N$

wherein R_{11} and R_{12} are independently selected from loweralkyl, aryl

hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, heterocyclic, and arylalkyl.

16. The compound according to Claim 1 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1 is

substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, R_2 is 1,3-benzodioxolyl, 1,4-

benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N$ wherein R_{11} and R_{12} are

independently selected from loweralkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, aryl, and heterocyclic.

17. The compound according to Claim 1 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1 is

substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, R_2 is 1,3-benzodioxolyl, 1,4-

benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$


wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N$ wherein R_{11} is loweralkyl and R_{12} is aryl.

18. The compound according to Claim 1 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, Z is -CH₂-, R₁ is substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 2-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ is alkyl and R₁₂ is selected from aryl, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

19. A compound according to Claim 1 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, Z is -CH₂-, R₁ is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, aryalkyl, aryl, (N-alkanoyl-N-alkyl)aminoalkyl, or alkylsulfonylamidoalkyl, and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from alkyl, aryl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

20. A compound according to Claim 1 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, Z is -CH₂-, R₁ is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, aryalkyl, aryl, (N-alkanoyl-N-alkyl)aminoalkyl, or alkylsulfonylamidoalkyl, and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from alkyl, aryl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic, with the proviso that one or R₁₁ and R₁₂ is alkyl.

21. The compound according to Claim 1 of the formula:



wherein

Z is $-C(R_{18})(R_{19})-$ or $-C(O)-$ wherein R_{18} and R_{19} are independently selected from hydrogen and loweralkyl;

5 n is 0 or 1;

R is $-(CH_2)_m-W$ wherein m is an integer from 0 to 6 and W is

(a) $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,

(b) $-PO_3H_2$,

(c) $-P(O)(OH)E$ wherein E is hydrogen, loweralkyl or arylalkyl,

10 (d) $-CN$,

(e) $-C(O)NHR_{17}$ wherein R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,

15 (i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

(l) $-C(O)NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl, aryl or dialkylamino,

20 (m) $-S(O)_2NHC(O)R_{16}$ wherein R_{16} is defined as above,



(n)




(o)



(p)



(q)



- 5 R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than
- 15 hydrogen;
 R_3 is (a) $R_4-C(O)-R_5$, R_4-R_{5a} , $R_6-S(O)_2-R_7$ or $R_{26}-S(O)-R_{27}$ wherein R_5 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene, (iv) $-N(R_{20})-R_8$ or $-R_{8a}-N(R_{20})-R_8$ wherein R_8 and R_{8a} are independently selected from the group consisting of
- 20 alkylene and alkenylene and R_{20} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl or cycloalkylalkyl or (v) $-O-R_9$ or $-R_{9a}-O-R_9$ wherein R_9 and R_{9a} are independently selected from alkylene;
 R_{5a} is (i) alkylene or (ii) alkenylene;
- 25 R_7 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene or (iv) $-N(R_{21})-R_{10}$ or $-R_{10a}-N(R_{21})-R_{10}$

wherein R_{10} and R_{10a} are independently selected from the group consisting of alkylene and alkenylene and R_{21} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl;

R_4 and R_6 are independently selected from the group consisting of

5 (i) $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from

- (1) hydrogen,
- (2) loweralkyl,
- (3) haloalkyl,
- 10 (4) alkoxyalkyl,
- (5) haloalkoxyalkyl,
- (6) alkenyl,
- (7) alkynyl,
- (8) cycloalkyl,
- 15 (9) cycloalkylalkyl,
- (10) aryl,
- (11) heterocyclic,
- (12) arylalkyl,
- (13) (heterocyclic)alkyl,
- 20 (14) hydroxyalkyl,
- (15) alkoxy,
- (16) aminoalkyl, and
- (17) trialkylaminoalkyl,

(ii) loweralkyl,

25 (iii) alkenyl,

(iv) alkynyl,

(v) cycloalkyl,

(vi) cycloalkylalkyl,

(vii) aryl,

30 (viii) arylalkyl,

(ix) heterocyclic,

(x) (heterocyclic)alkyl,

(xi) alkoxyalkyl,

(xii) hydroxyalkyl,


35 (xiii) haloalkyl,

(xiv) haloalkenyl,

(xv) haloalkoxyalkyl,

- (xvi) haloalkoxy,
- (xvii) alkoxyhaloalkyl,
- (xviii) alkylaminoalkyl,
- (xix) dialkylaminoalkyl,
- (xx) alkoxy, and

5



(xxi)

wherein z is 0-5 and R_{7a} is alkylene;

R₂₆ is (i) loweralkyl, (ii) haloalkyl, (iii) alkenyl, (iv) alkynyl, (v) cycloalkyl, (vi) cycloalkylalkyl, (vii) aryl, (viii) arylalkyl, (ix) heterocyclic, (x) (heterocyclic)alkyl, (xi) alkoxyalkyl or (xii) alkoxy-substituted haloalkyl; and R₂₇ is alkylene or alkenylene;

(b) R₂₂-O-C(O)-R₂₃- wherein R₂₂ is a carboxy protecting group or heterocyclic and R₂₃ is (i) a covalent bond, (ii) alkylene, (iii) alkenylene or (iv) -N(R₂₄)-R₂₅- wherein R₂₅ is alkylene and R₂₄ is hydrogen or loweralkyl,

(c) loweralkyl,

(d) alkenyl,

(e) alkynyl,

(f) cycloalkyl,

(g) cycloalkylalkyl,

(h) aryl,

(i) arylalkyl,

(j) aryloxyalkyl,

(k) heterocyclic,

(l) (heterocyclic)alkyl,

(m) alkoxyalkyl,

(n) alkoxyalkoxyalkyl, or

(o) R₁₃-C(O)-CH(R₁₄)-

wherein R₁₃ is amino, alkylamino or dialkylamino and R₁₄ is aryl or R₁₅-C(O)- wherein R₁₅ is amino, alkylamino or dialkylamino;

or a pharmaceutically acceptable salt thereof.

22. The compound according to Claim 21 wherein n is 0 and Z is -CH₂-.

23. The compound according to Claim 21 wherein n is 1 and Z is -CH₂-.

24. The compound according to Claim 21 wherein n is 0, Z is -CH₂-, and R₃ is R₄-C(O)-R₅-, R₆-SO₂-R₇- or R₂₆-S(O)-R₂₇- wherein R₄, R₅, R₆, R₇, R₂₆ and R₂₇ are as defined therein.

25. The compound according to Claim 21 wherein n is 0, Z is -CH₂-, and R₃ is alkoxyalkyl or alkoxyalkoxyalkyl.

26. The compound according to Claim 21 wherein n is 0, Z is -CH₂-, and R₃ is R₄-C(O)-R₅- wherein R₄ is (R₁₁)(R₁₂)N- as defined therein and R₅ is alkylene or R₃ is R₆-S(O)₂-R₇- or R₂₆-S(O)-R₂₇- wherein R₇ is alkylene, R₂₇ is alkylene and R₆ and R₂₆ are as defined therein.

27. The compound according to Claim 21 wherein n is 0, Z is -CH₂- and R₃ is R₄-C(O)-N(R₂₀)-R₈- or R₆-S(O)₂-N(R₂₁)-R₁₀- wherein R₈ and R₁₀ are alkylene and R₄, R₆, R₂₀ and R₂₁ are as defined therein.

28. The compound according to Claim 21 wherein n is 0, R is tetrazolyl or -C(O)₂-G wherein G is hydrogen or a carboxy protecting group or R is tetrazolyl or R is -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl, haloalkyl or aryl, Z is -CH₂-, R₁ and R₂ are independently selected from (i) loweralkyl, (ii) cycloalkyl, (iii) substituted and unsubstituted aryl wherein aryl is phenyl substituted with one, two or three substituents independently selected from loweralkyl, alkoxy, halo, alkoxyalkoxy and carboxyalkoxy and (iv) substituted or unsubstituted heterocyclic, (v) alkenyl, (vi) heterocyclic (alkyl), (vii) aryloxyalkyl, (viii) aryalkyl, (ix) (N-alkanoyl-N-alkyl)aminoalkyl, and (x) alkylsulfonylamidoalkyl, and R₃ is R₄-C(O)-R₅- wherein R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, heterocyclic, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, aryl and aryalkyl and R₅ is alkylene; or

R_3 is $R_4-C(O)-N(R_{20})-R_8-$ or $R_6-S(O)_2-N(R_{21})-R_{10}-$ wherein R_4 is loweralkyl, aryl, alkoxy, alkylamino, aryloxy or arylalkoxy and R_6 is loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl, R_8 and R_{10} are alkylene and R_{20} and R_{21} are loweralkyl; or

- 5 R_3 is $R_6-S(O)_2-R_7-$ or $R_{26}-S(O)-R_{27}-$ wherein R_6 is loweralkyl or haloalkyl, R_7 is alkylene, R_{26} is loweralkyl and R_{27} is alkylene.

10

29. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i)

- loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, 15 (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or 20 dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonfylamidoalkyl, R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 25 benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-N(R_{20})-R_8-$ or $R_6-S(O)_2-N(R_{21})-R_{10}-$ wherein R_8 and R_{10} are alkylene, R_{20} and R_{21} are loweralkyl, R_4 is loweralkyl, aryl, alkoxy, alkylamino, aryloxy or arylalkoxy and 30 R_6 is loweralkyl, haloalkyl, alkoxyalkyl, aryl or arylalkyl.

30. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is (i)

- 35 loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl,

4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) aryalkyl, (x) aryoxylalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonamidoalkyl, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ and R₁₂ are independently selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl and arylalkyl, heterocyclic, hydroxyalkyl, alkoxy, aminoalkyl, and trialkylaminoalkyl.

31. The compound according to Claim 21 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl, haloalkyl or aryl, Z is -CH₂-, R₁ is (i) loweralkyl or (ii) alkenyl, (iii) aryalkyl, (iv) aryoxylalkyl, (v) heterocyclic (alkyl), (vi) aryl, (vii) (N-alkanoyl-N-alkyl)aminoalkyl, or (viii) alkylsulfonamidoalkyl, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R₃ is R₄-C(O)-R₅- wherein R₅ is alkylene and R₄ is (R₁₁)(R₁₂)N- wherein R₁₁ is loweralkyl and R₁₂ is aryl or arylalkyl.

32. The compound according to Claim 21 wherein n is 0, R is -C(O)₂-G wherein G is hydrogen or a carboxy protecting group, tetrazolyl or -C(O)-NHS(O)₂R₁₆ wherein R₁₆ is loweralkyl, haloalkyl or aryl, Z is -CH₂-, R₁ is (i) phenyl or (ii) substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the

substituent is selected from loweralkyl, alkoxy and halogen and R_3 is $R_6-S(O)_2-N(R_{21})-R_{10}$ wherein R_{10} is alkylene, R_6 is loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl and R_{21} is loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl.

5

33 The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl or aryl, Z is $-CH_2-$, R_1 is substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl or 1,4-benzodioxanyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy and alkoxyalkoxy, R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R_3 is alkoxyalkyl or $R_6-S(O)_2-N(R_{21})-R_{10}$ wherein R_{10} is alkylene, R_6 is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{21} is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl.

20

34. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl or haloalkyl, Z is $-CH_2-$, R_1 is substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and carboxyalkoxy, R_2 is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from loweralkyl, aryl arylalkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

35

35. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, tetrazolyl or $-C(O)-NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl or haloalkyl, Z is $-CH_2-$, R_1 is

loweralkyl, alkoxyalkyl or alkenyl, R_2 is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from

5 loweralkyl, aryl, arylalkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic.

36. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1

10 is substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and

15 carboxyalkoxy, R_2 is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from

20 loweralkyl.

37. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1

25 is substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 2-fluorophenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-ethylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from alkoxy, alkoxyalkoxy and

30 carboxyalkoxy, R_2 is 1,3-benzodioxolyl, 1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} is loweralkyl and R_{12} is aryl.

38. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1

35 is substituted or unsubstituted 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 3-fluorophenyl, 2-fluorophenyl, 3-fluoro-4-ethoxyphenyl, 4-methoxymethoxyphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or

dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, R_2 is substituted or

unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen and R_3 is $R_6-S(O)_2-N(R_{21})-R_{10}$ wherein R_{10} is alkylene, R_6 is loweralkyl, haloalkyl, alkoxyalkyl or haloalkoxyalkyl and R_{21} is loweralkyl, haloalkyl or alkoxyalkyl.

39. The compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1 is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, aryalkyl, aryl, (N-alkanoyl-N-alkyl)aminoalkyl, or alkylsulfonylamidoalkyl, and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from alkyl, aryl, hydroxyalkyl, alkoxy, aminoalkyl, and heterocyclic.

40. A compound according to Claim 21 wherein n is 0, R is $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group, Z is $-CH_2-$, R_1 is loweralkyl, alkenyl, heterocyclic (alkyl), aryloxyalkyl, aryalkyl, aryl, (N-alkanoyl-N-alkyl)aminoalkyl, or alkylsulfonylamidoalkyl, and R_3 is $R_4-C(O)-R_5$ wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from alkyl, aryl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, and heterocyclic, with the proviso that one or R_{11} and R_{12} is alkyl

41. A compound selected from the group consisting of
trans-trans-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(3-(N-propyl-N-n-pentanesulfonylamino)propyl)pyrrolidine-3-carboxylic acid;
trans, trans-2-(4-Methoxymethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;
trans, trans-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;
trans, trans-2-(3,4-Dimethoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-hexanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;
trans, trans-2-(4-Propoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-n-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;

- trans,trans*-2-(3,4-Difluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(((N,N-di(*n*-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3,4-Difluorophenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-*n*-pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;
- 5 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-*n*-hexanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(3-chloropropanesulfonyl)amino)ethyl)-pyrrolidine-3-carboxylic acid;
- 10 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(3-chloropropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(4-methylbutanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;
- 15 *trans,trans*-2-(4-Methoxy-3-fluorophenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(*n*-pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2,2,3,3,3-pentafluoropropoxyethanesulfonyl)-amino)ethyl)pyrrolidine-3-carboxylic acid;
- 20 *trans,trans*-2-(1,4-Benzodioxan-6-yl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(*n*-pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(pentanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(3-chloropropanesulfonyl)amino)-ethyl)pyrrolidine-3-carboxylic acid;
- 25 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(pentanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid;
- 30 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-((2,2,2-trifluoroethoxyethane)sulfonyl)amino)-ethyl)pyrrolidine-3-carboxylic acid;
- 35 *trans,trans*-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-((2,2,2-trifluoroethoxyethane)sulfonyl)amino)-ethyl)pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-(2-methoxyethyl)-N-(butanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-(2-methylpropanesulfonyl)amino)ethyl)pyrrolidine-3-carboxylic acid; and

trans,trans-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-isobutyl-N-(butanesulfonylamino)ethyl)pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-Methylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Tetrahydro-2H-pyran)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,4-Trimethyl-3-pentenyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2,-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-((N-4-heptyl-N-(2-methyl-3-fluorophenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(1,3-Dioxo-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

trans,trans-2-(2-(2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(2-(2-Oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

(2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid

trans,trans-2-(2-(1,3-Dioxol-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl))aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;

- trans,trans*-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2,2-dimethylpentyl)-4-(2,3-dihydro-benzofuran-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 5 *trans,trans*-2-(2,2-Dimethyl-2-(1,3-dioxolan-2-yl)ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(2-Methoxyphenyl)-ethyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 10 *trans,trans*-2-(2,2-Dimethyl-3-(E)-pentenyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(2-pyridyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- (2S,3R,4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 15 (2S,3R,4S)-2-(2-(2-oxopyrrolidin-1-yl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N-4-heptyl-N-(4-fluoro-3-methylphenyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(2-(1-pyrazolyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 20 *trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-((N-butyl-N-(4-dimethylaminobutyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;
- (2R,3R,4S)-2-(3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonylamino)ethyl)-pyrrolidine-3-carboxylic acid;
- 25 (2S,3R,4S)-2-(2,2-Dimethylpentyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- (2S,3R,4S)-2-(2,2-Dimethylpent-(E)-3-enyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- 30 (2S,3R,4S)-2-(2,2-Dimethylpent-(E)-3-enyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid;
- (2S,3R,4S)-2-((2-Methoxyphenoxy)-methyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid;
- 35 (2S,3R,4S)-2-(2-(2-Methoxyphenyl)ethyl)-4-(1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)pyrrolidine-3-carboxylic acid;
- trans,trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-((bis-(o-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-methoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-(2,2-dimethyl-1-phenylpropyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-methoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-((bis-(o-tolyl)methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-isopropoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-(2,2-dimethyl-1-phenylpropyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;


trans,trans-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-(3,3-dimethyl-1-phenylbutyl)-1-amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-[4-(2-isopropoxyethoxy)phenyl]-4-(1,3-benzodioxol-5-yl)-1-(N-((1-(o-tolyl)-1-(o-ethylphenyl)-methyl)amino)carbonylmethyl)-pyrrolidine-3-carboxylic acid;

trans,trans-2-(4-(2-(2-propoxy)ethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-N-phenyl-N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid; and

trans,trans-2-(4-(2-methoxyethoxy)phenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-phenyl-N-t-butylhydrazino carbonylmethyl)-pyrrolidine-3-carboxylic acid; or a pharmaceutically acceptable salt thereof.

42. A compound of the formula:



wherein n is 0 or 1;

m is 0 to 6;

W is (a) -C(O)₂-G where G is hydrogen or a carboxy protecting group, (b) -PO₃H₂,

(c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,

(d) -CN,

(e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,









(h) tetrazolyl,

(i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

- (l) $-C(O)NHS(O)_2R_{16}$ where R_{16} is loweralkyl, haloalkyl, phenyl or dialkylamino,
 (m) $-S(O)_2NHC(O)R_{16}$,




R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl,

alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and (R_{aa})(R_{bb})N-R_{cc}- wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R₁ and R₂ is other than hydrogen; or a salt thereof.

43. The compound of Claim 42 wherein m is zero or 1; W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; or the substantially pure (+)- or (-)-isomer thereof.

44. The compound of Claim 42 wherein n and m are both 0; W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonylamidoalkyl, and R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen; or the substantially pure (+)- or (-)-isomer thereof.

45. The compound according to Claim 42 of the formula:



wherein n is 0 or 1;

m is 0 to 6;

- 5 W is (a) $-C(O)_2-G$ where G is hydrogen or a carboxy protecting group, (b) $-PO_3H_2$,

(c) $-P(O)(OH)E$ where E is hydrogen, loweralkyl or arylalkyl,

(d) $-CN$,

(e) $-C(O)NHR_{17}$ where R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,


(i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

(l) $-C(O)NHS(O)_2R_{16}$ where R_{16} is loweralkyl, haloalkyl, phenyl or dialkylamino,

(m) $-S(O)_2NHC(O)R_{16}$,








(n)



(o)



(p)



R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen; or a salt thereof.

46. The compound according to Claim 45 wherein

m is zero or 1;
 W is $-CO_2-G$ wherein G is hydrogen or a carboxy protecting group; or the substantially pure (+)- or (-)-isomer thereof.

47. The compound according to Claim 45 wherein


n and m are both 0;

W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonylamidoalkyl, and R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen; or the substantially pure (+)- or (-)-isomer thereof.

48. The substantially pure compound (+)-*trans,trans*-2-(4-Methoxyphenyl)-4-(1,3-benzodioxo-5-yl)pyrrolidine-3-carboxylic acid; or a salt or ester thereof.

49. The substantially pure compound (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(*N,N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid; or a salt or ester thereof.

50. A compound of the formula



5 wherein n is 0 or 1;

m is 0 to 6;

R_{5b} is alkylene;

Q is a leaving group;

W is (a) -C(O)₂-G where G is hydrogen or a carboxy protecting group, (b)

10 -PO₃H₂,

(c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,

(d) -CN,

(e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,

(f) alkylaminocarbonyl,

15 (g) dialkylaminocarbonyl,

(h) tetrazolyl,

(i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

20 (l) -C(O)NHS(O)₂R₁₆ where R₁₆ is loweralkyl, haloalkyl, phenyl or

dialkylamino,







(m) -S(O)₂NHC(O)R₁₆.



(n)



(o)



R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl,

alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl,

haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl,

10 cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl,

dialkylaminocarbonylalkyl, aminocarbonylalkenyl,

alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl,

aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl,

alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and

15 (R_{aa})(R_{bb})N-R_{cc}- wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl

and R_{cc} is alkylene, with the proviso that one or both of R₁ and R₂ is other

than hydrogen;

or a salt thereof.

51. The compound according to Claim 50 wherein

m is zero or 1;

R_{5b} is alkylene;

Q is a leaving group; and

- 5 W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; or the substantially pure (+)- or (-)-isomer thereof.

52. The compound according to Claim 50 wherein

n and m are both 0;

- 10 R_{5b} is alkylene;

Q is a leaving group;

W is -CO₂-G wherein G is hydrogen or a carboxy protecting group;

and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl,

- 15 (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl,


- 20 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy and R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl,

1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or

- 25 difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonylamidoalkyl; or the substantially pure (+)- or (-)-isomer thereof.

53. The compound according to Claim 50 of the formula

30



wherein n is 0 or 1;

m is 0 to 6;

R_{5b} is alkylene;

Q is a leaving group;

W is (a) -C(O)₂-G where G is hydrogen or a carboxy protecting group, (b)

5 -PO₃H₂,

(c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,

(d) -CN,

(e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,

(f) alkylaminocarbonyl,

10 (g) dialkylaminocarbonyl,

(h) tetrazolyl,

(i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

15 (l) -C(O)NHS(O)₂R₁₆ where R₁₆ is loweralkyl, haloalkyl, phenyl or dialkylamino,

(m) -S(O)₂NHC(O)R₁₆,



(n)



(o)






(p)



(q)



(r)



R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl,

- 5 alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl,
- 10 aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ - wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen; or a salt thereof.


54. The compound according to Claim 53 wherein m is zero or 1; R_{5b} is alkylene; Q is a leaving group; W is $-CO_2-G$ wherein G is hydrogen or a carboxy protecting group; or the substantially pure (+)- or (-)-isomer thereof.

55. The compound according to Claim 53 wherein n and m are both 0; R_{5b} is alkylene; Q is a leaving group; W is $-CO_2-G$ wherein G is hydrogen or a carboxy protecting group; and R_1 is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl,

4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-*t*-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy and R_2 is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl,

- 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen, (ix) aryalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-
5 N-alkyl)aminoalkyl, or (xiii) alkylsulfonylamidoalkyl; or the substantially pure (+)- or (-)-isomer thereof.

56. A compound of the formula



- 5 wherein n is 0 or 1; m is 0 to 6; R_{5b} is alkylene;
R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl,
cycloalkyl, cycloalkylalkyl, aryl or arylalkyl;
W is (a) -C(O)₂-G where G is hydrogen or a carboxy protecting group, (b)
-PO₃H₂,

- 10 (c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,
(d) -CN,
(e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,
(f) alkylaminocarbonyl,
(g) dialkylaminocarbonyl,
15 (h) tetrazolyl,
(i) hydroxy,
(j) alkoxy,
(k) sulfonamido,
(l) -C(O)NHS(O)₂R₁₆ where R₁₆ is loweralkyl, haloalkyl, phenyl or
20 dialkylamino,
(m) -S(O)₂NHC(O)R₁₆,








(n)



(o)



(p)



R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidoalkyl, heterocyclic, (heterocyclic)alkyl and (R_{aa})(R_{bb})N-R_{cc} wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R₁ and R₂ is other than hydrogen; or a salt thereof.

57. The compound according to Claim 56 wherein

m is zero or 1;

R_{5b} is alkylene;

R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl; and
W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; or the substantially pure (+)- or (-)-isomer thereof.

58. The compound according to Claim 56 wherein

n and m are both 0;

R_{5b} is alkylene;

R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl,

5 cycloalkyl, cycloalkylalkyl, aryl or arylalkyl;

W is -CO₂-G wherein G is hydrogen or a carboxy protecting group;

and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi)

pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-

fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-

10 trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-

4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-t-

butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein

the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and

carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-

15 N-alkyl)aminoalkyl, or (xiii) alkylsulfonylamidoalkyl, and R₂ is substituted or

unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-


methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl,

dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected

from loweralkyl, alkoxy and halogen; or

20 the substantially pure (+)- or (-)-isomer thereof.

59. The compound according to Claim 56 of the formula



25 wherein n is 0 or 1; m is 0 to 6; R_{5b} is alkylene; R_{20a} is hydrogen, loweralkyl, alkenyl,

haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl;

W is (a) -C(O)Z-G where G is hydrogen or a carboxy protecting group, (b)

-PO₃H₂,

30 (c) -P(O)(OH)E where E is hydrogen, loweralkyl or arylalkyl,

(d) -CN,

(e) -C(O)NHR₁₇ where R₁₇ is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

(h) tetrazolyl,


(i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

(l) $-\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}_{16}$ where R_{16} is loweralkyl, haloalkyl, phenyl or dialkylamino,

(m) $-\text{S}(\text{O})_2\text{NHC}(\text{O})\text{R}_{16}$.



(n) ,



(o) ,



(p) ,



(q) ,



(r) ,



(s) ,



(t) , or



(u) ; and

5

10

15

R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonfylamidoalkyl, heterocyclic, (heterocyclic)alkyl and (R_{aa})(R_{bb})N-R_{cc}- wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R₁ and R₂ is other than hydrogen; or a salt thereof.

60. The compound according to Claim 59 wherein

m is zero or 1;

R_{5b} is alkylene;

R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl; and W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; or the substantially pure (+)- or (-)-isomer thereof.

61. The compound according to Claim 58 wherein

n and m are both 0; R_{5b} is alkylene; R_{20a} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl or arylalkyl; W is -CO₂-G wherein G is hydrogen or a carboxy protecting group; and R₁ is (i) loweralkyl, (ii) alkenyl, (iii) alkoxyalkyl, (iv) cycloalkyl, (v) phenyl, (vi) pyridyl, (vii) furanyl or (viii) substituted or unsubstituted 4-methoxyphenyl, 4-fluorophenyl, 3-fluorophenyl, 4-ethoxyphenyl, 4-ethylphenyl, 4-methylphenyl, 4-trifluoromethylphenyl, 4-pentafluoroethylphenyl, 3-fluoro-4-methoxyphenyl, 3-fluoro-4-ethoxyphenyl, 2-fluorophenyl, 4-methoxymethoxyphenyl, 4-hydroxyphenyl, 4-t-butylphenyl, 1,3-benzodioxolyl, 1,4-benzodioxanyl or dihydrobenzofuranyl wherein the substituent is selected from loweralkyl, haloalkyl, alkoxy, alkoxyalkoxy and carboxyalkoxy, (ix) arylalkyl, (x) aryloxyalkyl, (xi) heterocyclic (alkyl), (xii) (N-alkanoyl-N-alkyl)aminoalkyl, or (xiii) alkylsulfonfylamidoalkyl, and R₂ is substituted or unsubstituted 1,3-benzodioxolyl, 7-methoxy-1,3-benzodioxolyl, 1,4-benzodioxanyl, 8-methoxy-1,4-benzodioxanyl, dihydrobenzofuranyl, benzofuranyl, 4-methoxyphenyl, dimethoxyphenyl, fluorophenyl or difluorophenyl wherein the substituent is selected from loweralkyl, alkoxy and halogen; or the substantially pure (+)- or (-)-isomer thereof.

62. A pharmaceutical composition for antagonizing the action of endothelin comprising a therapeutically effective amount of the compound of Claim 1 and a pharmaceutically acceptable carrier.

63. A pharmaceutical composition for antagonizing the action of endothelin comprising a therapeutically effective amount of the compound of Claim 21 and a pharmaceutically acceptable carrier.

64. A pharmaceutical composition for antagonizing the action of endothelin comprising a therapeutically effective amount of (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(*N*,*N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid and a pharmaceutically acceptable carrier.

65. A pharmaceutical composition for antagonizing the action of endothelin comprising a therapeutically effective amount of (2*S*,3*R*,4*S*)-2-3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(*N*-propyl-*N*-pentanesulfonyl)ethyl)-pyrrolidine-3-carboxylic acid and a pharmaceutically acceptable carrier.

66. A method for antagonizing the action of endothelin comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.

67. A method for antagonizing the action of endothelin comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 21.

68. A method for antagonizing the action of endothelin comprising administering to a mammal in need of such treatment a therapeutically effective amount of (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(*N*,*N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid.

69. A method for antagonizing the action of endothelin comprising administering to a mammal in need of such treatment a therapeutically effective amount of (2*S*,3*R*,4*S*)-2-3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(*N*-propyl-*N*-pentanesulfonyl)ethyl)-pyrrolidine-3-carboxylic acid.

70. A method for treating hypertension, congestive heart failure, restenosis following arterial injury, renal failure, cancer, colitis, reperfusion injury, angina, pulmonary hypertension, migraine, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.

71. A method for treating coronary angina, cerebral vasospasm, acute and chronic renal failure, gastric ulceration, cyclosporin-induced nephrotoxicity, endotoxin-induced toxicity, asthma, LPL-related lipoprotein disorders, proliferative diseases, acute or chronic pulmonary hypertension, platelet aggregation, thrombosis, IL-2 mediated cardiotoxicity, nociception, colitis, vascular permeability disorders, ischemia-reperfusion injury, raynaud's disease and migraine comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of claim 1.

72. A method for treating hypertension, congestive heart failure, restenosis following arterial injury, renal failure, cancer, colitis, reperfusion injury, angina, pulmonary hypertension, migraine, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 21.

73. A method for treating hypertension, congestive heart failure, restenosis following arterial injury, renal failure, cancer, colitis, reperfusion injury, angina, pulmonary hypertension, migraine, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(*N*,*N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid.

74. A method for treating hypertension, congestive heart failure, restenosis following arterial injury, renal failure, cancer, colitis, reperfusion injury, angina, pulmonary hypertension, migraine, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of (2*S*,3*R*,4*S*)-2-3-Fluoro-4-methoxyphenyl)-4-(1,3-

benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonyl)ethyl)-pyrrolidine-3-carboxylic acid.

75. A method for treating coronary angina, cerebral vasospasm, acute and chronic renal failure, gastric ulceration, cyclosporin-induced nephrotoxicity, endotoxin-induced toxicity, asthma, LPL-related lipoprotein disorders, proliferative diseases, acute or chronic pulmonary hypertension, platelet aggregation, thrombosis, IL-2 mediated cardiotoxicity, nociception, colitis, vascular permeability disorders, ischemia-reperfusion injury, raynaud's disease and migraine comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of claim 21.

76. A method for treating coronary angina, cerebral vasospasm, acute and chronic renal failure, gastric ulceration, cyclosporin-induced nephrotoxicity, endotoxin-induced toxicity, asthma, LPL-related lipoprotein disorders, proliferative diseases, acute or chronic pulmonary hypertension, platelet aggregation, thrombosis, IL-2 mediated cardiotoxicity, nociception, colitis, vascular permeability disorders, ischemia-reperfusion injury, raynaud's disease and migraine comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(N,N-di(n-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid.


77. A method for treating coronary angina, cerebral vasospasm, acute and chronic renal failure, gastric ulceration, cyclosporin-induced nephrotoxicity, endotoxin-induced toxicity, asthma, LPL-related lipoprotein disorders, proliferative diseases, acute or chronic pulmonary hypertension, platelet aggregation, thrombosis, IL-2 mediated cardiotoxicity, nociception, colitis, vascular permeability disorders, ischemia-reperfusion injury, raynaud's disease and migraine comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of (2*S*,3*R*,4*S*)-2-3-Fluoro-4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(2-(N-propyl-N-pentanesulfonyl)ethyl)-pyrrolidine-3-carboxylic acid.

78. A method for treating treating hypertension, congestive heart failure, restenosis following arterial injury, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1 in combination with one or more cardiovascular agents.


79. A method for treating treating hypertension, congestive heart failure, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 21 in combination with one or more cardiovascular agents.

80. A method for treating treating hypertension, congestive heart failure, cerebral or myocardial ischemia or atherosclerosis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of (2*S*,3*R*,4*S*)-2-(2,2-Dimethylpentyl)-4-(7-methoxy-1,3-benzodioxol-5-yl)-1-(*N*,*N*-di(*n*-butyl)aminocarbonylmethyl)-pyrrolidine-3-carboxylic acid in combination with one or more cardiovascular agents.

81. A process for the preparation of a compound of the formula:



wherein E is a carboxy-protecting group and R₁ and R₂ are independently selected from loweralkyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidoalkyl, heterocyclic and (heterocyclic)alkyl; or a salt thereof, comprising a) catalytic hydrogenation of a compound of the formula:




wherein E, R₁ and R₂ are defined as above and b) catalytic hydrogenation of the product of step a) in the presence of an acid or a mixture of acids.

82. The process of Claim 71 wherein E is loweralkyl, R₁ is aryl and R₂ is heterocyclic.

83. The process of Claim 71 wherein the hydrogenation catalyst is Raney nickel and the acid is a mixture of acetic acid and trifluoroacetic acid.


84. The process of Claim 71 wherein E is loweralkyl, R₁ is 4-methoxyphenyl and R₂ is 1,3-benzodioxol-5-yl.

85. A process for the preparation of a compound of the formula:



wherein E is a carboxy-protecting group and R₁ and R₂ are independently selected from loweralkyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic and (heterocyclic)alkyl; or a salt thereof, comprising

a) catalytic hydrogenation of a compound of the formula:



wherein E, R₁ and R₂ are defined as above,

b) catalytic hydrogenation of the product of step a) in the presence of an acid or a mixture of acids, and


c) epimerization of the product of step b) with a base.

86. The process of Claim 75 wherein E is loweralkyl, R₁ is aryl and R₂ is heterocyclic.

87. The process of Claim 75 wherein the hydrogenation catalyst is Raney nickel and the acid is a mixture of acetic acid and trifluoroacetic acid.

88. The process of Claim 75 wherein E is loweralkyl, R₁ is 4-methoxyphenyl and R₂ is 1,3-benzodioxol-5-yl.

89. A process for the preparation of a compound of the formula:




wherein E is a carboxy-protecting group, R₁ and R₂ are independently selected from loweralkyl, alkoxyalkyl, alkoxyalkonylalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidalkyl, heterocyclic and (heterocyclic)alkyl and R₃ is R₄-C(O)-R₅-

wherein R_5 is alkylene and R_4 is $(R_{11})(R_{12})N$ - wherein R_{11} and R_{12} are independently selected from

- (1) loweralkyl,
- (2) haloalkyl,
- (3) alkoxyalkyl,
- (4) haloalkoxyalkyl,
- (5) alkenyl,
- (6) alkynyl,
- (7) cycloalkyl,
- (8) cycloalkylalkyl,
- (9) aryl,
- (10) heterocyclic,
- (11) arylalkyl and
- (12) (heterocyclic)alkyl;
- (13) hydroxyalkyl,
- (14) alkoxy,
- (15) aminoalkyl, and
- (16) trialkylaminoalkyl,

or a salt thereof, comprising

a) catalytic hydrogenation of a compound of the formula:



wherein E, R_1 and R_2 are defined as above,

b) catalytic hydrogenation of the product of step a) in the presence of an acid or a mixture of acids,

c) epimerization of the product of step b) with a base and


d) alkylation of the product of step c) with a compound of the formula R_3-X wherein X is a leaving group and R_3 is defined as above.

90. The process of Claim 79 wherein E is loweralkyl, R_1 is aryl, R_2 is heterocyclic and R_3 is $-CH_2C(O)NR_{11}R_{12}$ wherein R_{11} and R_{12} are independently selected from the group consisting of loweralkyl.


91. The process of Claim 79 wherein the hydrogenation catalyst is Raney nickel and the acid is a mixture of acetic acid and trifluoroacetic acid.

5 92. The process of Claim 79 wherein E is loweralkyl, R_1 is 4-methoxyphenyl, R_2 is 1,3-benzodioxol-5-yl, R_3 is $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{n-Bu})_2$ and X is a halogen or sulfonate leaving group.

10 93. A process for the preparation of the substantially pure (+)-trans,trans optical isomer of the compound of the formula:




15 wherein E is loweralkyl, R_1 is 4-methoxyphenyl and R_2 is 1,3-benzodioxol-5-yl, or a salt thereof, comprising reacting a mixture of the (+) and (-) enantiomers of the compound of the formula:



20 wherein E is loweralkyl, R_1 is 4-methoxyphenyl and R_2 is 1,3-benzodioxol-5-yl with S-(+)- mandelic acid and separating the mandelate salt of the (+)-trans,trans optical isomer.

25 94. A compound of the formula:



wherein

Z is $-C(R_{18})(R_{19})-$ or $-C(O)-$ wherein R_{18} and R_{19} are independently selected from

5 hydrogen and loweralkyl;

n is 0 or 1;

R is $-(CH_2)_m-W$ wherein m is an integer from 0 to 6 and W is

(a) $-C(O)_2-G$ wherein G is hydrogen or a carboxy protecting group,

(b) $-PO_3H_2$,

10 (c) $-P(O)(OH)E$ wherein E is hydrogen, loweralkyl or arylalkyl,

(d) $-CN$,

(e) $-C(O)NHR_{17}$ wherein R_{17} is loweralkyl,

(f) alkylaminocarbonyl,

(g) dialkylaminocarbonyl,

15 (h) tetrazolyl,

(i) hydroxy,

(j) alkoxy,

(k) sulfonamido,

(l) $-C(O)NHS(O)_2R_{16}$ wherein R_{16} is loweralkyl, haloalkyl, aryl or

dialkylamino,

20 (m) $-S(O)_2NHC(O)R_{16}$ wherein R_{16} is defined as above,








(n)



(o)



(p)



R_1 and R_2 are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl and $(R_{aa})(R_{bb})N-R_{cc}$ wherein R_{aa} is aryl or arylalkyl, R_{bb} is hydrogen or alkanoyl and R_{cc} is alkylene, with the proviso that one or both of R_1 and R_2 is other than hydrogen;

R_3 is $(a)R_4-C(O)-R_5-$, $R_4-C(O)-R_5-N(R_6)-$, wherein R_5 is (i) a covalent bond, (ii) alkylene, (iii) alkenylene, (iv) $-N(R_{20})-R_8-$ or $-R_{8a}-N(R_{20})-R_8-$
















wherein R_8 and R_{8a} are independently selected from the group consisting of alkylene and alkenylene and R_{20} is hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl or cycloalkylalkyl or (v) $-O-R_9-$ or $-R_{9a}-O-R_9-$ wherein R_9 and R_{9a} are independently selected from alkylene;
















R_4 and R_6 are $(R_{11})(R_{12})N-$ wherein R_{11} and R_{12} are independently selected from


(1) hydrogen,

- (2) loweralkyl,
 - (3) haloalkyl,
 - (4) alkoxyalkyl,
 - (5) haloalkoxyalkyl,
 - (6) alkenyl,
 - (7) alkynyl,
 - (8) cycloalkyl,
 - (9) cycloalkylalkyl,
 - (10) aryl,
 - (11) heterocyclic,
 - (12) arylalkyl,
 - (13) (heterocyclic)alkyl,
 - (14) hydroxyalkyl,
 - (15) alkoxy,
 - (16) aminoalkyl,
 - (17) trialkylaminoalkyl,
 - (18) alkylaminoalkyl,
 - (19) dialkylaminoalkyl,
 - (25) carboxyalkyl,
 - (26) (cycloalkyl)aminoalkyl,
 - (27) (cycloalkyl)alkylaminoalkyl,
 - (28) (heterocyclic)aminoalkyl, and
 - (29) (heterocyclic)aminoalkyl, with the proviso that at least one of R_{11} and R_{12} is selected from heterocyclic, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, trialkylaminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, carboxyalkyl, (cycloalkyl)aminoalkyl, (cycloalkyl)alkylaminoalkyl, (heterocyclic)aminoalkyl, and (heterocyclic)alkylaminoalkyl;
- or a pharmaceutically acceptable salt thereof.


95. A compound selected from the group consisting of:








wherein R is selected from the group consisting of:





96. A method for treating hypertension, congestive heart failure, restenosis following arterial injury, renal failure, cancer, colitis, reperfusion injury, angina, pulmonary hypertension, migraine, cerebral or myocardial ischemia, atherosclerosis, coronary angina, cerebral vasospasm, acute and chronic renal failure, gastric ulceration, cyclosporin-induced nephrotoxicity, endotoxin-induced toxicity, asthma, LPL-related lipoprotein disorders, proliferative diseases, acute or chronic pulmonary hypertension, platelet aggregation, thrombosis, IL-2 mediated cardiotoxicity, nociception, colitis, vascular permeability disorders, ischemia-reperfusion injury, Raynaud's disease, prostatic hyperplasia, and migraine comprising a therapeutically effective amount of a compound of claim 94, wherein said compound has an

attached charged functionality which reduces the degree of plasma protein binding of the compound.

5 97. A method of improving the *in vivo* activity of compounds by reducing the amount of compound bound to protein by attaching a charged functionality to the compound.

98. A method of claim 97 wherein the charged functionality carries a positive charge at physiological pH.

10 99. A method for inhibiting bone metastases and metastatic growth in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

15 100. The method of Claim 99 wherein the bone metastases are osteoblastic.

20 101. The method of Claim 100 wherein the osteoblastic bone metastases result from the spread of a primary cancer selected from breast, prostate, lung, kidney, thyroid, myeloma, lymphoma, sarcoma, osteosarcoma, and ovarian.

102. The method of Claim 101 wherein the primary cancer is prostate cancer and the patient is male.

25 103. The method of Claim 99 which additionally comprises co-administration of an anticancer drug.

30 104. The method of Claim 103 wherein the anticancer drug agent is selected from leuprolide, goserelin, bicalutamide, nilutamide, flutamide, vitamin D, vitamin D analogues, estrogen, estrogen analogues, prednisone, hydrocortisone, ketoconazole, cyproterone acetate, and progesterone.

105. The method of Claim 99 which additionally comprises the administration of radiation therapy.

35 106. The method of Claim 99 which additionally comprises the administration of at least one therapeutic agent which impedes net bone loss.

107. The method of Claim 106 wherein the therapeutic agent is a bisphosphonate.

108. The method of Claim 99 wherein the endothelin antagonist is an ET_A-selective endothelin antagonist.

109. A method for the inhibition of bone loss in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

110. The method of Claim 109 wherein the patient has cancer.

111. The method of Claim 109 wherein the cancer is prostate cancer and the patient is male.

112. The method of Claim 109 which additionally comprises the administration of at least one therapeutic agent which impedes net bone loss.

113. The method of Claim 112 wherein the therapeutic agent is a bisphosphonate.

114. A method for the reduction of cancer-related pain in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.


115. The method of Claim 114 wherein the cancer is prostate cancer and the patient is male.

116. The method of Claim 114 which additionally comprises the administration of an anticancer drug.

117. The method of Claim 116 wherein the anticancer drug is selected from leuprolide, goserelin, bicalutamide, nilutamide, flutamide, vitamin D, vitamin D analogues, estrogen, estrogen analogues, prednisone, hydrocortisone, ketoconazole, cyproterone acetate, and progesterone.

118. The method of Claim 115 which additionally comprises the administration of radiation therapy.

119. A method for inhibiting bone metastases in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of a compound of formula I:



wherein

R is $-(CH_2)_m-W$;


Z is selected from $-C(R_{18})(R_{19})-$ and $-C(O)-$;

R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl, and $(R_{aa})(R_{bb})N-R_{cc}-$,

with the proviso that one or both of R₁ and R₂ is other than hydrogen;

R₃ is selected from R₄-C(O)-R₅-, R₄-R_{5a}-, R₄-C(O)-R₅-N(R₆)-, R₆-S(O)₂-R₇- R₂₆-S(O)-R₂₇-, R₂₂-O-C(O)-R₂₃-, loweralkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, aryloxyalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl, alkoxyalkoxyalkyl, and R₁₃-C(O)-CH(R₁₄)-

R₄ and R₆ are independently selected from (R₁₁)(R₁₂)N-, loweralkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkenyl, haloalkoxyalkyl, haloalkoxy, alkoxyhaloalkyl, alkylaminoalkyl, dialkylaminoalkyl, alkoxy, and



R₅ is selected from a covalent bond, alkylene, alkenylene, $-N(R_{20})-R_8-$, $-R_{8a}-N(R_{20})-R_8-$, $-O-R_9-$, and

-R_{9a}-O-R₉-;

R₆ is selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl;

R₇ is a covalent bond, alkylene, alkenylene -N(R₂₁)-R₁₀-, and -R_{10a}-N(R₂₁)-

5 R₁₀-;

R₈ is selected from alkylene and alkenylene;

R₉ is alkylene;

R₁₀ is selected from alkylene and alkenylene;

R₁₁ and R₁₂ are independently selected from hydrogen, loweralkyl, haloalkyl,

10 alkoxyalkyl, haloalkoxyalkylalkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, heterocyclic, arylalkyl, (heterocyclic)alkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, and carboxyalkyl;

R₁₃ is selected from amino, alkylamino and dialkylamino;

R₁₄ is selected from aryl and R₁₅-C(O)-;

15 R₁₅ is selected from amino, alkylamino and dialkylamino;

R₁₆ is selected from loweralkyl, haloalkyl, aryl and dialkylamino;

R₁₇ is loweralkyl;

R₁₈ and R₁₉ are independently selected from hydrogen and loweralkyl;

R₂₀ is selected from hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl,

20 haloalkoxyalkyl, cycloalkyl and cycloalkylalkyl;

R₂₁ is selected from hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl,

haloalkoxyalkyl, aryl and arylalkyl;

R₂₂ is selected from a carboxy protecting group and heterocyclic;

R₂₃ is selected from covalent bond, alkylene, alkenylene and -N(R₂₄)-R₂₅-;

25 R₂₄ is selected from hydrogen and loweralkyl;

R₂₅ is alkylene;

R₂₆ is selected from loweralkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl,

cycloalkylalkyl, aryl, arylalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl and alkoxy-substituted haloalkyl;

30 R₂₇ is selected from alkylene and alkenylene;

R_{5a} is selected from alkylene and alkenylene;

R_{7a} is alkylene;

R_{8a} is selected from alkylene and alkenylene;

R_{9a} is alkylene;

35 R_{10a} is selected from alkylene and alkenylene;

R_{aa} is selected from aryl and arylalkyl;

R_{bb} is selected from hydrogen and alkanoyl;

R_{CC} is alkylene;

m is 0-6;

n is 0 or 1;


z is 0-5;

5 E is selected from hydrogen, loweralkyl and arylalkyl;

G is selected from hydrogen and a carboxy protecting group; and

W is selected from $-C(O)_2-G$; $-PO_3H_2$; $-P(O)(OH)(E)$,

-CN, $-C(O)NHR_{17}$, alkylaminocarbonyl, dialkylaminocarbonyl, tetrazolyl, hydroxy, alkoxy, sulfonamido, $-C(O)NHS(O)_2R_{16}$, $-S(O)_2NHC(O)R_{16}$.



or a pharmaceutically acceptable salt thereof.

120. The method of Claim 119 wherein the bone metastases are
15 osteoblastic.

121. The method of Claim 120 wherein the osteoblastic bone metastases
result from the spread of a primary cancer selected from breast, prostate, lung,
kidney, thyroid, myeloma, lymphoma, sarcoma, osteosarcoma, and ovarian.

122. The method of Claim 121 wherein the primary cancer is prostate cancer
and the patient is male.

123. The method of Claim 119 which additionally comprises the
25 administration of an anticancer drug.


124. The method of Claim 123 wherein the additional anticancer drug is
selected from leuprolide, goserelin, bicalutamide, nilutamide, flutamide, vitamin D,
vitamin D analogues, estrogen, estrogen analogues, prednisone, hydrocortisone,
30 ketoconazole, cyproterone acetate, and progesterone.

125. The method of Claim 119 which additionally comprises the administration of radiation therapy.

126. The method of Claim 119 which additionally comprises the administration of at least one therapeutic agent which impedes net bone loss.

127. The method of Claim 126 wherein the therapeutic agent is a bisphosphonate.

128. A method for the inhibition of bone loss in cancer patients which comprises administering to the patient in need thereof a therapeutically effective amount of a compound of formula I:



wherein

R is $-(CH_2)_m-W$;

Z is selected from $-C(R_{18})(R_{19})-$ and $-C(O)-$;


R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonamidoalkyl, heterocyclic, (heterocyclic)alkyl, and (R_{aa})(R_{bb})N-R_{cc}-;

with the proviso that one or both of R₁ and R₂ is other than hydrogen;

R₃ is selected from R₄-C(O)-R₅-, R₄-R_{5a}-, R₄-C(O)-R₅-N(R₆)-, R₆-S(O)₂-R₇- R₂₆-S(O)-R₂₇-, R₂₂-O-C(O)-R₂₃-, loweralkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, aryloxyalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl, alkoxyalkoxyalkyl, and R₁₃-C(O)-CH(R₁₄)-;

R₄ and R₆ are independently selected from (R₁₁)(R₁₂)N-, loweralkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclic, (heterocyclic)alkyl,

alkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkenyl, haloalkoxyalkyl, haloalkoxy, alkoxyhaloalkyl, alkylaminoalkyl, dialkylaminoalkyl, alkoxy, and



R₅ is selected from a covalent bond, alkylene, alkenylene, -N(R₂₀)-R₈-, -R_{8a}-

- 5 N(R₂₀)-R₈-, -O-R₉-, and
-R_{9a}-O-R₉-;

R₆ is selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl;

R₇ is a covalent bond, alkylene, alkenylene -N(R₂₁)-R₁₀-, and -R_{10a}-N(R₂₁)-

- 10 R₁₀-;

R₈ is selected from alkylene and alkenylene;

R₉ is alkylene;

R₁₀ is selected from alkylene and alkenylene;

R₁₁ and R₁₂ are independently selected from hydrogen, loweralkyl, haloalkyl,

- 15 alkoxyalkyl, haloalkoxyalkylalkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, heterocyclic, arylalkyl, (heterocyclic)alkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, and carboxyalkyl;

R₁₃ is selected from amino, alkylamino and dialkylamino;

R₁₄ is selected from aryl and R₁₅-C(O)-;

- 20 R₁₅ is selected from amino, alkylamino and dialkylamino;

R₁₆ is selected from loweralkyl, haloalkyl, aryl and dialkylamino;

R₁₇ is loweralkyl;

R₁₈ and R₁₉ are independently selected from hydrogen and loweralkyl;

R₂₀ is selected from hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl,

- 25 haloalkoxyalkyl, cycloalkyl and cycloalkylalkyl;

R₂₁ is selected from hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl,

haloalkoxyalkyl, aryl and arylalkyl;

R₂₂ is selected from a carboxy protecting group and heterocyclic;

R₂₃ is selected from covalent bond, alkylene, alkenylene and -N(R₂₄)-R₂₅-;

- 30 R₂₄ is selected from hydrogen and loweralkyl;

R₂₅ is alkylene;

R₂₆ is selected from loweralkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl,

cycloalkylalkyl, aryl, arylalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl and alkoxy-substituted haloalkyl;

- 35 R₂₇ is selected from alkylene and alkenylene;

5

10


15

20

25

131. The method of Claim 130 wherein the therapeutic agent is a bisphosphonate.

132. A method for the reduction of cancer-related pain which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula I:



wherein

R is $-(CH_2)_m-W$;


Z is selected from $-C(R_{18})(R_{19})-$ and $-C(O)-$;

R₁ and R₂ are independently selected from hydrogen, loweralkyl, alkenyl, alkynyl, alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, thioalkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, aminocarbonylalkenyl, alkylaminocarbonylalkenyl, dialkylaminocarbonylalkenyl, hydroxyalkenyl, aryl, arylalkyl, aryloxyalkyl, arylalkoxyalkyl, (N-alkanoyl-N-alkyl)aminoalkyl, alkylsulfonylamidoalkyl, heterocyclic, (heterocyclic)alkyl, and (R_{aa})(R_{bb})N-R_{cc};

with the proviso that one or both of R₁ and R₂ is other than hydrogen;

R₃ is selected from R₄-C(O)-R₅-, R₄-R_{5a}-, R₄-C(O)-R₅-N(R₆)-, R₆-S(O)₂-R₇-R₂₆-S(O)-R₂₇-, R₂₂-O-C(O)-R₂₃-, loweralkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, aryloxyalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl, alkoxyalkoxyalkyl, and R₁₃-C(O)-CH(R₁₄)-

R₄ and R₆ are independently selected from (R₁₁)(R₁₂)N-, loweralkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl, hydroxyalkyl, haloalkyl, haloalkenyl, haloalkoxyalkyl, haloalkoxy, alkoxyhaloalkyl, alkylaminoalkyl, dialkylaminoalkyl, alkoxy, and



R₅ is selected from a covalent bond, alkylene, alkenylene, -N(R₂₀)-R₈-, -R_{8a}-N(R₂₀)-R₈-, -O-R₉-, and -R_{9a}-O-R₉;

R₆ is selected from loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl or arylalkyl;

R₇ is a covalent bond, alkylene, alkenylene -N(R₂₁)-R₁₀-, and -R_{10a}-N(R₂₁)-R₁₀-;

R₈ is selected from alkylene and alkenylene;

R₉ is alkylene;

5 R₁₀ is selected from alkylene and alkenylene;

R₁₁ and R₁₂ are independently selected from hydrogen, loweralkyl, haloalkyl, alkoxyalkyl, haloalkoxyalkylalkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, heterocyclic, arylalkyl, (heterocyclic)alkyl, hydroxyalkyl, alkoxy, aminoalkyl, trialkylaminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, and carboxyalkyl;

10 R₁₃ is selected from amino, alkylamino and dialkylamino;

R₁₄ is selected from aryl and R₁₅-C(O)-;

R₁₅ is selected from amino, alkylamino and dialkylamino;

R₁₆ is selected from loweralkyl, haloalkyl, aryl and dialkylamino;

R₁₇ is loweralkyl;

15 R₁₈ and R₁₉ are independently selected from hydrogen and loweralkyl;

R₂₀ is selected from hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, cycloalkyl and cycloalkylalkyl;

R₂₁ is selected from hydrogen, loweralkyl, alkenyl, haloalkyl, alkoxyalkyl, haloalkoxyalkyl, aryl and arylalkyl;

20 R₂₂ is selected from a carboxy protecting group and heterocyclic;

R₂₃ is selected from covalent bond, alkylene, alkenylene and -N(R₂₄)-R₂₅-;

R₂₄ is selected from hydrogen and loweralkyl;

R₂₅ is alkylene;

R₂₆ is selected from loweralkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl,

25 cycloalkylalkyl, aryl, arylalkyl, heterocyclic, (heterocyclic)alkyl, alkoxyalkyl and alkoxy-substituted haloalkyl;

R₂₇ is selected from alkylene and alkenylene;

R_{5a} is selected from alkylene and alkenylene;

R_{7a} is alkylene;

30 R_{8a} is selected from alkylene and alkenylene;

R_{9a} is alkylene;

R_{10a} is selected from alkylene and alkenylene;

R_{aa} is selected from aryl and arylalkyl;

R_{bb} is selected from hydrogen and alkanoyl;

35 R_{cc} is alkylene;

m is 0-6;

n is 0 or 1;


z is 0-5;

E is selected from hydrogen, loweralkyl and arylalkyl;

G is selected from hydrogen and a carboxy protecting group; and

W is selected from $-\text{C}(\text{O})_2\text{-G}$; $-\text{PO}_3\text{H}_2$, $-\text{P}(\text{O})(\text{OH})(\text{E})$,

- 5 -CN, -C(O)NHR₁₇, alkylaminocarbonyl, dialkylaminocarbonyl, tetrazolyl, hydroxy, alkoxy, sulfonamido, -C(O)NHS(O)₂R₁₆, -S(O)₂NHC(O)R₁₆,




or a pharmaceutically acceptable salt thereof.

133. The method of Claim 132 wherein the cancer is prostate cancer and the patient is male.

134. The method of Claim 132 which additionally comprises the administration of an anticancer drug.

135. The method of Claim 134 wherein the additional anticancer drug is selected from leuprolide, goserelin, bicalutamide, nilutamide, flutamide, vitamin D, vitamin D analogues, estrogen, estrogen analogues, prednisone, hydrocortisone, ketoconazole, cyproterone acetate, and progesterone.

136. A method for inhibiting bone metastases in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of a compound of formula IIIa



IIIa.

137. The method of Claim 136 wherein the bone metastases are osteoblastic.

138. The method of Claim 137 wherein the osteoblastic bone metastases result from the spread of a primary cancer selected from breast, prostate, lung, kidney, thyroid, myeloma, lymphoma, sarcoma, osteosarcoma, and ovarian.

139. The method of Claim 138 wherein the primary cancer is prostate cancer and the patient is male.

140. The method of Claim 138 which additionally comprises the administration of an anticancer drug.

141. The method of Claim 140 wherein the additional anticancer drug is selected from leuprolide, goserelin, bicalutamide, nilutamide, flutamide, vitamin D, vitamin D analogues, estrogen, estrogen analogues, prednisone, hydrocortisone, ketoconazole, cyproterone acetate, and progesterone.


142. The method of Claim 138 which additionally comprises the administration of radiation therapy.

143. The method of Claim 138 which additionally comprises the administration of at least one therapeutic agent which impedes net bone loss.

144. The method of Claim 143 wherein the agent is a bisphosphonate.

145. The method of Claim 138 wherein the endothelin antagonist is an ET_A-selective endothelin antagonist.

- 5 146. A method for the inhibition of bone loss in cancer patients which comprises administering to the patient in need thereof a therapeutically effective amount of a compound of formula IIIa




IIIa.

10 147. The method of Claim 146 wherein the cancer is prostate cancer and the patient is male.

15 148. The method of Claim 146 which additionally comprises the administration of at least one therapeutic agent which impedes net bone loss.

149. The method of Claim 148 wherein therapeutic agent is a bisphosphonate.

20 150. A method for the reduction of cancer-related pain which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula IIIa



IIIa.

151. The method of Claim 150 wherein the cancer is prostate cancer and the patient is male.

152. The method of Claim 150 which additionally comprises the administration of an anticancer drug.

153. The method of Claim 152 wherein the anticancer drug is selected from leuprolide, goserelin, bicalutamide, nilutamide, flutamide, vitamin D, vitamin D analogues, estrogen, estrogen analogues, prednisone, hydrocortisone, ketoconazole, cyproterone acetate, and progesterone.

154. A method for preventing new bone metastases in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

155. A method for inhibiting metastatic growth in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

156. A method for inhibiting bone turnover in a patient which comprises administering to the patient in need thereof a therapeutically effective amount of an endothelin ET-A receptor antagonist.

157. A compound according to Claim 1 of formula (I) wherein n is zero; Z is -CH₂- wherein R₁₈ and R₁₉ are hydrogen; R is C(O)-G wherein G is hydrogen; R₁ is aryl substituted with one substituent selected from methoxy, methoxyethoxy, and isopropoxyethoxy; R₂ is 1,3-benzodiox-5-yl; R₃ is R₄-C(O)-R₅- wherein R₅ is methylene and R₄ is selected from (R₁₁)(R₁₂)N- and (R_{11a})(R_{12a})N-N(H)-; one of R₁₁ and R₁₂ is hydrogen and the other is selected from arylalkyl and diarylalkyl wherein each aryl group of the diarylalkyl is substituted with methyl or ethyl; and one of R_{11a} or R_{12a} is alkyl and the other is aryl.


158. A compound according to Claim 1 of formula (I) wherein n is zero; Z is -CH₂- wherein R₁₈ and R₁₉ are hydrogen; R is C(O)-G wherein G is hydrogen; R₁ is phenyl substituted with one substituent selected from methoxy, methoxyethoxy, and isopropoxyethoxy; R₂ is 1,3-benzodiox-5-yl; R₃ is R₄-C(O)-R₅- wherein R₅ is methylene and R₄ is selected from (R₁₁)(R₁₂)N- and (R_{11a})(R_{12a})N-N(H)-; one of R₁₁ and R₁₂ is hydrogen and the other is selected from phenylalkyl and diphenylalkyl wherein each phenyl group of the diphenylalkyl is substituted with methyl or ethyl; and one of R_{11a} or R_{12a} is alkyl and the other is phenyl.

159. A compound according to Claim 1 of formula (II) wherein n is zero; Z is -CH₂- wherein R₁₈ and R₁₉ are hydrogen; R is C(O)-G wherein G is hydrogen; R₁ is aryl substituted with one substituent selected from methoxy, methoxyethoxy, and isopropoxyethoxy; R₂ is 1,3-benzodiox-5-yl; R₃ is R₄-C(O)-R₅- wherein R₅ is methylene and R₄ is selected from (R₁₁)(R₁₂)N- and (R_{11a})(R_{12a})N-N(H)-; one of R₁₁ and R₁₂ is hydrogen and the other is selected from arylalkyl and diarylalkyl wherein each aryl group of the diarylalkyl is substituted with methyl or ethyl; and one of R_{11a} or R_{12a} is alkyl and the other is aryl.

160. A compound according to Claim 1 of formula (II) wherein n is zero; Z is -CH₂- wherein R₁₈ and R₁₉ are hydrogen; R is C(O)-G wherein G is hydrogen; R₁ is phenyl substituted with one substituent selected from methoxy, methoxyethoxy, and isopropoxyethoxy; R₂ is 1,3-benzodiox-5-yl; R₃ is R₄-C(O)-R₅- wherein R₅ is methylene and R₄ is selected from (R₁₁)(R₁₂)N- and (R_{11a})(R_{12a})N-N(H)-; one of R₁₁ and R₁₂ is hydrogen and the other is selected from phenylalkyl and diphenylalkyl wherein each phenyl group of the diphenylalkyl is substituted with methyl or ethyl; and one of R_{11a} or R_{12a} is alkyl and the other is phenyl.

ABSTRACT OF THE DISCLOSURE

A compound of the formula (I):



or a pharmaceutically acceptable salt thereof is disclosed, as well as processes for and intermediates in the preparation thereof, a method of antagonizing endothelin. methods for the inhibition of bone metastases, methods for the prevention of growth of new metastases, methods for the inhibition of bone turnover, and methods for the prevention of bone loss in patients, including cancer patients, using an endothelin ET-A receptor antagonist.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: M. Winn, *et al.*

Serial No.:

Filed:

For: Endothelin Antagonists

Group Art Unit:

Examiner:

Case No.: 5994.US.P9

Express Mail No.:

Certificate of Mailing (37 CFR §1.10)

I hereby certify that this paper (along with any paper referred to as being attached or enclosed) is being deposited with the United States Postal Service on the date shown below with sufficient postage as Express Mail in an envelope addressed to:

Commissioner for Patents
Box Patent Application
Washington, D.C. 20231, on:

Date of Deposit:

Deborah L. Siedlecki

Date

**Declaration and Power of Attorney
for a United States Patent Application**

As a below-named inventor, I hereby declare:

My residence, post office address and citizenship are as stated below next to my name. I believe I am an original and first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled "METHODS OF TREATING CANCER AND THE PAIN ASSOCIATED THEREWITH USING ENDOTHELIN ANTAGONISTS", the specification of which is attached.

I hereby state that I have reviewed and understand the contents of the above-mentioned specification, including the claims.

I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

Claim to benefit of foreign application(s) as follows:

I hereby claim foreign priority benefits under 35 U.S.C. §119 for the following foreign applications for patent or inventor's certificate.

NONE

The following foreign applications for patent or inventor's certificate have a filing date earlier than the filing date of the applications identified above.

NONE

Claim to benefit of earlier U.S. application(s) as follows:

I hereby claim benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

NONE

I hereby claim the benefit under 35 U.S.C. §120 of the following earlier-filed United States patent applications. Insofar as the subject matter of each of the claims of this application is not disclosed in the prior U.S. applications in the manner required by 35 U.S.C. §112, first paragraph, I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which came into existence between the filing date(s) of the prior applications and the national or PCT filing date of this application.

U.S. Serial No. 09/634,661, filed August 7, 2000
U.S. Serial No. 09/048,955, filed March 27, 1998
U.S. Serial No. 08/794,506, filed February 4, 1997
U.S. Serial No. 08/600,625, filed February 13, 1996
U.S. Serial No. 08/497,998, filed August 2, 1995
U.S. Serial No. 08/442,575, filed May 30, 1995
U.S. Serial No. 08/334,717, filed November 4, 1994
U.S. Serial No. 08/293,349, filed August 19, 1994

I hereby appoint the following Attorneys and/or agents to prosecute this application and any continuation or divisional applications based hereon, and to transact all business in the Patent and Trademark Office connected therewith:

Regina M. Anderson, Reg. No. 35,820
Cheryl L. Becker, Reg. No. 35,441
Thomas D. Brainard, Reg. No. 32,459
Valerie Calloway, Reg. No. 40,546
Dianne Casuto, Reg. No. 40,943
Steven R. Crowley, Reg. No. 31,604
Andreas M. Danckers, Reg. No. 32,652
J. Michael Dixon, Reg. No. 32,410
Mimi C. Goller, Reg. No. 39,046
Daniel J. Hulseberg, Reg. No. 36,554

Neal D. Marcus, Reg. No. 35,267
James D. McNeil, Reg. No. 26,204
Lawrence S. Pope, Reg. No. 26,791
Nicholas A. Poulos, Reg. No. 30,209
Dugal S. Sickert, Reg. No. 33,784
Gregory W. Steele, Reg. No. 33,796
Michael J. Ward, Reg. No. 37,960
David M. Weinstein, Reg. No. 28,128
Steven F. Weinstock, Reg. No. 30,117
Brian R. Woodworth, Reg. No. 33,137
Paul D. Yasger, Reg. No. 37,477

Send correspondence to: Steven F. Weinstock
Abbott Laboratories
D-377 AP6D
100 Abbott Park Road
Abbott Park, IL 60064-6050

Direct telephone calls to: James D. McNeil
(847) 937-4556

Name: (first, middle, last): MARTIN WINN
Post Office Address: 1263 Carlisle Place, Deerfield, Illinois 60015
Residence: Deerfield, Illinois 60015
Citizenship: United States of America

Name: (first, middle, last):	STEVEN A. BOYD
Post Office Address:	410 Lakeview Drive, Mundelein, Illinois 60060
Residence:	Mundelein, Illinois 60060
Citizenship:	United States of America
Name: (first, middle, last):	CHARLES W. HUTCHINS
Post Office Address:	4120 Oglesby, Gurnee, Illinois 60031
Residence:	Gurnee, IL 60031
Citizenship:	United States of America
Name: (first, middle, last):	HWAN-SOO JAE
Post Office Address:	477 Woodland Avenue, Glencoe, Illinois 60022
Residence:	Glencoe, Illinois 60022
Citizenship:	United States of America
Name: (first, middle, last):	ANDREW S. TASKER
Post Office Address:	6251 Eagle Ridge Drive, Gurnee, Illinois 60031
Residence:	Gurnee, IL 60031
Citizenship:	British
Name: (first, middle, last):	THOMAS W. von GELDERN
Post Office Address:	4209 W. Solon Road, Richmond, Illinois 60071
Residence:	Richmond, Illinois 60071
Citizenship:	United States of America
Name: (first, middle, last):	JEFFREY A. KESTER
Post Office Address:	680 Sapling, Deerfield, Illinois 60015
Residence:	Deerfield, Illinois 60015
Citizenship:	United States of America
Name: (first, middle, last):	BRYAN K. SORENSEN
Post Office Address:	2620 North Lewis Ave., Waukegan, Illinois 60087
Residence:	Waukegan, Illinois 60076
Citizenship:	United States of America
Name: (first, middle, last):	BRUCE G. SZCZEPANKIEWICZ
Post Office Address:	33720 Royal Oak Ln., #209, Gages Lake, IL 60030
Residence:	Gages Lake, Illinois 60030
Citizenship:	United States of America
Name: (first, middle, last):	KENNETH J. HENRY
Post Office Address:	1047 S. West Avenue, Waukegan, Illinois 60085
Residence:	Waukegan, Illinois 60085
Citizenship:	United States of America
Name: (first, middle, last):	GANG LIU
Post Office Address:	520 Lakehurst Drive, Apt. 2L, Waukegan, IL 60085
Residence:	Waukegan, Illinois 60085
Citizenship:	People's Republic of China
Name: (first, middle, last):	STEVEN J. WITTENBERGER
Post Office Address:	317 N. Emerald Avenue, Mundelein, Illinois 60060
Residence:	Mundelein, Illinois 60060
Citizenship:	United States of America

Name: (first, middle, last): STEVEN A. KING
Post Office Address: 16713 Orchard Valley Drive, Gurnee, Illinois 60031
Residence: Gurnee, IL 60031
Citizenship: United States of America

Name: (first, middle, last): Todd J. Janus
Post Office Address: 270 Big Terra Lane, Gurnee, IL 60031
Residence: Gurnee, IL 60031
Citizenship: United States of America

Name: (first, middle, last): Robert J. Padley
Post Office Address: 770 Moffett Road, Lake Bluff, IL 60044
Residence: Lake Bluff, IL 60044
Citizenship: United States of America

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that all statements made herein were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

MARTIN WINN _____ Date STEVEN A. BOYD _____ Date

CHARLES W. HUTCHINS _____ Date HWAN-SOO JAE _____ Date

ANDREW S. TASKER _____ Date THOMAS W. von GELDERN _____ Date

JEFFREY A. KESTER _____ Date BRYAN K. SORENSEN _____ Date

BRUCE G. SZCZEPANKIEWICZ _____ Date KENNETH J. HENRY _____ Date

GANG LIU _____ Date STEVEN WITTENBERGER _____ Date

STEVEN A. KING _____ Date TODD J. JANUS _____ Date

ROBERT J. PADLEY _____ Date